

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

B9

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C12N 15/12, 15/62, 5/10, C07K 14/52, 16/22, C12Q 1/68, A61K 38/18</b>		<b>A1</b>	(11) International Publication Number: <b>WO 00/66736</b> (43) International Publication Date: <b>9 November 2000 (09.11.00)</b>
(21) International Application Number: <b>PCT/US00/40047</b> (22) International Filing Date: <b>3 May 2000 (03.05.00)</b>  (30) Priority Data: 09/304,216      3 May 1999 (03.05.99)      US 60/164,463      10 November 1999 (10.11.99)      US 60/180,169      4 February 2000 (04.02.00)      US  (71) Applicant: <b>ZYMOGENETICS, INC. [US/US]; 1201 Eastlake Avenue East, Seattle, WA 98102 (US).</b>  (72) Inventors: <b>GILBERT, Teresa; 246 12th Street Southeast, Seattle, WA 98002 (US). HART, Charles, E.; 20114 183rd Place Northeast, Woodinville, WA 98072 (US). SHEPPARD, Paul, O.; 13532 278th Drive Northeast, Granite Falls, WA 98252 (US). GILBERTSON, Debra, G.; 4025 Northeast 195th Place, Seattle, WA 98155 (US).</b>  (74) Agent: <b>PARKER, Gary, E.; ZymoGenetics, Inc., 1201 Eastlake Avenue East, Seattle, WA 98102 (US).</b>		(81) Designated States: <b>AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</b>  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: <b>GROWTH FACTOR HOMOLOG ZVEGF4</b>			
(57) Abstract			
<p>Polypeptide growth factors, methods of making them, polynucleotides encoding them, antibodies to them, and methods of using them are disclosed. The polypeptides comprise an amino acid segment that is at least 70 % identical to residues 52-179 of SEQ ID NO:2 or residues 258-370 of SEQ ID NO:2. Multimers of the polypeptides are also disclosed. The polypeptides, multimeric proteins, and polynucleotides can be used in the study and regulation of cell and tissue development, as components of cell culture media, and as diagnostic agents.</p>			

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Description

## GROWTH FACTOR HOMOLOG ZVEGF4

5

## BACKGROUND OF THE INVENTION

In multicellular animals, cell growth, differentiation, and migration are controlled by polypeptide growth factors. These growth factors play a role in both normal development and pathogenesis, including the development of solid tumors.

10

Polypeptide growth factors influence cellular events by binding to cell-surface receptors, many of which are tyrosine kinases. Binding initiates a chain of signalling events within the cell, which ultimately results in phenotypic changes, such as cell division, protease production, and cell migration.

15

Growth factors can be classified into families on the basis of structural similarities. One such family, the PDGF (platelet derived growth factor) family, is characterized by a dimeric structure stabilized by disulfide bonds. This family includes PDGF, the placental growth factors (PIGFs), and the vascular endothelial growth factors (VEGFs). The individual polypeptide chains of these proteins form characteristic higher-order structures having a bow tie-like configuration about a cystine knot, formed by disulfide bonding between pairs of cysteine residues. Hydrophobic interactions between loops contribute to the dimerization of the two monomers. See, Daopin et al., *Science* 257:369, 1992; Lapthorn et al., *Nature* 369:455, 1994. Members of this family are active as both homodimers and heterodimers. See, for example, Heldin et al., *EMBO J.* 7:1387-1393, 1988; Cao et al., *J. Biol. Chem.* 271:3154-3162, 1996. The cystine knot motif and "bow tie" fold are also characteristic of the growth factors transforming growth factor-beta (TGF- $\beta$ ) and nerve growth factor (NGF), and the glycoprotein hormones. Although their amino acid sequences are quite divergent, these proteins all contain the six conserved cysteine residues of the cystine knot.

20

25

30

Five vascular endothelial growth factors have been identified: VEGF, also known as vascular permeability factor (Dvorak et al., *Am. J. Pathol.* 146:1029-1039, 1995); VEGF-B (Olofsson et al., *Proc. Natl. Acad. Sci. USA* 93:2567-2581, 1996; Hayward et al., WIPO Publication WO 96/27007); VEGF-C (Joukov et al., *EMBO J.* 15:290-298, 1996); VEGF-D (Oliviero, WO 97/12972; Achen et al., WO 98/07832), and zveg3 (SEQ ID NO:32 and NO:33; co-pending U.S. Patent Applications Nos. 60/111,173, 60/142,576, and 60/161,653). Five VEGF polypeptides

35

(121, 145, 165, 189, and 206 amino acids) arise from alternative splicing of the VEGF mRNA.

VEGFs stimulate the development of vasculature through a process known as angiogenesis, wherein vascular endothelial cells re-enter the cell cycle, 5 degrade underlying basement membrane, and migrate to form new capillary sprouts. These cells then differentiate, and mature vessels are formed. This process of growth and differentiation is regulated by a balance of pro-angiogenic and anti-angiogenic factors. Angiogenesis is central to normal formation and repair of tissue, occurring in embryo development and wound healing. Angiogenesis is also a factor in the 10 development of certain diseases, including solid tumors, rheumatoid arthritis, diabetic retinopathy, macular degeneration, and atherosclerosis.

A number of proteins from vertebrates and invertebrates have been identified as influencing neural development. Among those molecules are members of the neuropilin family and the semaphorin/collapsin family.

15 Three receptors for VEGF have been identified: KDR/Flk-1 (Matthews et al., *Proc. Natl. Acad. Sci. USA* 88:9026-9030, 1991), Flt-1 (de Vries et al., *Science* 255:989-991, 1992), and neuropilin-1 (Soker et al., *Cell* 92:735-745, 1998). Neuropilin-1 is also a receptor for PlGF-2 (Migdal et al., *J. Biol. Chem.* 273: 22272-22278, 1998).

20 Neuropilin-1 is a cell-surface glycoprotein that was initially identified in *Xenopus* tadpole nervous tissues, then in chicken, mouse, and human. The primary structure of neuropilin-1 is highly conserved among these vertebrate species. Neuropilin-1 has been demonstrated to be a receptor for various members of the semaphorin family including semaphorin III (Kolodkin et al., *Cell* 90:753-762, 1997), 25 Sema E and Sema IV (Chen et al., *Neuron* 19:547-559, 1997). A variety of activities have been associated with the binding of neuropilin-1 to its ligands. For example, binding of semaphorin III to neuropilin-1 can induce neuronal growth cone collapse and repulsion of neurites *in vitro* (Kitsukawa et al., *Neuron* 19: 995-1005, 1997). Experiments with transgenic mice indicate the involvement of neuropilin-1 in the 30 development of the cardiovascular system, nervous system, and limbs. See, for example, Kitsukawa et al., *Development* 121:4309-4318, 1995; and Takashima et al., American Heart Association 1998 Meeting, Abstract # 3178.

Semaphorins are a large family of molecules which share the defining semaphorin domain of approximately 500 amino acids. Dimerization is believed to be 35 important for functional activity (Klostermann et al., *J. Biol. Chem.* 273:7326-7331, 1998). Collapsin-1, the first identified vertebrate member of the semaphorin family of axon guidance proteins, has also been shown to form covalent dimers, with

dimerization necessary for collapse activity (Koppel et al., *J. Biol. Chem.* 273:15708-15713, 1998). Semaphorin III has been associated *in vitro* with regulating growth cone collapse and chemorepulsion of neurites. Semaphorins have been shown to be responsible for a variety of developmental effects, including effects on sensory afferent innervation, skeletal and cardiac development (Fehar et al., *Nature* 383:525-528, 1996), immunosuppression via inhibition of cytokines (Mangasser-Stephan et al., *Biochem. Biophys. Res. Comm.* 234:153-156, 1997), and promotion of B-cell aggregation and differentiation (Hall et al., *Proc. Natl. Acad. Sci. USA* 93:11780-11785, 1996). CD100 has also been shown to be expressed in many T-cell lymphomas and may be a marker of malignant T-cell neoplasms (Dorfman et al., *Am. J. Pathol.* 153:255-262, 1998). Transcription of the mouse semaphorin gene, M-semaH, correlates with metastatic ability of mouse tumor cell lines (Christensen et al., *Cancer Res.* 58:1238-1244, 1998).

The role of growth factors, other regulatory molecules, and their receptors in controlling cellular processes makes them likely candidates and targets for therapeutic intervention. Platelet-derived growth factor, for example, has been disclosed for the treatment of periodontal disease (U.S. Patent No. 5,124,316), gastrointestinal ulcers (U.S. Patent No. 5,234,908), and dermal ulcers (Robson et al., *Lancet* 339:23-25, 1992). Inhibition of PDGF receptor activity has been shown to reduce intimal hyperplasia in injured baboon arteries (Giese et al., Restenosis Summit VIII, Poster Session #23, 1996; U.S. Patent No. 5,620,687). PDGF has also been shown to stimulate bone cell replication (reviewed by Canalis et al., *Endocrinology and Metabolism Clinics of North America* 18:903-918, 1989), to stimulate the production of collagen by bone cells (Centrella et al., *Endocrinology* 125:13-19, 1989) and to be useful in regenerating periodontal tissue (U.S. Patent No. 5,124,316; Lynch et al., *J. Clin. Periodontol.* 16:545-548, 1989). Vascular endothelial growth factors (VEGFs) have been shown to promote the growth of blood vessels in ischemic limbs (Isner et al., *The Lancet* 348:370-374, 1996), and have been proposed for use as wound-healing agents, for treatment of periodontal disease, for promoting endothelialization in vascular graft surgery, and for promoting collateral circulation following myocardial infarction (WIPO Publication No. WO 95/24473; U.S. Patent No. 5,219,739). VEGFs are also useful for promoting the growth of vascular endothelial cells in culture. A soluble VEGF receptor (soluble flt-1) has been found to block binding of VEGF to cell-surface receptors and to inhibit the growth of vascular tissue *in vitro* (*Biotechnology News* 16(17):5-6, 1996).

In view of the proven clinical utility of polypeptide growth factors, there is a need in the art for additional such molecules for use as therapeutic agents, diagnostic agents, and research tools and reagents.

The present invention provides such polypeptides for these and other uses that should be apparent to those skilled in the art from the teachings herein.

#### DESCRIPTION OF THE INVENTION

5           The present invention provides an isolated polypeptide of at least 15 amino acid residues comprising an epitope-bearing portion of a protein of SEQ ID NO:2. Within certain embodiments, the polypeptide comprises a segment that is at least 70% identical to residues 52-179 of SEQ ID NO:2 or residues 258-370 of SEQ ID NO:2. Within other embodiments, the polypeptide is selected from the group  
10           consisting of residues 19-179 of SEQ ID NO:2, residues 52-179 of SEQ ID NO:2, residues 19-257 of SEQ ID NO:2, residues 52-257 of SEQ ID NO:2, residues 19-253 of SEQ ID NO:2, residues 52-253 of SEQ ID NO:2, residues 19-370 of SEQ ID NO:2, residues 52-370 of SEQ ID NO:2, residues 258-370 of SEQ ID NO:2, and residues 180-370 of SEQ ID NO:2.

15           The invention also provides an isolated polypeptide comprising a sequence of amino acids of the formula  $R1_x-R2_y-R3_z$ , wherein R1 is a polypeptide of from 100 to 130 residues in length, is at least 70% identical to residues 52-179 of SEQ ID NO:2, and comprises a sequence motif C[KR]Y[DNE][WYF]X{11,15}G[KR][WYF]C (SEQ ID NO:4) corresponding to  
20           residues 109-131 of SEQ ID NO:2; R2 is a polypeptide at least 90% identical to residues 180-257 of SEQ ID NO:2; R3 is a polypeptide at least 70% identical in amino acid sequence to residues 258-370 of SEQ ID NO:2 and comprises cysteine residues at positions corresponding to residues 272, 302, 306, 318, 360, and 362 of SEQ ID NO:2, a glycine residue at a position corresponding to residue 304 of SEQ ID NO:2, and a  
25           sequence motif CX{18,33}CXGXCX{6,33}CX{20,50}CXC (SEQ ID NO:3) corresponding to residues 272-362 of SEQ ID NO:2; and each of x, y, and z is individually 0 or 1, subject to the limitations that at least one of x and z is 1, and, if x and z are each 1, then y is 1. There are thus provided isolated polypeptides of the above formula wherein (a) x=1, (b) z=1, and (c) x=1 and z=1. Within certain embodiments,  
30           x=1 and R1 is at least 90% identical to residues 52-179 of SEQ ID NO:2 or residues 19-179 of SEQ ID NO:2. Within related embodiments, x=1 and R1 comprises residues 52-179 of SEQ ID NO:2. Within other embodiments, z=1 and R3 is at least 90% identical to residues 258-370 of SEQ ID NO:2 or R3 comprises residues 258-370 of SEQ ID NO:2. Within other embodiments, x=1, z=1, and R3 is at least 90% identical to  
35           residues 258-370 of SEQ ID NO:2; and x=1, z=1, R1 is at least 90% identical to residues 52-179 of SEQ ID NO:2, and R2 is at least 90% identical to residues 180-257 of SEQ ID NO:2. Within additional embodiments, x=1, z=1, and the polypeptide

comprises residues 19-370 of SEQ ID NO:2 or residues 52-370 of SEQ ID NO:2. The isolated polypeptide may further comprise cysteine residues at positions corresponding to residues 308 and 316 of SEQ ID NO:2. Within other embodiments, the isolated polypeptide further comprises an affinity tag. Within a related embodiment, the isolated polypeptide comprises an immunoglobulin constant domain.

The present invention also provides an isolated protein comprising a first polypeptide operably linked to a second polypeptide, wherein the first polypeptide comprises a sequence of amino acids of the formula  $R1_x-R2_y-R3_z$  as disclosed above. The protein modulates cell proliferation, apoptosis, differentiation, metabolism, or migration. Within one embodiment, the protein is a heterodimer. Within related embodiments, the second polypeptide is selected from the group consisting of VEGF, VEGF-B, VEGF-C, VEGF-D, zveg3 (SEQ ID NO:33), PlGF, PDGF-A, and PDGF-B. Within other related embodiments,  $x=1$ ,  $z=1$ , and the second polypeptide comprises residues 46-345 of SEQ ID NO:33;  $x=1$  and the second polypeptide comprises residues 46-170 of SEQ ID NO:33; or  $z=1$  and the second polypeptide comprises residues 235-345 of SEQ ID NO:33.

Within another embodiment, the protein is a homodimer.

There is also provided an isolated protein produced by a method comprising the steps of (a) culturing a host cell containing an expression vector comprising the following operably linked elements: a transcription promoter; a DNA segment encoding a polypeptide selected from the group consisting of (i) residues 52-370 of SEQ ID NO:2, (ii) residues 52-253 of SEQ ID NO:2, (iii) residues 180-370 of SEQ ID NO:2, and (iv) residues 258-370 of SEQ ID NO:2; and a transcription terminator, under conditions whereby the DNA segment is expressed; and (b) recovering from the cell the protein product of expression of the DNA construct.

Within another aspect of the invention there is provided an isolated polynucleotide of up to approximately 4.4 kb in length, wherein said polynucleotide encodes a polypeptide as disclosed above. Within one embodiment of the invention, the polynucleotide is DNA.

Within a further aspect of the invention there is provided an expression vector comprising the following operably linked elements: (a) a transcription promoter; (b) a DNA polynucleotide as disclosed above; and (c) a transcription terminator. The vector may further comprise a secretory signal sequence operably linked to the DNA polynucleotide.

Also provided by the invention is a cultured cell into which has been introduced an expression vector as disclosed above, wherein the cell expresses the polypeptide encoded by the DNA polynucleotide. The cultured cell can be used within

a method of producing a polypeptide, the method comprising culturing the cell and recovering the expressed polypeptide.

The proteins provided herein can be combined with a pharmaceutically acceptable vehicle to provide a pharmaceutical composition.

5           The invention also provides an antibody that specifically binds to an epitope of a polypeptide as disclosed above. Antibodies of the invention include, *inter alia*, monoclonal antibodies and single chain antibodies, and may be linked to a reporter molecule.

10           The invention further provides a method for detecting a genetic abnormality in a patient, comprising the steps of (a) obtaining a genetic sample from a patient, (b) incubating the genetic sample with a polynucleotide comprising at least 14 contiguous nucleotides of SEQ ID NO:1 or the complement of SEQ ID NO:1, under conditions wherein the polynucleotide will hybridize to a complementary polynucleotide sequence, to produce a first reaction product, and (c) comparing the first  
15           reaction product to a control reaction product, wherein a difference between the first reaction product and the control reaction product is indicative of a genetic abnormality in the patient.

          The invention also provides a polypeptide comprising a sequence selected from the group consisting of: residues 46-234 of SEQ ID NO:33 operably  
20           linked to residues 258-370 of SEQ ID NO:2; residues 46-170 of SEQ ID NO:33 operably linked to residues 180-370 of SEQ ID NO:2; residues 52-257 of SEQ ID NO:2 operably linked to residues 235-345 of SEQ ID NO:33; and residues 52-179 of SEQ ID NO:2 operably linked to residues 171-345 of SEQ ID NO:33.

          The invention also provides a method of activating a cell-surface PDGF  
25           receptor, comprising exposing a cell comprising a cell-surface PDGF receptor to a polypeptide or protein as disclosed above, whereby the polypeptide or protein binds to and activates the receptor.

          The invention also provides a method of inhibiting a PDGF receptor-mediated cellular process, comprising exposing a cell comprising a cell-surface PDGF  
30           receptor to a compound that inhibits binding of a polypeptide or protein as disclosed above to the receptor.

          The invention also provides a method of stimulating the growth of bone tissue, comprising applying to bone a growth-stimulating amount of a polypeptide or protein as disclosed above.

35           The invention also provides a method of modulating the proliferation, differentiation, migration, or metabolism of bone cells, comprising exposing bone cells to an effective amount of a polypeptide or protein as disclosed above.



These and other aspects of the invention will become evident upon reference to the following detailed description of the invention and the attached drawings.

In the accompanying drawings:

5           Fig. 1 is a Hopp/Woods hydrophilicity profile of the amino acid sequence shown in SEQ ID NO:2. The profile is based on a sliding six-residue window. Buried G, S, and T residues and exposed H, Y, and W residues were ignored. These residues are indicated in the figure by lower case letters.

10           Fig. 2 is an illustration of the vector pHB12-8 for use in expressing cDNAs in transgenic animals.

          The term "affinity tag" is used herein to denote a polypeptide segment that can be attached to a second polypeptide to provide for purification or detection of the second polypeptide or provide sites for attachment of the second polypeptide to a substrate. In principal, any peptide or protein for which an antibody or other specific  
15   binding agent is available can be used as an affinity tag. Affinity tags include a poly-histidine tract, protein A (Nilsson et al., *EMBO J.* 4:1075, 1985; Nilsson et al., *Methods Enzymol.* 198:3, 1991), glutathione S transferase (Smith and Johnson, *Gene* 67:31, 1988), maltose binding protein (Kellerman and Ferenci, *Methods Enzymol.* 90:459-463, 1982; Guan et al., *Gene* 67:21-30, 1987), Glu-Glu affinity tag (Grussenmeyer et al.,  
20   *Proc. Natl. Acad. Sci. USA* 82:7952-4, 1985; see SEQ ID NO:5), substance P, Flag™ peptide (Hopp et al., *Biotechnology* 6:1204-10, 1988), streptavidin binding peptide, thioredoxin, ubiquitin, cellulose binding protein, T7 polymerase, or other antigenic epitope or binding domain. See, in general, Ford et al., Protein Expression and Purification 2: 95-107, 1991. DNAs encoding affinity tags and other reagents are  
25   available from commercial suppliers (e.g., Pharmacia Biotech, Piscataway, NJ; New England Biolabs, Beverly, MA; and Eastman Kodak, New Haven, CT).

          The term "allelic variant" is used herein to denote any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in phenotypic polymorphism within  
30   populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequence. The term allelic variant is also used herein to denote a protein encoded by an allelic variant of a gene.

          The terms "amino-terminal" and "carboxyl-terminal" are used herein to denote positions within polypeptides. Where the context allows, these terms are used  
35   with reference to a particular sequence or portion of a polypeptide to denote proximity or relative position. For example, a certain sequence positioned carboxyl-terminal to a reference sequence within a polypeptide is located proximal to the carboxyl terminus of

the reference sequence, but is not necessarily at the carboxyl terminus of the complete polypeptide.

A "beta-strand-like region" is a region of a protein characterized by certain combinations of the polypeptide backbone dihedral angles phi ( $\phi$ ) and psi ( $\psi$ ). Regions wherein  $\phi$  is less than  $-60^\circ$  and  $\psi$  is greater than  $90^\circ$  are beta-strand-like. Those skilled in the art will recognize that the limits of a  $\beta$ -strand are somewhat imprecise and may vary with the criteria used to define them. See, for example, Richardson and Richardson in Fasman, ed., *Prediction of Protein Structure and the Principles of Protein Conformation*, Plenum Press, New York, 1989; and Lesk, Protein Architecture: A Practical Approach, Oxford University Press, New York, 1991.

A "complement" of a polynucleotide molecule is a polynucleotide molecule having a complementary base sequence and reverse orientation as compared to a reference sequence. For example, the sequence 5' ATGCACGGG 3' is complementary to 5' CCCGTGCAT 3'.

"Corresponding to", when used in reference to a nucleotide or amino acid sequence, indicates the position in a second sequence that aligns with the reference position when two sequences are optimally aligned.

The term "degenerate nucleotide sequence" denotes a sequence of nucleotides that includes one or more degenerate codons (as compared to a reference polynucleotide molecule that encodes a polypeptide). Degenerate codons contain different triplets of nucleotides, but encode the same amino acid residue (i.e., GAU and GAC triplets each encode Asp).

The term "expression vector" is used to denote a DNA molecule, linear or circular, that comprises a segment encoding a polypeptide of interest operably linked to additional segments that provide for its transcription. Such additional segments include promoter and terminator sequences, and may also include one or more origins of replication, one or more selectable markers, an enhancer, a polyadenylation signal, etc. Expression vectors are generally derived from plasmid or viral DNA, or may contain elements of both.

The term "isolated", when applied to a polynucleotide, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such isolated molecules are those that are separated from their natural environment and include cDNA and genomic clones. Isolated polynucleotide molecules of the present invention are free of other genes with which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as promoters and terminators. The identification of

associated regions will be evident to one of ordinary skill in the art (see, for example, Dynan and Tijan, *Nature* 316:774-78, 1985).

An "isolated" polypeptide or protein is a polypeptide or protein that is found in a condition other than its native environment, such as apart from blood and animal tissue. Within one embodiment, the isolated polypeptide or protein is substantially free of other polypeptides or proteins, particularly other polypeptides or proteins of animal origin. The polypeptides or proteins may be provided in a highly purified form, i.e. greater than 95% pure or greater than 99% pure. When used in this context, the term "isolated" does not exclude the presence of the same polypeptide or protein in alternative physical forms, such as dimers or alternatively glycosylated or derivatized forms.

A "motif" is a series of amino acid positions in a protein sequence for which certain amino acid residues are required. A motif defines the set of possible residues at each such position.

"Operably linked" means that two or more entities are joined together such that they function in concert for their intended purposes. When referring to DNA segments, the phrase indicates, for example, that coding sequences are joined in the correct reading frame, and transcription initiates in the promoter and proceeds through the coding segment(s) to the terminator. When referring to polypeptides, "operably linked" includes both covalently (e.g., by disulfide bonding) and non-covalently (e.g., by hydrogen bonding, hydrophobic interactions, or salt-bridge interactions) linked sequences, wherein the desired function(s) of the sequences are retained.

The term "ortholog" denotes a polypeptide or protein obtained from one species that is the functional counterpart of a polypeptide or protein from a different species. Sequence differences among orthologs are the result of speciation.

A "polynucleotide" is a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Polynucleotides include RNA and DNA, and may be isolated from natural sources, synthesized *in vitro*, or prepared from a combination of natural and synthetic molecules. Sizes of polynucleotides are expressed as base pairs (abbreviated "bp"), nucleotides ("nt"), or kilobases ("kb"). Where the context allows, the latter two terms may describe polynucleotides that are single-stranded or double-stranded. When the term is applied to double-stranded molecules it is used to denote overall length and will be understood to be equivalent to the term "base pairs". It will be recognized by those skilled in the art that the two strands of a double-stranded polynucleotide may differ slightly in length and that the ends thereof may be staggered as a result of enzymatic cleavage; thus all

nucleotides within a double-stranded polynucleotide molecule may not be paired. Such unpaired ends will in general not exceed 20 nt in length.

A "polypeptide" is a polymer of amino acid residues joined by peptide bonds, whether produced naturally or synthetically. Polypeptides of less than about 10 amino acid residues are commonly referred to as "peptides".

The term "promoter" is used herein for its art-recognized meaning to denote a portion of a gene containing DNA sequences that provide for the binding of RNA polymerase and initiation of transcription. Promoter sequences are commonly, but not always, found in the 5' non-coding regions of genes.

A "protein" is a macromolecule comprising one or more polypeptide chains. A protein may also comprise non-peptidic components, such as carbohydrate groups. Carbohydrates and other non-peptidic substituents may be added to a protein by the cell in which the protein is produced, and will vary with the type of cell. Proteins are defined herein in terms of their amino acid backbone structures; substituents such as carbohydrate groups are generally not specified, but may be present nonetheless.

A "secretory signal sequence" is a DNA sequence that encodes a polypeptide (a "secretory peptide") that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger polypeptide is commonly cleaved to remove the secretory peptide during transit through the secretory pathway.

A "segment" is a portion of a larger molecule (e.g., polynucleotide or polypeptide) having specified attributes. For example, a DNA segment encoding a specified polypeptide is a portion of a longer DNA molecule, such as a plasmid or plasmid fragment, that, when read from the 5' to the 3' direction, encodes the sequence of amino acids of the specified polypeptide.

Molecular weights and lengths of polymers determined by imprecise analytical methods (e.g., gel electrophoresis) will be understood to be approximate values. When such a value is expressed as "about" X or "approximately" X, the stated value of X will be understood to be accurate to  $\pm 20\%$ .

All references cited herein are incorporated by reference in their entirety.

The present invention is based in part upon the discovery of a novel DNA molecule that encodes a polypeptide comprising a growth factor domain and a CUB domain. The growth factor domain is characterized by an arrangement of cysteine residues and beta strands that is characteristic of the "cystine knot" structure of the PDGF family. The CUB domain shows sequence homology to CUB domains in the neuropilins (Takagi et al., *Neuron* 7:295-307, 1991; Soker et al., *ibid.*), human bone

morphogenetic protein-1 (Wozney et al., *Science* 242:1528-1534, 1988), porcine seminal plasma protein and bovine acidic seminal fluid protein (Romero et al., *Nat. Struct. Biol.* 4:783-788, 1997), and *X. laevis* tolloid-like protein (Lin et al., *Dev. Growth Differ.* 39:43-51, 1997). Analysis of the tissue distribution of the mRNA corresponding to this novel DNA showed that expression was widespread in adult human tissues. The polypeptide has been designated "zveg4" in view of its homology to the VEGFs in the growth factor domain.

Structural predictions based on the zveg4 sequence and its homology to other growth factors suggests that the polypeptide can form homomultimers or heteromultimers that act on tissues to control organ development by modulating cell proliferation, migration, differentiation, or metabolism. Experimental evidence supports these predictions. Zveg4 heteromultimers may comprise a polypeptide from another member of the PDGF/VEGF family of proteins, including VEGF, VEGF-B, VEGF-C, VEGF-D, zveg3, PIGF (Maglione et al., *Proc. Natl. Acad. Sci. USA* 88:9267-9271, 1991), PDGF-A (Murray et al., U.S. Patent No. 4,899,919; Heldin et al., U.S. Patent No. 5,219,759), or PDGF-B (Chiu et al., *Cell* 37:123-129, 1984; Johnsson et al., *EMBO J.* 3:921-928, 1984). Members of this family of polypeptides regulate organ development and regeneration, post-developmental organ growth, and organ maintenance, as well as tissue maintenance and repair processes. These factors are also involved in pathological processes where therapeutic treatments are required, including cancer, rheumatoid arthritis, diabetic retinopathy, ischemic limb disease, peripheral vascular disease, myocardial ischemia, vascular intimal hyperplasia, atherosclerosis, and hemangioma formation. To treat these pathological conditions it will often be required to develop compounds to antagonize the members of the PDGF/VEGF family of proteins, or their respective receptors. This may include the development of neutralizing antibodies, small molecule antagonists, modified forms of the growth factors that maintain receptor binding activity but lack receptor activating activity, soluble receptors (including receptor-immunoglobulin fusion proteins) or antisense or ribozyme molecules to block polypeptide production.

A representative human zveg4 polypeptide sequence is shown in SEQ ID NO:2, and a representative mouse zveg4 polypeptide sequence is shown in SEQ ID NO:53. DNAs encoding these polypeptides are shown in SEQ ID NOS:1 and 52, respectively. Analysis of the amino acid sequence shown in SEQ ID NO:2 indicates that residues 1 to 18 form a secretory peptide. The CUB domain extends from residue 52 to residue 179. A propeptide-like sequence extends from residue 180 to either residue 245, residue 249 or residue 257, and includes four potential cleavage sites at its carboxyl terminus, monobasic sites at residue 245 and residue 249, a dibasic site at

residues 254-255, and a target site for furin or a furin-like protease at residues 254-257. Protein produced in a baculovirus expression system showed cleavage between residues 250 and 259, as well as longer species with amino termini at residues 19 and 35. The growth factor domain extends from residue 258 to residue 370, and may include  
5 additional residues at the N-terminus (for instance, this domain may include residues 250 to 370 or residues 246 to 370). Those skilled in the art will recognize that domain boundaries are somewhat imprecise and can be expected to vary by up to  $\pm 5$  residues from the specified positions. Cleavage of full-length zveg4 with plasmin resulted in  
10 activation of the zveg4 polypeptide. By Western analysis, a band migrating at approximately the same size as the growth factor domain was observed. A matched, uncleaved full-length zveg4 sample demonstrated no activation.

Signal peptide cleavage is predicted to occur in human zveg4 after residue 18 ( $\pm 3$  residues). Upon comparison of human and mouse zveg4 sequences, alternative signal peptide cleavage sites are predicted after residue 23 and/or residue 24.  
15 This analysis suggests that the zveg4 polypeptide chain may be cleaved to produce a plurality of monomeric species, some of which are shown in Table 1. In certain host cells, cleavage after Lys-255 is expected to result in subsequent removal of residues 254-255, although polypeptides with a carboxyl terminus at residue 255 may also be prepared. Cleavage after Lys-257 is expected to result in subsequent removal of residue  
20 257. These cleavage sites can be modified to prevent proteolysis and thus provide for the production of uncleaved zveg4 polypeptides and multimers comprising them. Actual cleavage patterns are expected to vary among host cells.

Table 1

<u>Monomer</u>	<u>Residues (SEQ ID NO:2)</u>
Cub domain	19 - 179
	35 - 179
	52 - 179
CUB domain + interdomain region	
	19 - 257
	35 - 257
	52 - 257
	19 - 255
	35 - 255
	52 - 255
	19 - 253
	35 - 253
	52 - 253
	19 - 249
	35 - 249
	52 - 249
	19 - 245
	35 - 245
	52 - 245
Cub domain + interdomain region + growth factor domain	
	19 - 370
	35 - 370
	52 - 370
Growth factor domain	
	246 - 370
	250 - 370
	258 - 370
Growth factor domain + interdomain region	
	180 - 370

Also included within the present invention are polypeptides that are at least 70%, 80%, 90%, and 95% identical to the polypeptides disclosed in Table 1, wherein these additional polypeptides retain certain characteristic sequence motifs as disclosed below.

5                Zveg4 polypeptides are designated herein with a subscript indicating the amino acid residues. For example, the CUB domain polypeptides disclosed in Table 1 are designated "zveg4<sub>19-179</sub>", "zveg4<sub>35-179</sub>", and "zveg4<sub>52-179</sub>".

Higher order structure of zveg4 polypeptides can be predicted by sequence alignment with known homologs and computer analysis using available  
 10 software (e.g., the Insight II® viewer and homology modeling tools; MSI, San Diego, CA). Analysis of SEQ ID NO:2 predicts that the secondary structure of the growth factor domain is dominated by the cystine knot, which ties together variable beta strand-like regions and loops into a bow tie-like structure. Sequence alignment indicates that Cys residues within the growth factor domain at positions 272, 302, 306, 318, 360, and  
 15 362, and Gly 304 are highly conserved within the family. Further analysis suggests pairing (disulfide bond formation) of Cys residues 272 and 318, 302 and 360, and 306 and 362 to form the cystine knot. This arrangement of conserved residues can be represented by the formula CX{18,33}CXGXCX{6,33}CX{20,50}CXC (SEQ ID NO:3), wherein amino acid residues are represented by the conventional single-letter  
 20 code, X is any amino acid residue, and {y,z} indicates a region of variable residues (X) from y to z residues in length. A consensus bow tie structure is formed as: amino terminus to cystine knot → beta strand-like region 1 → variable loop 1 → beta strand-like region 2 → cystine knot → beta strand-like region 3 → variable loop 2 → beta strand-like region 4 → cystine knot → beta strand-like region 5 → variable loop 3 →  
 25 beta strand-like region 6 → cystine knot. Variable loops 1 and 2 form one side of the bow tie, with variable loop 3 forming the other side. The structure of the zveg4 growth factor domain appears to diverge from the consensus structure of other family members in loop 2 and beta strand-like regions 3 and 4, wherein all are abbreviated and essentially replaced by a cysteine cluster comprising residues 307 (Gly) through 317  
 30 (Thr), which includes Cys residues at positions 308 and 316 of SEQ ID NO:2. The approximate boundaries of the beta strand-like regions in SEQ ID NO:2 are: region 1, residues 273-281; region 2, residues 297-301; region 5, residues 319-324; region 6, residues 355-358. Loops separate regions 1 and 2, and regions 5 and 6.

The CUB domain of zveg4 is believed to form a beta barrel structure  
 35 with nine distinct beta strand-like regions. These regions comprise residues 54-57, 61-65, 79-84, 90-95, 97-99, 112-115, 126-130, 146-150, and 163-170 of SEQ ID NO:2. A multiple alignment of CUB domains of *Xenopus laevis* neuropilin precursor (Takagi et



al., *ibid.*), human BMP-1 (Wozney et al., *ibid.*), and *X. laevis* tolloid-like protein (Lin et al., *ibid.*) indicates the presence of a conserved motif corresponding to residues 109-131 of SEQ ID NO:2. This motif is represented by the formula C[KR]Y[DNE][WYF]X{11,15}G[KR][WYF]C (SEQ ID NO:4), wherein square  
 5 brackets indicate the allowable residues at a given position and X{y,z} is as defined above.

The proteins of the present invention include proteins comprising CUB domains homologous to the CUB domain of zveg4. These homologous domains are from 100 to 120 residues in length and comprise a motif of the sequence  
 10 C[KR]Y[DNE][WYF]X{11,15}G[KR][WYF]C (SEQ ID NO:4) corresponding to residues 109-131 of SEQ ID NO:2. These homologous CUB domains are at least 70% identical to residues 52-179 of SEQ ID NO:2, at least 80% identical, at least 90% identical, or at least 95% identical to residues 52-179 of SEQ ID NO:2.

CUB domain-containing proteins of the present invention may further  
 15 include a zveg4 interdomain region or homolog thereof. The interdomain region is at least 70% identical to residues 180 to 253 of SEQ ID NO:2.

As noted above, residues 254-257 of SEQ ID NO:2 are believed to provide cleavage sites for furin or other proteases. However, polypeptides comprising a C-terminal interdomain region (e.g., zveg4<sub>52-257</sub>) can be prepared with or without one  
 20 or more of residues 254-257 at the amino terminus. In addition, polypeptides comprising another C-terminal interdomain region (e.g., zveg4<sub>52-245</sub>) can be prepared.

Additional proteins of the present invention comprise the zveg4 growth factor domain or a homolog thereof. These proteins thus comprise a polypeptide segment that is at least 70%, 80%, 90% or 95% identical to residues 258-370 of SEQ  
 25 ID NO:2, wherein the polypeptide segment comprises Cys residues at positions corresponding to residues 272, 302, 306, 318, 360, and 362 of SEQ ID NO:2; a glycine at a position corresponding to residue 304 of SEQ ID NO:2; and the sequence motif CX{18,33}CXGXCX{6,33}CX{20,50}CXC (SEQ ID NO:3) corresponding to residues 272-362 of SEQ ID NO:2.

30 Additional proteins comprising combinations of the CUB domain, interdomain region, and growth factor domain are shown above in Table 1. In each case, the invention also includes homologous proteins comprising homologous domains as disclosed above. More particularly, domains or regions in the mouse zveg4 protein corresponding to domains or regions in the human zveg4 protein are included within  
 35 the present invention.

Structural analysis and homology predict that zveg4 polypeptides complex with a second polypeptide to form multimeric proteins. These proteins

include homodimers and heterodimers. In the latter case, the second polypeptide can be a truncated or other variant zveg4 polypeptide or another polypeptide, such as a PIGF, PDGF-A, PDGF-B, VEGF, VEGF-B, VEGF-C, VEGF-D, or zveg3 polypeptide. Among the dimeric proteins within the present invention are dimers formed by non-covalent association (e.g., hydrophobic interactions) with a second subunit, either a second zveg4 polypeptide or other second subunit, or by covalent association stabilized by intermolecular disulfide bonds between cysteine residues of the component monomers. Within SEQ ID NO:2, the Cys residues at positions 296, 308, 316, and 364 may form intramolecular or intermolecular disulfide bonds.

The present invention thus provides a variety of multimeric proteins comprising a zveg4 polypeptide as disclosed above. These zveg4 polypeptides include zveg4<sub>19-179</sub>, zveg4<sub>35-179</sub>, zveg4<sub>52-179</sub>, zveg4<sub>19-245</sub>, zveg4<sub>35-245</sub>, zveg4<sub>52-245</sub>, zveg4<sub>19-249</sub>, zveg4<sub>35-249</sub>, zveg4<sub>52-249</sub>, zveg4<sub>19-253</sub>, zveg4<sub>35-253</sub>, zveg4<sub>52-253</sub>, zveg4<sub>19-255</sub>, zveg4<sub>35-255</sub>, zveg4<sub>52-255</sub>, zveg4<sub>19-257</sub>, zveg4<sub>35-257</sub>, zveg4<sub>52-257</sub>, zveg4<sub>19-370</sub>, zveg4<sub>35-370</sub>, zveg4<sub>52-370</sub>, zveg4<sub>246-370</sub>, zveg4<sub>250-370</sub>, and zveg4<sub>258-370</sub>, as well as variants and derivatives of these polypeptides as disclosed herein. These zveg4 polypeptides can be prepared as homodimers or as heterodimers with corresponding regions of related family members. For example, a zveg4 CUB domain polypeptide can be dimerized with a polypeptide comprising residues 46-170 of SEQ ID NO:33; a zveg4 growth factor domain polypeptide can be dimerized with a polypeptide comprising residues 235-345 of SEQ ID NO:33; and a zveg4 CUB domain-interdomain-growth factor domain polypeptide can be dimerized with a polypeptide comprising residues 46-345 of SEQ ID NO:33.

Percent sequence identity is determined by conventional methods. See, for example, Altschul et al., *Bull. Math. Bio.* 48:603-616, 1986, and Henikoff and Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915-10919, 1992. Briefly, two amino acid sequences are aligned to optimize the alignment scores using a gap opening penalty of 10, a gap extension penalty of 1, and the "BLOSUM62" scoring matrix of Henikoff and Henikoff (*ibid.*) as shown in Table 2 (amino acids are indicated by the standard one-letter codes). The percent identity is then calculated as:

$$\frac{\text{Total number of identical matches}}{[\text{length of the longer sequence plus the number of gaps introduced into the longer sequence in order to align the two sequences}]} \times 100$$

Table 2

	A	R	N	D	C	Q	E	G	H	I	L	K	M	F	P	S	T	W	Y	V
5	A 4	R -1 5	N -2 0 6	D -2 -2 1 6	C 0 -3 -3 -3 9	Q -1 1 0 0 -3 5	E -1 0 0 2 -4 2 5	G 0 -2 0 -1 -3 -2 -2 6	H -2 0 1 -1 -3 0 0 -2 8	I -1 -3 -3 -3 -1 -3 -3 -4 -3 4	L -1 -2 -3 -4 -1 -2 -3 -4 -3 2 4	K -1 2 0 -1 -3 1 1 -2 -1 -3 -2 5	M -1 -1 -2 -3 -1 0 -2 -3 -2 1 2 -1 5	F -2 -3 -3 -3 -2 -3 -3 -3 -1 0 0 -3 0 6	P -1 -2 -2 -1 -3 -1 -1 -2 -2 -3 -3 -1 -2 -4 7	S 1 -1 1 0 -1 0 0 0 -1 -2 -2 0 -1 -2 -1 4	T 0 -1 0 -1 -1 -1 -1 -1 -2 -2 -1 -1 -1 -1 -2 -1 1 5	W -3 -3 -4 -4 -2 -2 -3 -2 -2 -3 -2 -3 -1 1 -4 -3 -2 11	Y -2 -2 -2 -3 -2 -1 -2 -3 2 -1 -1 -2 -1 3 -3 -2 -2 2 7	V 0 -3 -3 -3 -1 -2 -2 -3 -3 3 1 -2 1 -1 -2 -2 0 -3 -1 4
10																				
15																				
20																				

The level of identity between amino acid sequences can be determined using the "FASTA" similarity search algorithm of Pearson and Lipman (*Proc. Natl. Acad. Sci. USA* 85:2444, 1988) and Pearson (*Meth. Enzymol.* 183:63, 1990). Briefly, FASTA first characterizes sequence similarity by identifying regions shared by the query sequence (e.g., SEQ ID NO:2) and a test sequence that have either the highest density of identities (if the ktup variable is 1) or pairs of identities (if ktup=2), without considering conservative amino acid substitutions, insertions, or deletions. The ten regions with the highest density of identities are then rescored by comparing the similarity of all paired amino acids using an amino acid substitution matrix, and the ends of the regions are "trimmed" to include only those residues that contribute to the highest score. If there are several regions with scores greater than the "cutoff" value (calculated by a predetermined formula based upon the length of the sequence and the ktup value), then the trimmed initial regions are examined to determine whether the regions can be joined to form an approximate alignment with gaps. Finally, the highest scoring regions of the two amino acid sequences are aligned using a modification of the Needleman-Wunsch-Sellers algorithm (Needleman and Wunsch, *J. Mol. Biol.* 48:444, 1970; Sellers, *SIAM J. Appl. Math.* 26:787, 1974), which allows for amino acid insertions and deletions. Illustrative parameters for FASTA analysis are: ktup=1, gap opening penalty=10, gap extension penalty=1, and substitution matrix=BLOSUM62. These parameters can be introduced into a FASTA program by modifying the scoring matrix file ("SMATRIX"), as explained in Appendix 2 of Pearson, 1990 (*ibid.*).

FASTA can also be used to determine the sequence identity of nucleic acid molecules using a ratio as disclosed above. For nucleotide sequence comparisons, the ktup value can range between one to six, preferably from four to six.

Within certain embodiments of the invention amino acid substitutions as compared with the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:53 are conservative substitutions. The BLOSUM62 matrix (Table 2) is an amino acid substitution matrix derived from about 2,000 local multiple alignments of protein sequence segments, representing highly conserved regions of more than 500 groups of related proteins (Henikoff and Henikoff, *ibid.*). Thus, the BLOSUM62 substitution frequencies can be used to define conservative amino acid substitutions that may be introduced into the amino acid sequences of the present invention. As used herein, the term "conservative amino acid substitution" refers to a substitution represented by a BLOSUM62 value of greater than -1. For example, an amino acid substitution is conservative if the substitution is characterized by a BLOSUM62 value of 0, 1, 2, or 3. More conservative amino acid substitutions are characterized by a BLOSUM62 value

of at least 1 (e.g., 1, 2 or 3), while still more conservative amino acid substitutions are characterized by a BLOSUM62 value of at least 2 (e.g., 2 or 3).

Polypeptides of the present invention can be prepared with one or more amino acid substitutions, deletions or additions as compared to SEQ ID NO:2 or SEQ ID NO:53. These changes can be of a minor nature, that is conservative amino acid substitutions and other changes that do not significantly affect the folding or activity of the protein or polypeptide, and include amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue, an amino or carboxyl-terminal cysteine residue to facilitate subsequent linking to maleimide-activated keyhole limpet hemocyanin, a small linker peptide of up to about 20-25 residues, or an affinity tag as disclosed above. Two or more affinity tags may be used in combination. Polypeptides comprising affinity tags can further comprise a polypeptide linker and/or a proteolytic cleavage site between the zvegf4 polypeptide and the affinity tag. Exemplary cleavage sites include, without limitation, thrombin cleavage sites and factor Xa cleavage sites.

The present invention further provides a variety of other polypeptide fusions and related multimeric proteins comprising one or more polypeptide fusions. For example, a zvegf4 polypeptide can be prepared as a fusion to a dimerizing protein as disclosed in U.S. Patents Nos. 5,155,027 and 5,567,584. Exemplary dimerizing proteins in this regard include immunoglobulin constant region domains. Dimerization can also be stabilized by fusing a zvegf4 polypeptide to a leucine zipper sequence (Riley et al., *Protein Eng.* 9:223-230, 1996; Mohamed et al., *J. Steroid Biochem. Mol. Biol.* 51:241-250, 1994). Immunoglobulin-zvegf4 polypeptide fusions and leucine zipper fusions can be expressed in genetically engineered cells to produce a variety of multimeric zvegf4 analogs. Auxiliary domains can be fused to zvegf4 polypeptides to target them to specific cells, tissues, or macromolecules (e.g., collagen). For example, a zvegf4 polypeptide or protein can be targeted to a predetermined cell type by fusing a zvegf4 polypeptide to a ligand that specifically binds to a receptor on the surface of the target cell. In this way, polypeptides and proteins can be targeted for therapeutic or diagnostic purposes. A zvegf4 polypeptide can be fused to two or more moieties, such as an affinity tag for purification and a targeting domain. Polypeptide fusions can also comprise one or more cleavage sites, particularly between domains. See, Tuan et al., *Connective Tissue Research* 34:1-9, 1996.

Zvegf4 polypeptide fusions will generally contain not more than about 1,500 amino acid residues, often not more than about 1,200 residues, more often not more than about 1,000 residues, and will in many cases be considerably smaller. For example, a zvegf4 polypeptide of 352 residues (residues 19-370 of SEQ ID NO:2) can be fused to *E. coli*  $\beta$ -galactosidase (1,021 residues; see Casadaban et al., *J. Bacteriol.*

143:971-980, 1980), a 10-residue spacer, and a 4-residue factor Xa cleavage site to yield a polypeptide of 1,387 residues. In a second example, residues 250-370 of SEQ ID NO:2 can be fused to maltose binding protein (approximately 370 residues), a 4-residue cleavage site, and a 6-residue polyhistidine tag.

5           A polypeptide comprising the zvegf4 growth factor domain (e.g., zvegf4<sub>258-370</sub> or zvegf4<sub>180-370</sub>) may be fused to a non-zvegf4 CUB domain. Within a related embodiment of the invention, a zvegf4 polypeptide comprising zvegf4 growth factor and CUB domains is fused to a non-zvegf4 CUB domain, such as a CUB-domain-comprising neuropilin polypeptide.

10           The present invention further provides polypeptide fusions comprising the zvegf4 CUB domain (e.g., zvegf4<sub>52-179</sub>). The CUB domain, with its homology to neuropilin-1, may be used to target zvegf4 or other proteins containing it to cells having cell-surface semaphorins, including endothelial cells, neuronal cells, lymphocytes, and tumor cells. The zvegf4 CUB domain can thus be joined to other moieties, including  
15 polypeptides (e.g., other growth factors, antibodies, and enzymes) and non-peptidic moieties (e.g., radionuclides, contrast agents, and the like), to target them to cells expressing cell-surface semaphorins. The cleavage sites between the CUB and growth factor domains of zvegf4 may allow for proteolytic release of the growth factor domain or other moiety through existing local proteases within tissues, or by proteases added  
20 from exogenous sources. The release of the targeted moiety may provide more localized biological effects.

          The polypeptide fusions of the present invention further include fusions between zvegf4 and zvegf3, wherein a domain of zvegf4 is replaced with the corresponding domain of zvegf3 or a variant thereof. A representative human zvegf3  
25 polypeptide sequence is shown in SEQ ID NO:33, and a representative mouse sequence is shown in SEQ ID NO:35. Within SEQ ID NO:33, the CUB domain comprises residues 46-170, the interdomain region comprises residues 171-234, and the growth factor domain comprises residues 235-345 (all  $\pm$  5 residues). A secretory peptide is predicted to be cleaved from the polypeptide after residue 14 ( $\pm$ 3 residues). Cleavage  
30 sites are predicted at residue 249, residues 254-255, and residues 254-257. Domain boundaries in mouse zvegf3 and other orthologous sequences can be determined readily by those of ordinary skill in the art by alignment with the human sequence disclosed herein. Of particular interest are fusions in which the zvegf3 CUB domain is combined with the zvegf4 growth factor domain, and fusions in which the zvegf4 CUB domain is  
35 combined with the zvegf3 growth factor domain. Within these polypeptide fusions the interdomain region may be derived from either zvegf3 or zvegf4. Polypeptide fusions

comprising zveg3 and zveg4 sequences include both full-length and truncated sequences, for example sequences analogous to those disclosed in Table 1, above.

Proteins comprising the wild-type zveg4 CUB domain and variants thereof may be used to modulate activities mediated by cell-surface semaphorins. While not wishing to be bound by theory, zveg4 may bind to semaphorins via its CUB domain. The observation that semaphorin III is involved in vascular development suggests that members of the vascular growth factor family of proteins may also be involved, especially due to the co-binding activity of VEGF and semaphorin III to neuropilin-1. Zveg4 may thus be used to design agonists and antagonist of neuropilin-semaphorin interactions. For example, the zveg4 sequence disclosed herein provides a starting point for the design of molecules that antagonize semaphorin-stimulated activities, including neurite growth, cardiovascular development, cartilage and limb development, and T and B-cell function. Additional applications include intervention in various pathologies, including rheumatoid arthritis, various forms of cancer, autoimmune disease, inflammation, retinopathies, hemangiomas, ischemic events within tissues including the heart, kidney and peripheral arteries, neuropathies, acute nerve damage, and diseases of the central and peripheral nervous systems, including stroke.

The isolated CUB domain of either mouse or human zveg4 (and multimers thereof) may also be useful to block binding of other zveg4 molecules (e.g., full-length polypeptide, isolated growth factor domain, or multimers thereof) to cell-surface molecules and/or extracellular binding sites by itself binding to such molecules or sites. In addition, the isolated CUB domain of either mouse or human zveg4 may be useful to block zveg4 binding, and/or more generally vascular endothelial growth factor binding, to neuropilin-1 (see M.L. Gagnon et al., Proc. Natl. Acad. Sci. USA 97:2573-78, 2000). Further, the second major loop of zveg4 (residues 308-316) may represent the receptor-binding loop of zveg4 (see, for instance, WO 99/13329; WO 98/10795; and W.J. LaRochelle et al., J. Biol. Chem. 267:17074-77, 1992), and thus may be useful as an antagonist of zveg4 activity. Within this peptide (zveg4 residues 308-316), Cys308 and Cys316 may or may not be disulfide bonded. Also, dimers of this peptide may be constructed such that residue Cys308 is disulfide bonded to either Cys308 or Cys316 of the homodimer partner peptide.

Amino acid sequence changes are made in zveg4 polypeptides so as to minimize disruption of higher order structure essential to biological activity. As noted above, conservative amino acid changes are generally less likely to negate activity than are non-conservative changes. Changes in amino acid residues will be made so as not to disrupt the cystine knot and "bow tie" arrangement of loops in the growth factor

domain that is characteristic of the protein family. Conserved motifs will also be maintained. The effects of amino acid sequence changes can be predicted by computer modeling as disclosed above or determined by analysis of crystal structure (see, e.g., Laphorn et al., *ibid.*). A hydrophilicity profile of SEQ ID NO:2 is shown in Fig. 1.

5 Those skilled in the art will recognize that this hydrophilicity will be taken into account when designing alterations in the amino acid sequence of a zveg4 polypeptide, so as not to disrupt the overall profile. Additional guidance in selecting amino acid substitutions is provided by a comparison of the mouse (SEQ ID NO:53) and human (SEQ ID NO:2) zveg4 sequences. The amino acid sequence is highly conserved

10 between mouse and human zveg4s, with an overall amino acid sequence identity of 85.1%.

The polypeptides of the present invention can also comprise non-naturally occurring amino acid residues. Non-naturally occurring amino acids include, without limitation, *trans*-3-methylproline, 2,4-methanoproline, *cis*-4-hydroxyproline, *trans*-4-hydroxyproline, *N*-methylglycine, *allo*-threonine, methylthreonine, hydroxyethylcysteine, hydroxyethylhomocysteine, nitroglutamine, homoglutamine, pipecolic acid, *tert*-leucine, norvaline, 2-azaphenylalanine, 3-azaphenylalanine, 4-azaphenylalanine, and 4-fluorophenylalanine. Several methods are known in the art for incorporating non-naturally occurring amino acid residues into proteins. For example,

20 an *in vitro* system can be employed wherein nonsense mutations are suppressed using chemically aminoacylated suppressor tRNAs. Methods for synthesizing amino acids and aminoacylating tRNA are known in the art. Transcription and translation of plasmids containing nonsense mutations is carried out in a cell-free system comprising an *E. coli* S30 extract and commercially available enzymes and other reagents. Proteins

25 are purified by chromatography. See, for example, Robertson et al., J. Am. Chem. Soc. 113:2722, 1991; Ellman et al., Methods Enzymol. 202:301, 1991; Chung et al., Science 259:806-809, 1993; and Chung et al., Proc. Natl. Acad. Sci. USA 90:10145-10149, 1993). In a second method, translation is carried out in *Xenopus* oocytes by microinjection of mutated mRNA and chemically aminoacylated suppressor tRNAs

30 (Turcatti et al., J. Biol. Chem. 271:19991-19998, 1996). Within a third method, *E. coli* cells are cultured in the absence of a natural amino acid that is to be replaced (e.g., phenylalanine) and in the presence of the desired non-naturally occurring amino acid(s) (e.g., 2-azaphenylalanine, 3-azaphenylalanine, 4-azaphenylalanine, or 4-fluorophenylalanine). The non-naturally occurring amino acid is incorporated into the

35 protein in place of its natural counterpart. See, Koide et al., Biochem. 33:7470-7476, 1994. Naturally occurring amino acid residues can be converted to non-naturally occurring species by *in vitro* chemical modification. Chemical modification can be



combined with site-directed mutagenesis to further expand the range of substitutions (Wynn and Richards, *Protein Sci.* 2:395-403, 1993).

Essential amino acids in the polypeptides of the present invention can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, *Science* 244, 1081-1085, 1989; Bass et al., *Proc. Natl. Acad. Sci. USA* 88:4498-4502, 1991). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for biological activity of other properties to identify amino acid residues that are critical to the activity of the molecule.

Multiple amino acid substitutions can be made and tested using known methods of mutagenesis and screening, such as those disclosed by Reidhaar-Olson and Sauer (*Science* 241:53-57, 1988) or Bowie and Sauer (*Proc. Natl. Acad. Sci. USA* 86:2152-2156, 1989). Briefly, these authors disclose methods for simultaneously randomizing two or more positions in a polypeptide, selecting for functional polypeptide, and then sequencing the mutagenized polypeptides to determine the spectrum of allowable substitutions at each position. Other methods that can be used include phage display (e.g., Lowman et al., *Biochem.* 30:10832-10837, 1991; Ladner et al., U.S. Patent No. 5,223,409; Huse, WIPO Publication WO 92/06204) and region-directed mutagenesis (Derbyshire et al., *Gene* 46:145, 1986; Ner et al., *DNA* 7:127, 1988).

Variants of the disclosed zveg4 DNA and polypeptide sequences can be generated through DNA shuffling as disclosed by Stemmer, *Nature* 370:389-391, 1994 and Stemmer, *Proc. Natl. Acad. Sci. USA* 91:10747-10751, 1994. Briefly, variant genes are generated by *in vitro* homologous recombination by random fragmentation of a parent gene followed by reassembly using PCR, resulting in randomly introduced point mutations. This technique can be modified by using a family of parent genes, such as allelic variants or genes from different species, to introduce additional variability into the process. Selection or screening for the desired activity, followed by additional iterations of mutagenesis and assay provides for rapid "evolution" of sequences by selecting for desirable mutations while simultaneously selecting against detrimental changes.

Mutagenesis methods as disclosed above can be combined with high volume or high-throughput screening methods to detect biological activity of zveg4 variant polypeptides, in particular biological activity in modulating cell proliferation or cell differentiation. For example, mitogenesis assays that measure dye incorporation or <sup>3</sup>H-thymidine incorporation can be carried out on large numbers of samples, as can cell-based assays that detect expression of a reporter gene (e.g., a luciferase gene).

Mutagenesis of the CUB domain can be used to modulate its binding to members of the semaphorin family, including enhancing or inhibiting binding to selected family members. A modified spectrum of binding activity may be desirable for optimizing therapeutic and/or diagnostic utility of proteins comprising a zveg4 CUB domain.

- 5 Direct binding utilizing labeled CUB protein can be used to monitor changes in CUB domain binding activity to selected semaphorin family members. Semaphorins of interest in this regard include isolated proteins, proteins present in cell membranes, and proteins present on cell-surfaces. The CUB domain can be labeled by a variety of methods including radiolabeling with isotopes, such as <sup>125</sup>I, conjugation to enzymes  
10 such as alkaline phosphatase or horseradish peroxidase, conjugation with biotin, and conjugation with various fluorescent markers including FITC. These and other assays are disclosed in more detail below. Mutagenized DNA molecules that encode active zveg4 polypeptides can be recovered from the host cells and rapidly sequenced using modern equipment. These methods allow the rapid determination of the importance of  
15 individual amino acid residues in a polypeptide of interest, and can be applied to polypeptides of unknown structure.

Using the methods discussed above, one of ordinary skill in the art can identify and/or prepare a variety of polypeptides that are homologous to the zveg4 polypeptides disclosed above in Table 1 and retain the biological properties of the wild-  
20 type protein. Such polypeptides can also include additional polypeptide segments as generally disclosed above.

The present invention also provides polynucleotide molecules, including DNA and RNA molecules, that encode the zveg4 polypeptides disclosed above. The polynucleotides of the present invention include the sense strand; the anti-sense strand;  
25 and the DNA as double-stranded, having both the sense and anti-sense strands annealed together by hydrogen bonds. A representative DNA sequence encoding human zveg4 polypeptides is set forth in SEQ ID NO:1, and a representative DNA sequence encoding mouse zveg4 polypeptides is set forth in SEQ ID NO:52. Additional DNA sequences encoding zveg4 polypeptides can be readily generated by those of ordinary skill in the  
30 art based on the genetic code. Counterpart RNA sequences can be generated by substitution of U for T.

Those skilled in the art will readily recognize that, in view of the degeneracy of the genetic code, considerable sequence variation is possible among polynucleotide molecules encoding zveg4 polypeptides. SEQ ID NO:6 is a degenerate  
35 DNA sequence that encompasses all DNAs that encode the zveg4 polypeptide of SEQ ID NO: 2. Those skilled in the art will recognize that the degenerate sequence of SEQ ID NO:6 also provides all RNA sequences encoding SEQ ID NO:2 by substituting U

for T. Thus, zveg4 polypeptide-encoding polynucleotides comprising nucleotides 1 - 1110, 1 - 537, 55 - 537, 103 - 537, 154 - 537, 55 - 771, 103 - 771, 154 - 771, 55 - 765, 103 - 765, 154 - 765, 55 - 759, 103 - 759, 154 - 759, 55 - 747, 103 - 747, 154 - 747, 55 - 735, 103 - 735, 154 - 735, 55 - 1110, 103 - 1110, 154 - 1110, 772 - 1110, 748 - 1110, 736 - 1110, and 538 - 1110 of SEQ ID NO:6 and their RNA equivalents are contemplated by the present invention. Table 3 sets forth the one-letter codes used within SEQ ID NO:6 to denote degenerate nucleotide positions. "Resolutions" are the nucleotides denoted by a code letter. "Complement" indicates the code for the complementary nucleotide(s). For example, the code Y denotes either C or T, and its complement R denotes A or G, A being complementary to T, and G being complementary to C.

Table 3

Nucleotide	Resolutions	Complement	Resolutions
A	A	T	T
C	C	G	G
G	G	C	C
T	T	A	A
R	A G	Y	C T
Y	C T	R	A G
M	A C	K	G T
K	G T	M	A C
S	C G	S	C G
W	A T	W	A T
H	A C T	D	A G T
B	C G T	V	A C G
V	A C G	B	C G T
D	A G T	H	A C T
N	A C G T	N	A C G T

The degenerate codons used in SEQ ID NO:6, encompassing all possible codons for a given amino acid, are set forth in Table 4, below.

TABLE 4

Amino Acid	One-Letter Code	Codons	Degenerate Codon
Cys	C	TGC TGT	TGY
Ser	S	AGC AGT TCA TCC TCG TCT	WSN
Thr	T	ACA ACC ACG ACT	CAN
Pro	P	CCA CCC CCG CCT	CCN
Ala	A	GCA GCC GCG GCT	GCN
Gly	G	GGA GGC GGG GGT	GGN
Asn	N	AAC AAT	AAY
Asp	D	GAC GAT	GAY
Glu	E	GAA GAG	GAR
Gln	Q	CAA CAG	CAR
His	H	CAC CAT	CAY
Arg	R	AGA AGG CGA CGC CGG CGT	MGN
Lys	K	AAA AAG	AAR
Met	M	ATG	ATG
Ile	I	ATA ATC ATT	ATH
Leu	L	CTA CTC CTG CTT TTA TTG	YTN
Val	V	GTA GTC GTG GTT	GTN
Phe	F	TTC TTT	TTY
Tyr	Y	TAC TAT	TAY
Trp	W	TGG	TGG
Ter	.	TAA TAG TGA	TRR
Asn Asp	B		RAY
Glu Gln	Z		SAR
Any	X		NNN
Gap	-	---	

One of ordinary skill in the art will appreciate that some ambiguity is introduced in determining a degenerate codon, representative of all possible codons encoding each amino acid. For example, the degenerate codon for serine (WSN) can, in some circumstances, encode arginine (AGR), and the degenerate codon for arginine (MGN) can, in some circumstances, encode serine (AGY). A similar relationship exists between codons encoding phenylalanine and leucine. Thus, some polynucleotides encompassed by the degenerate sequences may encode variant amino acid sequences,

but one of ordinary skill in the art can easily identify such variant sequences by reference to the amino acid sequence of SEQ ID NO: 2 and of SEQ ID NO:53. Variant sequences can be readily tested for functionality as described herein.

Within certain embodiments of the invention the isolated  
5 polynucleotides will hybridize to similar sized regions of SEQ ID NO:1 or SEQ ID NO:52, or a sequence complementary thereto, under stringent conditions. In general, stringent conditions are selected to be about 5°C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence  
10 hybridizes to a perfectly matched probe. Typical stringent conditions are those in which the salt concentration is up to about 0.03 M at pH 7 and the temperature is at least about 60°C.

As previously noted, the isolated polynucleotides of the present invention include DNA and RNA. Methods for preparing DNA and RNA are well  
15 known in the art. Complementary DNA (cDNA) clones are prepared from RNA that is isolated from a tissue or cell that produces large amounts of zveg4 RNA. Such tissues and cells are identified by Northern blotting (Thomas, *Proc. Natl. Acad. Sci. USA* 77:5201, 1980), and include heart, pancreas, stomach, and adrenal gland. Total RNA can be prepared using guanidine HCl extraction followed by isolation by centrifugation  
20 in a CsCl gradient (Chirgwin et al., *Biochemistry* 18:52-94, 1979). Poly (A)<sup>+</sup> RNA is prepared from total RNA using the method of Aviv and Leder (*Proc. Natl. Acad. Sci. USA* 69:1408-1412, 1972). Complementary DNA (cDNA) is prepared from poly(A)<sup>+</sup> RNA using known methods. In the alternative, genomic DNA can be isolated. For some applications (e.g., expression in transgenic animals) it may be advantageous to  
25 use a genomic clone, or to modify a cDNA clone to include at least one genomic intron. Methods for identifying and isolating cDNA and genomic clones are well known and within the level of ordinary skill in the art, and include the use of the sequence disclosed herein, or parts thereof, for probing or priming a library. Polynucleotides encoding zveg4 polypeptides are identified and isolated by, for example, hybridization  
30 or polymerase chain reaction ("PCR", Mullis, U.S. Patent 4,683,202). Expression libraries can be probed with antibodies to zveg4, receptor fragments, or other specific binding partners.

Those skilled in the art will recognize that the sequences disclosed in SEQ ID NOS:1 and 2 represent a single allele of human zveg4, and that the sequences  
35 disclosed in SEQ ID NOS:52 and 53 represent a single allele of mouse zveg4. Allelic variants of these sequences can be cloned by probing cDNA or genomic libraries from

different individuals according to standard procedures. Alternatively spliced forms of zveg4 are also expected to exist.

The zveg4 polynucleotide sequence disclosed herein can be used to isolate polynucleotides encoding other zveg4 proteins. Such other polynucleotides include allelic variants, alternatively spliced cDNAs and counterpart polynucleotides from other species (orthologs). These orthologous polynucleotides can be used, *inter alia*, to prepare the respective orthologous proteins. Other species of interest include, but are not limited to, mammalian, avian, amphibian, reptile, fish, insect and other vertebrate and invertebrate species. Of particular interest are zveg4 polynucleotides and proteins from other mammalian species, including non-human primate, murine, porcine, ovine, bovine, canine, feline, and equine polynucleotides and proteins. Orthologs of human zveg4 can be cloned using information and compositions provided by the present invention in combination with conventional cloning techniques. For example, a cDNA can be cloned using mRNA obtained from a tissue or cell type that expresses zveg4 as disclosed herein. Suitable sources of mRNA can be identified by probing Northern blots with probes designed from the sequences disclosed herein. A library is then prepared from mRNA of a positive tissue or cell line. A zveg4-encoding cDNA can then be isolated by a variety of methods, such as by probing with a complete or partial human cDNA or with one or more sets of degenerate probes based on the disclosed sequences. Hybridization will generally be done under low stringency conditions, wherein washing is carried out in 1 x SSC with an initial wash at 40°C and with subsequent washes at 5°C higher intervals until background is suitably reduced. A cDNA can also be cloned using the polymerase chain reaction, or PCR (Mullis, U.S. Patent No. 4,683,202), using primers designed from the representative human zveg4 sequence disclosed herein. Within an additional method, the cDNA library can be used to transform or transfect host cells, and expression of the cDNA of interest can be detected with an antibody to zveg4 polypeptide. Similar techniques can also be applied to the isolation of genomic clones.

For any zveg4 polypeptide, including variants and fusion proteins, one of ordinary skill in the art can readily generate a fully degenerate polynucleotide sequence encoding that variant using the information set forth in Tables 3 and 4, above.

Conserved regions of zveg4, identified by alignment with sequences of other family members, can be used to identify related polynucleotides and proteins. For instance, reverse transcription-polymerase chain reaction (RT-PCR) and other techniques known in the art can be used to amplify sequences encoding the conserved motifs present in zveg4 from RNA obtained from a variety of tissue sources. In particular, highly degenerate primers as shown below in Table 5 (designed from an

alignment of zveg4 with PDGF A and B chains, VEGF, VEGF-B, VEGF-C, VEGF-D, and zveg3) are useful for cloning polynucleotides encoding homologous growth factor domains. Primers shown in Table 6, designed from an alignment of zveg4 with *X. laevis* neuropilin precursor, human BMP-1, human zveg3, and *X. laevis* tolloid-like protein, are useful for cloning polynucleotides encoding CUB domains. The primers of Tables 5 and 6 can thus be used to obtain additional polynucleotides encoding homologs of the zveg4 sequence of SEQ ID NO:1 and NO:2.

Table 5

10 zveg4 residues 301-305

degenerate: MGN TGY GGN GGN AAY TG (SEQ ID NO:7)  
 consensus: MGN TGY DSN GGN WRY TG (SEQ ID NO:8)  
 complement: CAR YWN CCN SHR CAN CK (SEQ ID NO:9)

15 zveg4 residues 292-297

degenerate: TTY TTY CCN MGN TGY YT (SEQ ID NO:10)  
 consensus: NTN DDN CCN NSN TGY BT (SEQ ID NO:11)  
 complement: AVR CAN SNN GGN HHN AN (SEQ ID NO:12)

20 zveg4 residues 357-362

degenerate CAY GAR MGN TGY GAY TG (SEQ ID NO:13)  
 consensus: CAY NNN NVN TGY VVN TG (SEQ ID NO:14)  
 complement: CAN BBR CAN BNN NNR TG (SEQ ID NO:15)

25 zveg4 residues 250-255

degenerate: TGY ACN CCN MGN AAY TA (SEQ ID NO:16)  
 consensus: TGY HNN MCN MKN RMN DH (SEQ ID NO:17)  
 complement: DHN KYN MKN GKN NDR CA (SEQ ID NO:18)

30

Table 6

zveg4 residues 110-115

consensus: N TAY GAY TWY GTN GAR GT (SEQ ID NO:19)

complement: N ATR CTR AWR CAN CTY CA (SEQ ID NO:20)

5

zveg4 residues 68-73

consensus: GN TDB CCN MAN DVN TAY C (SEQ ID NO:21)

complement: CN AHV GGN KTN HBN ATR G (SEQ ID NO:22)

10 zveg4 residues 126-131

consensus: TN HDN GGN MRN TDB TGY G (SEQ ID NO:23)

complement: AN DHN CCN KYN AHV ACR C (SEQ ID NO:24)

15 Zveg4 polynucleotide sequences disclosed herein can also be used as probes or primers to clone 5' non-coding regions of a zveg4 gene, including promoter sequences. A human zveg4 genomic fragment, comprising 5' non-coding and coding sequences, is shown in SEQ ID NO:36. These flanking sequences can be used to direct the expression of zveg4 and other recombinant proteins. In addition, 5' flanking sequences can be used as targeting sites for regulatory constructs to activate or increase  
20 expression of endogenous zveg4 genes as disclosed by Treco et al., U.S. Patent No. 5,641,670. A human zveg4 genomic sequence comprising 5' non-coding sequence and approximately 100 nucleotides of coding sequence is shown in SEQ ID NO:36.

The polynucleotides of the present invention can also be prepared by automated synthesis. The production of short, double-stranded segments (60 to 80 bp)  
25 is technically straightforward and can be accomplished by synthesizing the complementary strands and then annealing them. Longer segments (typically >300 bp) are assembled in modular form from single-stranded fragments that are from 20 to 100 nucleotides in length. Automated synthesis of polynucleotides is within the level of ordinary skill in the art, and suitable equipment and reagents are available from  
30 commercial suppliers. See, in general, Glick and Pasternak, Molecular Biotechnology, Principles & Applications of Recombinant DNA, ASM Press, Washington, D.C., 1994; Itakura et al., *Ann. Rev. Biochem.* 53: 323-56, 1984; and Climie et al., *Proc. Natl. Acad. Sci. USA* 87:633-7, 1990.

35 The polypeptides of the present invention, including full-length polypeptides, biologically active fragments, and fusion polypeptides can be produced in genetically engineered host cells according to conventional techniques. Suitable host cells are those cell types that can be transformed or transfected with exogenous DNA



and grown in culture, and include bacteria, fungal cells, and cultured higher eukaryotic cells, including cultured cells of multicellular organisms. Techniques for manipulating cloned DNA molecules and introducing exogenous DNA into a variety of host cells are disclosed by Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989, and Ausubel et al., eds., *Current Protocols in Molecular Biology*, Green and Wiley and Sons, NY, 1993.

In general, a DNA sequence encoding a zveg4 polypeptide is operably linked to other genetic elements required for its expression, generally including a transcription promoter and terminator, within an expression vector. The vector will also commonly contain one or more selectable markers and one or more origins of replication, although those skilled in the art will recognize that within certain systems selectable markers may be provided on separate vectors, and replication of the exogenous DNA may be provided by integration into the host cell genome. Selection of promoters, terminators, selectable markers, vectors, and other elements is a matter of routine design within the level of ordinary skill in the art. Many such elements are described in the literature and are available through commercial suppliers.

To direct a zveg4 polypeptide into the secretory pathway of a host cell, a secretory signal sequence (also known as a leader sequence, prepro sequence or pre sequence) is provided in the expression vector. The secretory signal sequence may be that of zveg4, or may be derived from another secreted protein (e.g., t-PA; see, U.S. Patent No. 5,641,655) or synthesized *de novo*. The secretory signal sequence is operably linked to the zveg4 DNA sequence, i.e., the two sequences are joined in the correct reading frame and positioned to direct the newly synthesized polypeptide into the secretory pathway of the host cell. Secretory signal sequences are commonly positioned 5' to the DNA sequence encoding the polypeptide of interest, although certain signal sequences may be positioned elsewhere in the DNA sequence of interest (see, e.g., Welch et al., U.S. Patent No. 5,037,743; Holland et al., U.S. Patent No. 5,143,830).

Expression of zveg4 polypeptides via a host cell secretory pathway is expected to result in the production of multimeric proteins. As noted above, such multimers include both homomultimers and heteromultimers, the latter including proteins comprising only zveg4 polypeptides and proteins including zveg4 and heterologous polypeptides. For example, a heteromultimer comprising a zveg4 polypeptide and a polypeptide from a related family member (e.g., VEGF, VEGF-B, VEGF-C, VEGF-D, zveg3, PlGF, PDGF-A, or PDGF-B) can be produced by co-expression of the two polypeptides in a host cell. Sequences encoding these other family members are known. See, for example, Dvorak et al, *ibid.*; Olofsson et al, *ibid.*;

Hayward et al., *ibid.*; Joukov et al., *ibid.*; Oliviero et al., *ibid.*; Achen et al., *ibid.*; Maglione et al., *ibid.*; Heldin et al., U.S. Patent No. 5,219,759; and Johnsson et al., *ibid.* If a mixture of proteins results from expression, individual species are isolated by conventional methods. Monomers, dimers, and higher order multimers are separated  
5 by, for example, size exclusion chromatography. Heteromultimers can be separated from homomultimers by conventional chromatography or by immunoaffinity chromatography using antibodies specific for individual dimers or by sequential immunoaffinity steps using antibodies specific for individual component polypeptides. See, in general, U.S. Patent No. 5,094,941.

10 Cultured mammalian cells are suitable hosts for use within the present invention. Methods for introducing exogenous DNA into mammalian host cells include calcium phosphate-mediated transfection (Wigler et al., *Cell* 14:725, 1978; Corsaro and Pearson, *Somatic Cell Genetics* 7:603, 1981; Graham and Van der Eb, *Virology* 52:456, 1973), electroporation (Neumann et al., *EMBO J.* 1:841-845, 1982), DEAE-dextran  
15 mediated transfection (Ausubel et al., *ibid.*), and liposome-mediated transfection (Hawley-Nelson et al., *Focus* 15:73, 1993; Ciccarone et al., *Focus* 15:80, 1993). The production of recombinant polypeptides in cultured mammalian cells is disclosed by, for example, Levinson et al., U.S. Patent No. 4,713,339; Hagen et al., U.S. Patent No. 4,784,950; Palmiter et al., U.S. Patent No. 4,579,821; and Ringold, U.S. Patent No.  
20 4,656,134. Suitable cultured mammalian cells include the COS-1 (ATCC No. CRL 1650), COS-7 (ATCC No. CRL 1651), BHK (ATCC No. CRL 1632), BHK 570 (ATCC No. CRL 10314), 293 (ATCC No. CRL 1573; Graham et al., *J. Gen. Virol.* 36:59-72, 1977) and Chinese hamster ovary (e.g. CHO-K1; ATCC No. CCL 61) cell lines. Additional suitable cell lines are known in the art and available from public  
25 depositories such as the American Type Culture Collection, Manassas, Virginia. Strong transcription promoters can be used, such as promoters from SV-40 or cytomegalovirus. See, e.g., U.S. Patent No. 4,956,288. Other suitable promoters include those from metallothionein genes (U.S. Patent Nos. 4,579,821 and 4,601,978) and the adenovirus major late promoter. Expression vectors for use in mammalian cells include pZP-1 and  
30 pZP-9, which have been deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, VA USA under accession numbers 98669 and 98668, respectively.

Drug selection is generally used to select for cultured mammalian cells into which foreign DNA has been inserted. Such cells are commonly referred to as  
35 "transfectants". Cells that have been cultured in the presence of the selective agent and are able to pass the gene of interest to their progeny are referred to as "stable transfectants." An exemplary selectable marker is a gene encoding resistance to the

antibiotic neomycin. Selection is carried out in the presence of a neomycin-type drug, such as G-418 or the like. Selection systems can also be used to increase the expression level of the gene of interest, a process referred to as "amplification." Amplification is carried out by culturing transfectants in the presence of a low level of the selective agent and then increasing the amount of selective agent to select for cells that produce high levels of the products of the introduced genes. An exemplary amplifiable selectable marker is dihydrofolate reductase, which confers resistance to methotrexate. Other drug resistance genes (e.g. hygromycin resistance, multi-drug resistance, puromycin acetyltransferase) can also be used.

Other higher eukaryotic cells can also be used as hosts, including insect cells, plant cells and avian cells. The use of *Agrobacterium rhizogenes* as a vector for expressing genes in plant cells has been reviewed by Sinkar et al., *J. Biosci. (Bangalore)* 11:47-58, 1987. Transformation of insect cells and production of foreign polypeptides therein is disclosed by Guarino et al., U.S. Patent No. 5,162,222 and WIPO publication WO 94/06463.

Insect cells can be infected with recombinant baculovirus, commonly derived from *Autographa californica* nuclear polyhedrosis virus (AcNPV). See, King and Possee, The Baculovirus Expression System: A Laboratory Guide, London, Chapman & Hall; O'Reilly et al., Baculovirus Expression Vectors: A Laboratory Manual, New York, Oxford University Press., 1994; and Richardson, Ed., Baculovirus Expression Protocols. Methods in Molecular Biology, Humana Press, Totowa, NJ, 1995. Recombinant baculovirus can also be produced through the use of a transposon-based system described by Luckow et al. (*J. Virol.* 67:4566-4579, 1993). This system, which utilizes transfer vectors, is commercially available in kit form (Bac-to-Bac™ kit; Life Technologies, Rockville, MD). The transfer vector (e.g., pFastBac1™; Life Technologies) contains a Tn7 transposon to move the DNA encoding the protein of interest into a baculovirus genome maintained in *E. coli* as a large plasmid called a "bacmid." See, Hill-Perkins and Possee, *J. Gen. Virol.* 71:971-976, 1990; Bonning et al., *J. Gen. Virol.* 75:1551-1556, 1994; and Chazenbalk and Rapoport, *J. Biol. Chem.* 270:1543-1549, 1995. In addition, transfer vectors can include an in-frame fusion with DNA encoding a polypeptide extension or affinity tag as disclosed above. Using techniques known in the art, a transfer vector containing a zvegf4-encoding sequence is transformed into *E. coli* host cells, and the cells are screened for bacmids which contain an interrupted lacZ gene indicative of recombinant baculovirus. The bacmid DNA containing the recombinant baculovirus genome is isolated, using common techniques, and used to transfect *Spodoptera frugiperda* cells, such as Sf9 cells. Recombinant virus

that expresses zveg4 protein is subsequently produced. Recombinant viral stocks are made by methods commonly used the art.

For protein production, the recombinant virus is used to infect host cells, typically a cell line derived from the fall armyworm, *Spodoptera frugiperda* (e.g., Sf9 or Sf21 cells) or *Trichoplusia ni* (e.g., High Five™ cells; Invitrogen, Carlsbad, CA). See, in general, Glick and Pasternak, Molecular Biotechnology: Principles and Applications of Recombinant DNA, ASM Press, Washington, D.C., 1994. See also, U.S. Patent No. 5,300,435. Serum-free media are used to grow and maintain the cells. Suitable media formulations are known in the art and can be obtained from commercial suppliers. The cells are grown up from an inoculation density of approximately  $2-5 \times 10^5$  cells to a density of  $1-2 \times 10^6$  cells, at which time a recombinant viral stock is added at a multiplicity of infection (MOI) of 0.1 to 10, more typically near 3. Procedures used are generally described in available laboratory manuals (e.g., King and Possee, *ibid.*; O'Reilly et al., *ibid.*; Richardson, *ibid.*).

Fungal cells, including yeast cells, can also be used within the present invention. Yeast species of particular interest in this regard include *Saccharomyces cerevisiae*, *Pichia pastoris*, and *Pichia methanolica*. Methods for transforming *S. cerevisiae* cells with exogenous DNA and producing recombinant polypeptides therefrom are disclosed by, for example, Kawasaki, U.S. Patent No. 4,599,311; Kawasaki et al., U.S. Patent No. 4,931,373; Brake, U.S. Patent No. 4,870,008; Welch et al., U.S. Patent No. 5,037,743; and Murray et al., U.S. Patent No. 4,845,075. Transformed cells are selected by phenotype determined by the selectable marker, commonly drug resistance or the ability to grow in the absence of a particular nutrient (e.g., leucine). An exemplary vector system for use in *Saccharomyces cerevisiae* is the *POT1* vector system disclosed by Kawasaki et al. (U.S. Patent No. 4,931,373), which allows transformed cells to be selected by growth in glucose-containing media. Suitable promoters and terminators for use in yeast include those from glycolytic enzyme genes (see, e.g., Kawasaki, U.S. Patent No. 4,599,311; Kingsman et al., U.S. Patent No. 4,615,974; and Bitter, U.S. Patent No. 4,977,092) and alcohol dehydrogenase genes. See also U.S. Patents Nos. 4,990,446; 5,063,154; 5,139,936 and 4,661,454. Transformation systems for other yeasts, including *Hansenula polymorpha*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Kluyveromyces fragilis*, *Ustilago maydis*, *Pichia pastoris*, *Pichia methanolica*, *Pichia guilliermondii* and *Candida maltosa* are known in the art. See, for example, Gleeson et al., *J. Gen. Microbiol.* 132:3459-3465, 1986; Cregg, U.S. Patent No. 4,882,279; and Raymond et al., *Yeast* 14, 11-23, 1998. Aspergillus cells may be utilized according to the methods of McKnight et al., U.S. Patent No. 4,935,349. Methods for transforming *Acremonium chrysogenum*

are disclosed by Sumino et al., U.S. Patent No. 5,162,228. Methods for transforming *Neurospora* are disclosed by Lambowitz, U.S. Patent No. 4,486,533. Production of recombinant proteins in *Pichia methanolica* is disclosed in U.S. Patents No. 5,716,808, 5,736,383, 5,854,039, and 5,888,768.

5 Prokaryotic host cells, including strains of the bacteria *Escherichia coli*, *Bacillus* and other genera are also useful host cells within the present invention. Techniques for transforming these hosts and expressing foreign DNA sequences cloned therein are well known in the art (see, e.g., Sambrook et al., *ibid.*). When expressing a zveg4 polypeptide in bacteria such as *E. coli*, the polypeptide may be retained in the  
10 cytoplasm, typically as insoluble granules, or may be directed to the periplasmic space by a bacterial secretion sequence. In the former case, the cells are lysed, and the granules are recovered and denatured using, for example, guanidine isothiocyanate or urea. The denatured polypeptide can then be refolded and dimerized by diluting the denaturant, such as by dialysis against a solution of urea and a combination of reduced  
15 and oxidized glutathione, followed by dialysis against a buffered saline solution. In the alternative, the protein may be recovered from the cytoplasm in soluble form and isolated without the use of denaturants. The protein is recovered from the cell as an aqueous extract in, for example, phosphate buffered saline. To capture the protein of interest, the extract is applied directly to a chromatographic medium, such as an  
20 immobilized antibody or heparin-Sepharose column. Secreted polypeptides can be recovered from the periplasmic space in a soluble and functional form by disrupting the cells (by, for example, sonication or osmotic shock) to release the contents of the periplasmic space and recovering the protein, thereby obviating the need for denaturation and refolding.

25 Transformed or transfected host cells are cultured according to conventional procedures in a culture medium containing nutrients and other components required for the growth of the chosen host cells. A variety of suitable media, including defined media and complex media, are known in the art and generally include a carbon source, a nitrogen source, essential amino acids, vitamins and  
30 minerals. Media may also contain such components as growth factors or serum, as required. The growth medium will generally select for cells containing the exogenously added DNA by, for example, drug selection or deficiency in an essential nutrient which is complemented by the selectable marker carried on the expression vector or co-transfected into the host cell. *P. methanolica* cells, for example, are cultured in a  
35 medium comprising adequate sources of carbon, nitrogen and trace nutrients at a temperature of about 25°C to 35°C. Liquid cultures are provided with sufficient

aeration by conventional means, such as shaking of small flasks or sparging of fermentors.

Zveg4 polypeptides or fragments thereof can also be prepared through chemical synthesis according to methods known in the art, including exclusive solid phase synthesis, partial solid phase methods, fragment condensation or classical solution synthesis. See, for example, Merrifield, *J. Am. Chem. Soc.* 85:2149, 1963; Stewart et al., Solid Phase Peptide Synthesis (2nd edition), Pierce Chemical Co., Rockford, IL, 1984; Bayer and Rapp, *Chem. Pept. Prot.* 3:3, 1986; and Atherton et al., Solid Phase Peptide Synthesis: A Practical Approach, IRL Press, Oxford, 1989.

Covalent, multimeric complexes can also be made by isolating the desired component polypeptides and combining them *in vitro*. Covalent complexes that can be prepared in this manner include homodimers of zveg4 polypeptides, heterodimers of two different zveg4 polypeptides, and heterodimers of a zveg4 polypeptide and a polypeptide from another family member of the VEGF/PDGF family of proteins. The two polypeptides are mixed together under denaturing and reducing conditions, followed by renaturation of the proteins by removal of the denaturants. Removal can be done by, for example, dialysis or size exclusion chromatography to provide for buffer exchange. When combining two different polypeptides, the resulting renaturated proteins may form homodimers of the individual components as well as heterodimers of the two polypeptide components. See, Cao et al., *J. Biol. Chem.* 271:3154-3162, 1996.

Non-covalent complexes comprising a zveg4 polypeptide can be prepared by incubating a zveg4 polypeptide and a second polypeptide (e.g., a zveg4 polypeptide or another peptide of the PDGF/VEGF family) at near-physiological pH. In a typical reaction, polypeptides at a concentration of about 0.1-0.5  $\mu\text{g}/\mu\text{l}$  are incubated at pH $\approx$ 7.4 in a weak buffer (e.g., 0.01 M phosphate or acetate buffer); sodium chloride may be included at a concentration of about 0.1 M. At 37°C the reaction is essentially complete with 4-24 hours. See, for example, Weintraub et al., *Endocrinology* 101:225-235, 1997.

Depending upon the intended use, the polypeptides and proteins of the present invention can be purified to  $\geq 80\%$  purity,  $\geq 90\%$  purity,  $\geq 95\%$  purity, or to a pharmaceutically pure state, that is greater than 99.9% pure with respect to contaminating macromolecules, particularly other proteins and nucleic acids, and free of infectious and pyrogenic agents.

Zveg4 proteins (including chimeric polypeptides and polypeptide multimers) can be purified using fractionation and/or conventional purification methods and media, such as by a combination of chromatographic techniques. See, in general,

Affinity Chromatography: Principles & Methods, Pharmacia LKB Biotechnology, Uppsala, Sweden, 1988; and Scopes, Protein Purification: Principles and Practice, Springer-Verlag, New York, 1994. Proteins comprising a polyhistidine affinity tag (typically about 6 histidine residues) are purified by affinity chromatography on a nickel or cobalt chelate resin. See, for example, Houchuli et al., *Bio/Technol.* 6: 1321-1325, 1988. Proteins comprising a Glu-Glu tag can be purified by immunoaffinity chromatography according to conventional procedures. See, for example, Grussenmeyer et al., *ibid.* Maltose binding protein fusions are purified on an amylose column according to methods known in the art.

10           Using methods known in the art, zveg4 proteins can be prepared as monomers or multimers, glycosylated or non-glycosylated, pegylated or non-pegylated, and may or may not include an initial methionine amino acid residue.

          The invention further provides polypeptides that comprise an epitope-bearing portion of a protein as shown in SEQ ID NO:2. An "epitope" is a region of a protein to which an antibody can bind. See, for example, Geysen et al., *Proc. Natl. Acad. Sci. USA* 81:3998-4002, 1984. Epitopes can be linear or conformational, the latter being composed of discontinuous regions of the protein that form an epitope upon folding of the protein. Linear epitopes are generally at least 6 amino acid residues in length. Relatively short synthetic peptides that mimic part of a protein sequence are routinely capable of eliciting an antiserum that reacts with the partially mimicked protein. See, Sutcliffe et al., *Science* 219:660-666, 1983. Antibodies that recognize short, linear epitopes are particularly useful in analytic and diagnostic applications that employ denatured protein, such as Western blotting (Tobin, *Proc. Natl. Acad. Sci. USA* 76:4350-4356, 1979). Anti-peptide antibodies are not conformation-dependent and can be used to detect proteins in fragmented or otherwise altered forms (Niman et al., *Proc. Natl. Acad. Sci. USA* 82:7924-7928, 1985), such as might occur in body fluids or cell culture media. Antibodies to short peptides may also recognize proteins in native conformation and will thus be useful for monitoring protein expression and protein isolation, and in detecting zveg4 proteins in solution, such as by ELISA or in immunoprecipitation studies.

          Antigenic, epitope-bearing polypeptides of the present invention are useful for raising antibodies, including monoclonal antibodies, that specifically bind to a zveg4 protein. Antigenic, epitope-bearing polypeptides contain a sequence of at least six, within other embodiments at least nine, within other embodiments from 15 to about 30 contiguous amino acid residues of a zveg4 protein (e.g., SEQ ID NO:2). Polypeptides comprising a larger portion of a zveg4 protein, i.e., from 30 to 50 or 100 residues or up to the entire sequence are included. It is preferred that the amino acid

sequence of the epitope-bearing polypeptide is selected to provide substantial solubility in aqueous solvents, that is the sequence includes relatively hydrophilic residues, and hydrophobic residues are substantially avoided. Such regions of SEQ ID NO:2 include, for example, residues 39-44, 252-257, 102-107, 264-269, and 339-344. Exemplary  
5 longer peptide immunogens include peptides comprising residues (i) 131-148, (ii) 230-253, or (iii) 333-355 of SEQ ID NO:2. Peptide (ii) can be prepared with an additional C-terminal cys residue and peptide (iii) with an additional N-terminal cys residue to facilitate coupling.

As used herein, the term "antibodies" includes polyclonal antibodies,  
10 affinity-purified polyclonal antibodies, monoclonal antibodies, and antigen-binding fragments, such as F(ab')<sub>2</sub> and Fab proteolytic fragments. Genetically engineered intact antibodies or fragments, such as chimeric antibodies, Fv fragments, single chain antibodies and the like, as well as synthetic antigen-binding peptides and polypeptides, are also included. Non-human antibodies may be humanized by grafting non-human  
15 CDRs onto human framework and constant regions, or by incorporating the entire non-human variable domains (optionally "cloaking" them with a human-like surface by replacement of exposed residues, wherein the result is a "veneered" antibody). In some instances, humanized antibodies may retain non-human residues within the human variable region framework domains to enhance proper binding characteristics. Through  
20 humanizing antibodies, biological half-life may be increased, and the potential for adverse immune reactions upon administration to humans is reduced. Monoclonal antibodies can also be produced in mice that have been genetically altered to produce antibodies that have a human structure.

Methods for preparing and isolating polyclonal and monoclonal  
25 antibodies are well known in the art. See, for example, Cooligan, et al. (eds.), Current Protocols in Immunology, National Institutes of Health, John Wiley and Sons, Inc., 1995; Sambrook et al., Molecular Cloning: A Laboratory Manual, second edition, Cold Spring Harbor, NY, 1989; and Hurrell, J. G. R. (ed.), Monoclonal Hybridoma Antibodies: Techniques and Applications, CRC Press, Inc., Boca Raton, FL, 1982. As  
30 would be evident to one of ordinary skill in the art, polyclonal antibodies can be generated from inoculating a variety of warm-blooded animals such as horses, cows, goats, sheep, dogs, chickens, rabbits, mice, and rats with a zveg4 polypeptide or a fragment thereof. The immunogenicity of a zveg4 polypeptide may be increased through the use of an adjuvant, such as alum (aluminum hydroxide) or Freund's  
35 complete or incomplete adjuvant. Polypeptides useful for immunization also include fusion polypeptides, such as fusions of zveg4 or a portion thereof with an immunoglobulin polypeptide or with maltose binding protein. If the polypeptide



portion is "hapten-like", such portion may be advantageously joined or linked to a macromolecular carrier (such as keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA), or tetanus toxoid) for immunization.

Alternative techniques for generating or selecting antibodies useful  
5 herein include *in vitro* exposure of lymphocytes to zveg4 protein or peptide, and selection of antibody display libraries in phage or similar vectors (for instance, through use of immobilized or labeled zveg4 protein or peptide). Genes encoding polypeptides having potential zveg4 polypeptide binding domains can be obtained by screening random peptide libraries displayed on phage (phage display) or on bacteria, such as *E.*  
10 *coli*. Nucleotide sequences encoding the polypeptides can be obtained in a number of ways, such as through random mutagenesis and random polynucleotide synthesis. These random peptide display libraries can be used to screen for peptides that interact with a known target, which can be a protein or polypeptide, such as a ligand or receptor, a biological or synthetic macromolecule, or organic or inorganic substance.  
15 Techniques for creating and screening such random peptide display libraries are known in the art (Ladner et al., US Patent No. 5,223,409; Ladner et al., US Patent No. 4,946,778; Ladner et al., US Patent No. 5,403,484; and Ladner et al., US Patent No. 5,571,698), and random peptide display libraries and kits for screening such libraries are available commercially, for instance from Clontech Laboratories (Palo Alto, CA),  
20 Invitrogen Inc. (San Diego, CA), New England Biolabs, Inc. (Beverly, MA), and Pharmacia LKB Biotechnology Inc. (Piscataway, NJ). Random peptide display libraries can be screened using the zveg4 sequences disclosed herein to identify proteins which bind to zveg4. These "binding proteins", which interact with zveg4 polypeptides, can be used for tagging cells or for isolating homologous polypeptides by affinity  
25 purification, or they can be directly or indirectly conjugated to drugs, toxins, radionuclides, and the like. Binding proteins can also be used in analytical methods, such as for screening expression libraries and for neutralizing zveg4 activity; for diagnostic assays for determining circulating levels of polypeptides; for detecting or quantitating soluble polypeptides as marker of underlying pathology or disease; and as  
30 zveg4 antagonists to block zveg4 binding and signal transduction *in vitro* and *in vivo*.

Antibodies are determined to be specifically binding if they bind to a zveg4 polypeptide, peptide or epitope with an affinity at least 10-fold greater than the binding affinity to control (non-zveg4) polypeptide or protein. In this regard, a "non-zveg4 polypeptide" includes the related molecules VEGF, VEGF-B, VEGF-C, VEGF-  
35 D, zveg3, PIGF, PDGF-A, and PDGF-B, but excludes zveg4 polypeptides from non-human species. Due to the high level of amino acid sequence identity expected between zveg4 orthologs, antibodies specific for human zveg4 may also bind to

zveg4 from other species. The binding affinity of an antibody can be readily determined by one of ordinary skill in the art, for example, by Scatchard analysis (Scatchard, G., Ann. NY Acad. Sci. 51: 660-672, 1949). Methods for screening and isolating specific antibodies are well known in the art. See, for example, Paul (ed.),  
5 Fundamental Immunology, Raven Press, 1993; Getzoff et al., Adv. in Immunol. 43:1-98, 1988; Goding, J.W. (ed.), Monoclonal Antibodies: Principles and Practice, Academic Press Ltd., 1996; Benjamin et al., Ann. Rev. Immunol. 2:67-101, 1984.

A variety of assays known to those skilled in the art can be utilized to detect antibodies which specifically bind to zveg4 proteins or peptides. Exemplary  
10 assays are described in detail in Antibodies: A Laboratory Manual, Harlow and Lane (Eds.), Cold Spring Harbor Laboratory Press, 1988. Representative examples of such assays include: concurrent immunoelectrophoresis, radioimmunoassay, radioimmuno-precipitation, enzyme-linked immunosorbent assay (ELISA), dot blot or Western blot assay, inhibition or competition assay, and sandwich assay. In addition, antibodies can  
15 be screened for binding to wild-type versus mutant zveg4 protein or polypeptide.

Of particular interest are neutralizing antibodies, that is antibodies that block zveg4 biological activity. Within the present invention, an antibody is considered to be neutralizing if the antibody blocks at least 50% of the biological activity of a zveg4 protein when the antibody is present in a 1000-fold molar excess.  
20 Within certain embodiments of the invention the antibody will neutralize 50% of biological activity when present in a 100-fold molar excess or in a 10-fold molar excess. Within other embodiments the antibody neutralizes at least 60% of zveg4 activity, at least 70% of zveg4 activity, at least 80% of zveg4 activity, or at least 90% of zveg4 activity.

25 Antibodies to zveg4 may be used for tagging cells that express zveg4; for isolating zveg4 by affinity purification; for diagnostic assays for determining circulating levels of zveg4 polypeptides; for detecting or quantitating soluble zveg4 as a marker of underlying pathology or disease; in analytical methods employing FACS; for screening expression libraries; for generating anti-idiotypic antibodies; and as  
30 neutralizing antibodies or as antagonists to block zveg4 activity *in vitro* and *in vivo*. Suitable direct tags or labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent markers, chemiluminescent markers, magnetic particles and the like; indirect tags or labels may feature use of biotin-avidin or other complement/anti-complement pairs as intermediates. Antibodies may also be directly or indirectly  
35 conjugated to drugs, toxins, radionuclides and the like, and these conjugates used for *in vivo* diagnostic or therapeutic applications. Moreover, antibodies to zveg4 or fragments thereof may be used *in vitro* to detect denatured zveg4 or fragments thereof

in assays, for example, Western Blots or other assays known in the art. Antibodies can also be used to target an attached therapeutic or diagnostic moiety to cells expressing zveg4 or receptors for zveg4. Experimental data suggest that zveg4 may bind PDGF alpha and/or beta receptors.

5           Anti-zveg4 antibodies may be administered to recipients that would benefit from a decrease in bone proliferation or differentiation, such as those recipients suffering from osteosarcoma or osteopetrosis. In animals overexpressing zveg4, histological analysis showed proliferation of endosteal bone (particularly in trabecular  
10 of stromal cells in bone. Anti-zveg4 antibodies would interfere with these processes, and/or would diminish osteoblast proliferation and bone growth stimulation. Anti-zveg4 antibodies may also be used to antagonize production of cartilage by interfering with the ability of zveg4 to stimulate the development or proliferation of chondrocytes.

          In addition, anti-zveg4 antibodies may be used to diminish pro-fibrotic  
15 responses. Histological analysis of animals overexpressing zveg4 detected pro-fibrotic responses in certain organs, particularly liver, kidney and lung. Several diseases or conditions involve fibrosis in liver, lung and kidney. More particularly, alcoholism and viral hepatitis generally involve liver fibrosis, which is often a precursor to cirrhosis, which in turn may lead to an irreversible state of liver failure. Lung fibrosis resulting  
20 from exposure to environmental agents (e.g., asbestosis, silicosis) will often manifest as alveolitis or interstitial inflammation. Also, lung fibrosis may occur as a side effect of some cancer therapies, such as ionizing radiation or chemotherapeutic agents. Further, collagen vascular diseases, such as scleroderma and lupus, may also lead to lung fibrosis. In the kidney, the human condition of membranoproliferative  
25 glomerulonephritis may correspond to the pro-fibrotic response observed in animals overexpressing zveg4. Chronic immune complex deposition, as seen in lupus, hepatitis B and C, and chronic abscesses, may also lead to pro-fibrotic responses in the kidney. Administration of anti-zveg4 antibodies may beneficially interfere with zveg4-stimulated pro-fibrotic responses. Such responses include: sclerosing  
30 peritonitis, adhesions following surgery, particularly laparoscopic surgery, and restenosis.

          Activity of zveg4 proteins can be measured *in vitro* using cultured cells or *in vivo* by administering molecules of the claimed invention to an appropriate animal model. Target cells for use in zveg4 activity assays include vascular cells (especially  
35 endothelial cells, pericytes and smooth muscle cells), hematopoietic (myeloid and lymphoid) cells, liver cells (including hepatocytes, fenestrated endothelial cells, Kupffer cells, and Ito cells), fibroblasts (including human dermal fibroblasts and lung

fibroblasts), neurite cells (including astrocytes, glial cells, dendritic cells, and PC-12 cells), fetal lung cells, articular synoviocytes, pericytes, chondrocytes, osteoblasts, kidney mesangial cells, bone marrow stromal cells (see K. Satomura et al., J. Cell. Physiol. 177:426-38, 1998), and other cells having cell-surface PDGF receptors.

5                   Zveg4 proteins can be analyzed for receptor binding activity by a variety of methods well known in the art, including receptor competition assays (Bowen-Pope and Ross, *Methods Enzymol.* 109:69-100, 1985), use of soluble receptors, and use of receptors produced as IgG fusion proteins (U.S. Patent No. 5,750,375). Receptor binding assays can be performed on cell lines that contain known cell-surface  
10 receptors for evaluation. The receptors can be naturally present in the cell, or can be recombinant receptors expressed by genetically engineered cells. Cell types that are able to bind zveg4 can be identified through the use of a zveg4 polypeptide conjugated to a cytotoxin or other detectable molecule. Suitable detectable molecules include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent markers,  
15 chemiluminescent markers, magnetic particles, and the like. Suitable cytotoxic molecules include bacterial or plant toxins (for instance, diphtheria toxin, *Pseudomonas* exotoxin, ricin, abrin, saporin, and the like), as well as therapeutic radionuclides, such as iodine-131, rhenium-188 or yttrium-90. These can be either directly attached to the polypeptide or indirectly attached according to known methods, such as through a  
20 chelating moiety. Polypeptides can also be conjugated to cytotoxic drugs, such as adriamycin. For indirect attachment of a detectable or cytotoxic molecule, the detectable or cytotoxic molecule may be conjugated with a member of a complementary/anticomplementary pair, where the other member is bound to the polypeptide or antibody portion. For these purposes, biotin/streptavidin is an  
25 exemplary complementary/anticomplementary pair. Binding of a zveg4-toxin conjugate by cells, either in tissue culture, in organ culture, or *in vivo* will allow for the incorporation of the conjugate into the cell, causing cell death. This activity can be used to identify cell types that are able to bind and internalize zveg4. In addition to allowing for the identification of responsive cell types, toxin conjugates can be used in  
30 *in vivo* studies to identify organs and tissues where zveg4 has a biological activity by looking for pathology within the animal following injection of the conjugate.

Activity of zveg4 proteins can be measured *in vitro* using cultured cells. Mitogenic activity can be measured using known assays, including <sup>3</sup>H-thymidine incorporation assays (as disclosed by, e.g., Raines and Ross, *Methods Enzymol.*  
35 109:749-773, 1985 and Wahl et al., *Mol. Cell Biol.* 8:5016-5025, 1988), dye incorporation assays (as disclosed by, for example, Mosman, *J. Immunol. Meth.* 65:55-63, 1983 and Raz et al., *Acta Trop.* 68:139-147, 1997) or cell counts. Exemplary

mitogenesis assays measure incorporation of  $^3\text{H}$ -thymidine into (1) 20% confluent cultures to look for the ability of zveg4 proteins to further stimulate proliferating cells, and (2) quiescent cells held at confluence for 48 hours to look for the ability of zveg4 proteins to overcome contact-induced growth inhibition. See also, Gospodarowicz et al., *J. Cell. Biol.* 70:395-405, 1976; Ewton and Florini, *Endocrinol.* 106:577-583, 1980; and Gospodarowicz et al., *Proc. Natl. Acad. Sci. USA* 86:7311-7315, 1989. Cell differentiation can be assayed using suitable precursor cells that can be induced to differentiate into a more mature phenotype. For example, endothelial cells and hematopoietic cells are derived from a common ancestral cell, the hemangioblast (Choi et al., *Development* 125:725-732, 1998). Mesenchymal stem cells can also be used to measure the ability of zveg4 protein to stimulate differentiation into osteoblasts. Differentiation is indicated by the expression of osteocalcin, the ability of the cells to mineralize, and the expression of alkaline phosphatase, all of which can be measured by routine methods known in the art. Effects of zveg4 proteins on tumor cell growth and metastasis can be analyzed using the Lewis lung carcinoma model, for example as described by Cao et al., *J. Exp. Med.* 182:2069-2077, 1995. Activity of zveg4 proteins on cells of neural origin can be analyzed using assays that measure effects on neurite growth. Zveg4 can also be assayed in an aortic ring outgrowth assay (Nicosia and Ottinetti, *Laboratory Investigation* 63:115, 1990; Villaschi and Nicosia, *Am. J. Pathology* 143:181-190, 1993).

Zveg4 activity may also be detected using assays designed to measure zveg4-induced production of one or more additional growth factors or other macromolecules. Such assays include those for determining the presence of hepatocyte growth factor (HGF), epidermal growth factor (EGF), transforming growth factor alpha (TGF $\alpha$ ), interleukin-6 (IL-6), VEGF, acidic fibroblast growth factor (aFGF), and angiogenin. Suitable assays include mitogenesis assays using target cells responsive to the macromolecule of interest, receptor-binding assays, competition binding assays, immunological assays (e.g., ELISA), and other formats known in the art. Metalloprotease secretion is measured from treated primary human dermal fibroblasts, synoviocytes and chondrocytes. The relative levels of collagenase, gelatinase and stromelysin produced in response to culturing in the presence of a zveg4 protein is measured using zymogram gels (Loita and Stetler-Stevenson, *Cancer Biology* 1:96-106, 1990). Procollagen/collagen synthesis by dermal fibroblasts and chondrocytes in response to a test protein is measured using  $^3\text{H}$ -proline incorporation into nascent secreted collagen.  $^3\text{H}$ -labeled collagen is visualized by SDS-PAGE followed by autoradiography (Unemori and Amento, *J. Biol. Chem.* 265: 10681-10685, 1990). Glycosaminoglycan (GAG) secretion from dermal fibroblasts and chondrocytes is

measured using a 1,9-dimethylmethylene blue dye binding assay (Farndale et al., *Biochim. Biophys. Acta* 883:173-177, 1986). Collagen and GAG assays are also carried out in the presence of IL-1 $\beta$  or TGF- $\beta$  to examine the ability of zvegf4 protein to modify the established responses to these cytokines.

- 5                    Monocyte activation assays are carried out (1) to look for the ability of zvegf4 proteins to further stimulate monocyte activation, and (2) to examine the ability of zvegf4 proteins to modulate attachment-induced or endotoxin-induced monocyte activation (Fuhlbrigge et al., *J. Immunol.* 138: 3799-3802, 1987). IL-1 $\beta$  and TNF $\alpha$  levels produced in response to activation are measured by ELISA (Biosource, Inc. Camarillo, CA). Monocyte/macrophage cells, by virtue of CD14 (LPS receptor), are  
10                    exquisitely sensitive to endotoxin, and proteins with moderate levels of endotoxin-like activity will activate these cells.

- Hematopoietic activity of zvegf4 proteins can be assayed on various hematopoietic cells in culture. Suitable assays include primary bone marrow or  
15                    peripheral blood leukocyte colony assays, and later stage lineage-restricted colony assays, which are known in the art (e.g., Holly et al., WIPO Publication WO 95/21920). Marrow cells plated on a suitable semi-solid medium (e.g., 50% methylcellulose containing 15% fetal bovine serum, 10% bovine serum albumin, and 0.6% PSN antibiotic mix) are incubated in the presence of test polypeptide, then examined  
20                    microscopically for colony formation. Known hematopoietic factors are used as controls. Mitogenic activity of zvegf4 polypeptides on hematopoietic cell lines can be measured using <sup>3</sup>H-thymidine incorporation assays, dye incorporation assays, or cell counts (Raines and Ross, *Methods Enzymol.* 109:749-773, 1985 and Foster et al., U.S. Patent No. 5,641,655). For example, cells are cultured in multi-well microtiter plates.  
25                    Test samples and <sup>3</sup>H-thymidine are added, and the cells are incubated overnight at 37°C. Contents of the wells are transferred to filters, dried, and counted to determine incorporation of label. Cell proliferation can also be measured using a colorimetric assay based on the metabolic breakdown of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) (Mosman, *ibid.*). Briefly, a solution of MTT is added to  
30                    100  $\mu$ l of assay cells, and the cells are incubated at 37° C. After 4 hours, 200  $\mu$ l of 0.04 N HCl in isopropanol is added, the solution is mixed, and the absorbance of the sample is measured at 570 nm.

- Cell migration is assayed essentially as disclosed by Kähler et al. (*Arteriosclerosis, Thrombosis, and Vascular Biology* 17:932-939, 1997). A protein is  
35                    considered to be chemotactic if it induces migration of cells from an area of low protein concentration to an area of high protein concentration. The assay is performed using modified Boyden chambers with a polystyrene membrane separating the two chambers

(Transwell; Corning Costar Corp.). The test sample, diluted in medium containing 1% BSA, is added to the lower chamber of a 24-well plate containing Transwells. Cells are then placed on the Transwell insert that has been pretreated with 0.2% gelatin. Cell migration is measured after 4 hours of incubation at 37°C. Non-migrating cells are  
5 wiped off the top of the Transwell membrane, and cells attached to the lower face of the membrane are fixed and stained with 0.1% crystal violet. Stained cells are then counted directly using a microscope, or extracted with 10% acetic acid and absorbance is measured at 600 nm. Migration is then calculated from a standard calibration curve.

Cell adhesion activity is assayed essentially as disclosed by LaFleur et al.  
10 (*J. Biol. Chem.* 272:32798-32803, 1997). Briefly, microtiter plates are coated with the test protein, non-specific sites are blocked with BSA, and cells (such as smooth muscle cells, leukocytes, or endothelial cells) are plated at a density of approximately  $10^4$  -  $10^5$  cells/well. The wells are incubated at 37°C (typically for about 60 minutes), then non-adherent cells are removed by gentle washing. Adhered cells are quantitated by  
15 conventional methods (e.g., by staining with crystal violet, lysing the cells, and determining the optical density of the lysate). Control wells are coated with a known adhesive protein, such as fibronectin or vitronectin.

Assays for angiogenic activity are also known in the art. For example, the effect of zveg4 proteins on primordial endothelial cells in angiogenesis can be  
20 assayed in the chick chorioallantoic membrane angiogenesis assay (Leung, *Science* 246:1306-1309, 1989; Ferrara, *Ann. NY Acad. Sci.* 752:246-256, 1995). Briefly, a small window is cut into the shell of an eight-day old fertilized egg, and a test substance is applied to the chorioallantoic membrane. After 72 hours, the membrane is examined for neovascularization. Other suitable assays include microinjection of early stage quail  
25 (*Coturnix coturnix japonica*) embryos as disclosed by Drake et al. (*Proc. Natl. Acad. Sci. USA* 92:7657-7661, 1995); the rodent model of corneal neovascularization disclosed by Muthukkaruppan and Auerbach (*Science* 205:1416-1418, 1979), wherein a test substance is inserted into a pocket in the cornea of an inbred mouse; and the hamster cheek pouch assay (Höckel et al., *Arch. Surg.* 128:423-429, 1993). Induction  
30 of vascular permeability, which is indicative of angiogenic activity, is measured in assays designed to detect leakage of protein from the vasculature of a test animal (e.g., mouse or guinea pig) after administration of a test compound (Miles and Miles, *J. Physiol.* 118:228-257, 1952; Feng et al., *J. Exp. Med.* 183:1981-1986, 1996). *In vitro* assays for angiogenic activity include the tridimensional collagen gel matrix model  
35 (Pepper et al. *Biochem. Biophys. Res. Comm.* 189:824-831, 1992 and Ferrara et al., *Ann. NY Acad. Sci.* 732:246-256, 1995), which measures the formation of tube-like structures by microvascular endothelial cells; and basement membrane matrix models

(Grant et al., "Angiogenesis as a component of epithelial-mesenchymal interactions" in Goldberg and Rosen, *Epithelial-Mesenchymal Interaction in Cancer*, Birkhäuser Verlag, 1995, 235-248; Baatout, *Anticancer Research* 17:451-456, 1997), which are used to determine effects on cell migration and tube formation by endothelial cells  
5 seeded in a basement membrane extract enriched in laminin (e.g., Matrigel®; Becton Dickinson, Franklin Lakes, NJ). Angiogenesis assays can be carried out in the presence and absence of VEGF to assess possible combinatorial effects. VEGF can be used as a control within *in vivo* assays.

The activity of zvegf4 proteins, agonists, antagonists, and antibodies of  
10 the present invention can be measured, and compounds screened to identify agonists and antagonists, using assays that measure axon guidance and growth. Of particular interest are assays that indicate changes in neuron growth patterns, for example those disclosed in Hastings, WIPO Publication WO 97/29189 and Walter et al., *Development* 101:685-96, 1987. Assays to measure the effects on neuron growth are well known in  
15 the art. For example, the C assay (e.g., Raper and Kapfhammer, *Neuron* 4:21-9, 1990 and Luo et al., *Cell* 75:217-27, 1993) can be used to determine collapsing activity of zvegf4 on growing neurons. Other methods that can assess zvegf4-induced inhibition of neurite extension or divert such extension are also known. See, Goodman, *Annu. Rev. Neurosci.* 19:341-77, 1996. Conditioned media from cells expressing a zvegf4  
20 protein, a zvegf4 agonist, or a zvegf4 antagonist, or aggregates of such cells, can be placed in a gel matrix near suitable neural cells, such as dorsal root ganglia (DRG) or sympathetic ganglia explants, which have been co-cultured with nerve growth factor. Compared to control cells, zvegf4-induced changes in neuron growth can be measured (as disclosed by, for example, Messersmith et al., *Neuron* 14:949-59, 1995 and Puschel  
25 et al., *Neuron* 14:941-8, 1995). Likewise neurite outgrowth can be measured using neuronal cell suspensions grown in the presence of molecules of the present invention. See, for example, O'Shea et al., *Neuron* 7:231-7, 1991 and DeFreitas et al., *Neuron* 15:333-43, 1995. PC12 Pheochromocytoma cells (see Banker and Goslin, in *Culturing Nerve Cells*, chapter 6, "Culture and experimental use of the PC12 rat  
30 Pheochromocytoma cell line"; also, see Rydel and Greene, *J. Neuroscience* 7(11): 3639-53, November 1987) can be grown in the presence of zvegf4 to examine effects on neurite outgrowth. PC12 cells pre-treated with NGF to induce differentiation into a neuronal population can also be exposed to zvegf4 to determine the ability of zvegf4 to promote survival of neuronal cells.

35 The biological activities of zvegf4 proteins can be studied in non-human animals by administration of exogenous protein, by expression of zvegf4-encoding polynucleotides, and by suppression of endogenous zvegf4 expression through



antisense or knock-out techniques. Zveg4 proteins can be administered or expressed individually, in combination with other zveg4 proteins, or in combination with non-veg3 proteins, including other growth factors (e.g., other VEGFs, PlGFs, or PDGFs). For example, a combination of zveg4 polypeptides (e.g., a combination of zveg4<sub>19-179</sub> and zveg4<sub>258-370</sub>) can be administered to a test animal or expressed in the animal. Test animals are monitored for changes in such parameters as clinical signs, body weight, blood cell counts, clinical chemistry, histopathology, and the like.

Stimulation of coronary collateral growth can be measured in known animal models, including a rabbit model of peripheral limb ischemia and hind limb ischemia and a pig model of chronic myocardial ischemia (Ferrara et al., *Endocrine Reviews* 18:4-25, 1997). Zveg4 proteins are assayed in the presence and absence of VEGFs, angiopoietins, and basic FGF to test for combinatorial effects. These models can be modified by the use of adenovirus or naked DNA for gene delivery as disclosed in more detail below, resulting in local expression of the test protein(s).

Efficacy of zveg4 polypeptides in promoting wound healing can be assayed in animal models. One such model is the linear skin incision model of Mustoe et al. (*Science* 237:1333, 1987). In a typical procedure, a 6-cm incision is made in the dorsal pelt of an adult rat, then closed with wound clips. Test substances and controls (in solution, gel, or powder form) are applied before primary closure. Although administration is commonly limited to a single application, additional applications can be made on succeeding days by careful injection at several sites under the incision. Wound breaking strength is evaluated between 3 and 21 days post-wounding. In a second model, multiple, small, full-thickness excisions are made on the ear of a rabbit. The cartilage in the ear splints the wound, removing the variable of wound contraction from the evaluation of closure. Experimental treatments and controls are applied. The geometry and anatomy of the wound site allow for reliable quantification of cell ingrowth and epithelial migration, as well as quantitative analysis of the biochemistry of the wounds (e.g., collagen content). See, Mustoe et al., *J. Clin. Invest.* 87:694, 1991. The rabbit ear model can be modified to create an ischemic wound environment, which more closely resembles the clinical situation (Ahn et al., *Ann. Plast. Surg.* 24:17, 1990). Within a third model, healing of partial-thickness skin wounds in pigs or guinea pigs is evaluated (LeGrand et al., *Growth Factors* 8:307, 1993). Experimental treatments are applied daily on or under dressings. Seven days after wounding, granulation tissue thickness is determined. This model is commonly used for dose-response studies, as it is more quantitative than other *in vivo* models of wound healing. A full thickness excision model can also be employed. Within this model, the epidermis and dermis are removed down to the panniculus carnosum in rodents or the

- subcutaneous fat in pigs. Experimental treatments are applied topically on or under a dressing, and can be applied daily if desired. The wound closes by a combination of contraction and cell ingrowth and proliferation. Measurable endpoints include time to wound closure, histologic score, and biochemical parameters of wound tissue.
- 5 Impaired wound healing models are also known in the art (e.g., Cromack et al., *Surgery* 113:36, 1993; Pierce et al., *Proc. Natl. Acad. Sci. USA* 86:2229, 1989; Greenhalgh et al., *Amer. J. Pathol.* 136:1235, 1990). Delay or prolongation of the wound healing process can be induced pharmacologically by treatment with steroids, irradiation of the wound site, or by concomitant disease states (e.g., diabetes). Linear incisions or full-
- 10 thickness excisions are most commonly used as the experimental wound. Endpoints are as disclosed above for each type of wound. Subcutaneous implants can be used to assess compounds acting in the early stages of wound healing (Broadley et al., *Lab. Invest.* 61:571, 1985; Sprugel et al., *Amer. J. Pathol.* 129: 601, 1987). Implants are prepared in a porous, relatively non-inflammatory container (e.g., polyethylene sponges
- 15 or expanded polytetrafluoroethylene implants filled with bovine collagen) and placed subcutaneously in mice or rats. The interior of the implant is empty of cells, producing a "wound space" that is well-defined and separable from the preexisting tissue. This arrangement allows the assessment of cell influx and cell type as well as the measurement of vasculogenesis/angiogenesis and extracellular matrix production.
- 20 Expression of zvegf4 proteins in animals provides models for study of the biological effects of overproduction or inhibition of protein activity *in vivo*. Zvegf4-encoding polynucleotides can be introduced into test animals, such as mice, using viral vectors or naked DNA, or transgenic animals can be produced. A zvegf4 protein will commonly be expressed with a secretory peptide. Suitable secretory
- 25 peptides include the zvegf4 secretory peptide (e.g., residues 1-18 of SEQ ID NO:2) and heterologous secretory peptides. An exemplary heterologous secretory peptide is that of human tissue plasminogen activator (t-PA). The t-PA secretory peptide may be modified to reduce undesired proteolytic cleavage as disclosed in U.S. Patent No. 5,641,655.
- 30 One *in vivo* approach for assaying proteins of the present invention utilizes viral delivery systems. Exemplary viruses for this purpose include adenovirus, herpesvirus, retroviruses, vaccinia virus, and adeno-associated virus (AAV). Adenovirus, a double-stranded DNA virus, is currently the best studied gene transfer vector for delivery of heterologous nucleic acids. For review, see Becker et al., *Meth.*
- 35 *Cell Biol.* 43:161-89, 1994; and Douglas and Curiel, *Science & Medicine* 4:44-53, 1997. The adenovirus system offers several advantages. Adenovirus can (i) accommodate relatively large DNA inserts; (ii) be grown to high-titer; (iii) infect a

broad range of mammalian cell types; and (iv) be used with many different promoters including ubiquitous, tissue specific, and regulatable promoters. Because adenoviruses are stable in the bloodstream, they can be administered by intravenous injection.

Using adenovirus vectors where portions of the adenovirus genome are  
5 deleted, inserts are incorporated into the viral DNA by direct ligation or by homologous recombination with a co-transfected plasmid. In an exemplary system, the essential E1 gene has been deleted from the viral vector, and the virus will not replicate unless the E1 gene is provided by the host cell (the human 293 cell line is exemplary). When intravenously administered to intact animals, adenovirus primarily targets the liver. If  
10 the adenoviral delivery system has an E1 gene deletion, the virus cannot replicate in the host cells. However, the host's tissue (e.g., liver) will express and process (and, if a secretory signal sequence is present, secrete) the heterologous protein. Secreted proteins will enter the circulation in the highly vascularized liver, and effects on the infected animal can be determined. Intranasal delivery of adenovirus expressing zvegf4  
15 will target the zvegf4 protein to lung tissue. Further, adenovirus expressing zvegf4 can be administered directly into brain tissue. Adenoviral vectors containing various deletions of viral genes can be used in an attempt to reduce or eliminate immune responses to the vector. Such adenoviruses are E1 deleted, and in addition contain deletions of E2A or E4 (Lusky et al., *J. Virol.* 72:2022-2032, 1998; Raper et al., *Human*  
20 *Gene Therapy* 9:671-679, 1998). In addition, deletion of E2b is reported to reduce immune responses (Amalfitano, et al., *J. Virol.* 72:926-933, 1998). Generation of so-called "gutless" adenoviruses where all viral transcription units are deleted is particularly advantageous for insertion of large inserts of heterologous DNA. For review, see Yeh and Perricaudet, *FASEB J.* 11:615-623, 1997.

25 In another embodiment, a zvegf4 gene can be introduced in a retroviral vector as described, for example, by Anderson et al., U.S. Patent No. 5,399,346; Mann et al., *Cell* 33:153, 1983; Temin et al., U.S. Patent No. 4,650,764; Temin et al., U.S. Patent No. 4,980,289; Markowitz et al., *J. Virol.* 62:1120, 1988; Temin et al., U.S. Patent No. 5,124,263; Dougherty et al., WIPO publication WO 95/07358; and Kuo et  
30 al., *Blood* 82:845, 1993.

In an alternative method, the vector can be introduced by "lipofection" *in vivo* using liposomes. Synthetic cationic lipids can be used to prepare liposomes for *in vivo* transfection of a gene encoding a marker (Felgner et al., *Proc. Natl. Acad. Sci. USA* 84:7413-7, 1987; Mackey et al., *Proc. Natl. Acad. Sci. USA* 85:8027-31, 1988).  
35 The use of lipofection to introduce exogenous genes into specific organs *in vivo* has certain practical advantages. Molecular targeting of liposomes to specific cells represents one area of benefit. For instance, directing transfection to particular cell

types is particularly advantageous in a tissue with cellular heterogeneity, such as the pancreas, liver, kidney, and brain. Lipids may be chemically coupled to other molecules for the purpose of targeting. Targeted peptides (e.g., hormones or neurotransmitters), proteins such as antibodies, or non-peptide molecules can be coupled to liposomes chemically.

Within another embodiment target cells are removed from the animal, and the DNA is introduced as a naked DNA plasmid. The transformed cells are then re-implanted into the body of the animal. Naked DNA vectors can be introduced into the desired host cells by methods known in the art, e.g., transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, use of a gene gun or use of a DNA vector transporter. See, e.g., Wu et al., *J. Biol. Chem.* 267:963-7, 1992; Wu et al., *J. Biol. Chem.* 263:14621-4, 1988.

Mice engineered to express the *zveg4* gene, referred to as "transgenic mice," and mice that exhibit a complete absence of *zveg4* gene function, referred to as "knockout mice," can also be generated (Snouwaert et al., *Science* 257:1083, 1992; Lowell et al., *Nature* 366:740-42, 1993; Capecchi, *Science* 244:1288-1292, 1989; Palmiter et al., *Ann. Rev. Genet.* 20:465-499, 1986). Transgenesis experiments can be performed using normal mice or mice with genetic disease or other altered phenotypes. Transgenic mice that over-express *zveg4*, either ubiquitously or under a tissue-specific or tissue-restricted promoter, can be used to determine whether or not over-expression causes a phenotypic change. Exemplary promoters include metallothionein, albumin, ApoA1 and enolase gene promoters. The metallothionein-1 (MT-1) promoter provides expression in liver and other tissues, often leading to high levels of circulating protein. Over-expression of a wild-type *zveg4* polypeptide, polypeptide fragment or a mutant thereof may alter normal cellular processes, resulting in a phenotype that identifies a tissue in which *zveg4* expression is functionally relevant and may indicate a therapeutic target for the *zveg4*, its agonists or antagonists. For example, a transgenic mouse can be engineered to over-expresses a full-length *zveg4* sequence, which may result in a phenotype that shows similarity with human diseases. Similarly, knockout *zveg4* mice can be used to determine where *zveg4* is absolutely required *in vivo*. The phenotype of knockout mice is predictive of the *in vivo* effects of *zveg4* antagonists. Knockout mice can also be used to study the effects of *zveg4* proteins in models of disease, including, for example, cancer, atherosclerosis, rheumatoid arthritis, ischemia, and cardiovascular disease. The human *zveg4* cDNA can be used to isolate murine *zveg4* mRNA, cDNA and genomic DNA as disclosed above, which are subsequently used to generate knockout mice. These mice may be employed to study the *zveg4* gene and the protein encoded thereby in an *in vivo* system, and can be used as *in vivo* models

for corresponding human diseases. Moreover, transgenic mice expressing zveg4 antisense polynucleotides or ribozymes directed against zveg4, described herein, can be used analogously to knockout mice described above.

Antisense methodology can be used to inhibit zveg4 gene transcription to examine the effects of such inhibition *in vivo*. Polynucleotides that are complementary to a segment of a zveg4-encoding polynucleotide (e.g., a polynucleotide as set forth in SEQ ID NO:1) are designed to bind to zveg4-encoding mRNA and to inhibit translation of such mRNA. Such antisense oligonucleotides can also be used to inhibit expression of zveg4 polypeptide-encoding genes in cell culture.

10 Zveg4 proteins may be used therapeutically in human and veterinary medicine to stimulate tissue development or repair, or cellular differentiation or proliferation. Specific applications include, without limitation: the treatment of full-thickness skin wounds, including venous stasis ulcers and other chronic, non-healing wounds, particularly in cases of compromised wound healing due to diabetes mellitus, 15 connective tissue disease, smoking, burns, and other exacerbating conditions; fracture repair; skin grafting; within reconstructive surgery to promote neovascularization and increase skin flap survival; to establish vascular networks in transplanted cells and tissues, such as transplanted islets of Langerhans; to treat female reproductive tract disorders, including acute or chronic placental insufficiency (an important factor 20 causing perinatal morbidity and mortality) and prolonged bleeding; to promote the growth of tissue damaged by periodontal disease; to promote endothelialization of vascular grafts and stents; in the treatment of acute and chronic lesions of the gastrointestinal tract, including duodenal ulcers, which are characterized by a deficiency of microvessels; to promote angiogenesis and prevent neuronal degeneration due to 25 acute or chronic cerebral ischemia; to accelerate the formation of collateral blood vessels in ischemic limbs; to promote vessel re-endothelialization and to reduce intimal hyperplasia following invasive procedures such as balloon angioplasty and stent placement; to promote vessel repair and development of collateral circulation following myocardial infarction so as to limit ischemic injury; and to stimulate hematopoiesis. 30 The polypeptides are also useful additives in tissue adhesives for promoting revascularization of the healing tissue.

Of particular interest is the use of zveg4 for the treatment or repair of liver damage, including damage due to chronic liver disease, including chronic active hepatitis and many other types of cirrhosis. Widespread, massive necrosis, including 35 destruction of virtually the entire liver, can be caused by, *inter alia*, fulminant viral hepatitis; overdoses of the analgesic acetaminophen; exposure to other drugs and chemicals such as halothane, monoamine oxidase inhibitors, agents employed in the

treatment of tuberculosis, phosphorus, carbon tetrachloride, and other industrial chemicals. Conditions associated with ultrastructural lesions that do not necessarily produce obvious liver cell necrosis include Reye's syndrome in children, tetracycline toxicity, and acute fatty liver of pregnancy. Cirrhosis, a diffuse process characterized by fibrosis and a conversion of normal architecture into structurally abnormal nodules, can come about for a variety reasons including alcohol abuse, post necrotic cirrhosis (usually due to chronic active hepatitis), biliary cirrhosis, pigment cirrhosis, cryptogenic cirrhosis, Wilson's disease, and alpha-1-antitrypsin deficiency. Zveg4 may also be useful for the treatment of hepatic chronic passive congestion (CPC) and central hemorrhagic necrosis (CHN), which are two circulatory changes representing a continuum encountered in right-sided heart failure. Other circulatory disorders that may be treated with zveg4 include hepatic vein thrombosis, portal vein thrombosis, and cardiac sclerosis. In cases of liver fibrosis, it may be beneficial to administer a zveg4 antagonist to suppress the activation of stellate cells, which have been implicated in the production of extracellular matrix in fibrotic liver (Li and Friedman, *J. Gastroenterol. Hepatol.* 14:618-633, 1999). More generally, zveg4 may be beneficially used as an anti-fibrotic agent. Conditions that are characterized by a pro-fibrotic response include sclerosing peritonitis; adhesions following surgery (particularly laparoscopic surgery), which may lead to small bowel obstruction, difficulties on re-operation, pelvic adhesions and pelvic pain (see N. Panay and A.M. Lower, *Curr. Opin. Obstet. Gynecol.* 11:379-85, 1999); pulmonary fibrosis; kidney fibrosis; and restenosis.

Zveg4 polypeptides can be administered alone or in combination with other vasculogenic or angiogenic agents, including VEGF and angiopoietins 1 and 2. For example, basic and acidic FGFs, Ang-1, Ang-2, and VEGF have been found to play a role in the development of collateral circulation, and the combined use of zveg4 with one or more of these factors may be advantageous. VEGF has also been implicated in the survival of transplanted islet cells (Gorden et al. *Transplantation* 63:436-443, 1997; Pepper, *Arteriosclerosis, Throm. and Vascular Biol.* 17:605-619, 1997). Basic FGF has been shown to induce angiogenesis and accelerate healing of ulcers in experimental animals (reviewed by Folkman, *Nature Medicine* 1:27-31, 1995). VEGF has been shown to promote vessel re-endothelialization and to reduce intimal hyperplasia in animal models of restenosis (Asahara et al., *Circulation* 91:2802-2809, 1995; Callow et al., *Growth Factors* 10:223-228, 1994); efficacy of zveg4 polypeptides can be tested in these and other known models. When using zveg4 in combination with an additional agent, the two compounds can be administered simultaneously or sequentially as appropriate for the specific condition being treated.

Zveg4 proteins may be used either alone or in combination with other hematopoietic factors such as IL-3, G-CSF, GM-CSF, or stem cell factor to enhance expansion and mobilization of hematopoietic stem cells, including endothelial precursor stem cells. Cells that can be expanded in this manner include cells isolated from bone marrow, including bone marrow stromal cells (see K. Satomura et al., *J. Cell. Physiol.* 177:426-38, 1998), or cells isolated from blood. Zveg4 proteins may also be given directly to an individual to enhance endothelial stem cell production and differentiation within the treated individual. The stem cells, either developed within the patient, or provided back to a patient, may then play a role in modulating areas of ischemia within the body, thereby providing a therapeutic effect. These cells may also be useful in enhancing re-endothelialization of areas devoid of endothelial coverage, such as vascular grafts, vascular stents, and areas where the endothelial coverage has been damaged or removed (e.g., areas of angioplasty). Zveg4 proteins may also be used in combination with other growth and differentiation factors such as angiopoietin-1 (Davis et al., *Cell* 87:1161-1169, 1996) to help create and stabilize new vessel formation in areas requiring neovascularization, including areas of ischemia (cardiac or peripheral ischemia), organ transplants, wound healing, and tissue grafting.

Zveg4 proteins, agonists and antagonists may be used to modulate neurite growth and development and demarcate nervous system structures. As such, Zveg4 proteins, agonists, and antagonists would be useful as a treatment of peripheral neuropathies by increasing spinal cord and sensory neurite outgrowth. A zveg4 antagonist could be part of a therapeutic treatment for the regeneration of neurite outgrowths following strokes, brain damage caused by head injuries and paralysis caused by spinal injuries. Application may also be made in treating neurodegenerative diseases such as multiple sclerosis, Alzheimer's disease and Parkinson's disease. Application may also be made in mediating development and innervation pattern of stomach tissue.

Zveg4 has been found to have PDGF-like activity, including mitogenic activity on fibroblasts, vascular smooth muscle cells, and pericytes. Zveg4 has also been found to stimulate bone growth in an animal model. These results suggest that zveg4 proteins will be useful in promoting the growth of bone and ligament. Such uses include, for example, treatment of periodontal disease, fractures (including non-union fractures), implant recipient sites, bone grafts, and joint injuries involving cartilage and/or ligament damage. Zveg4 may be used in combination with other bone stimulating factors, such as IGF-1, EGF, TGF- $\beta$ , PDGF, and BMPs. Methods for using growth factors in the treatment of periodontal disease are known in the art. See, for example, U.S. Patent No. 5,124,316 and Lynch et al., *ibid*.

For pharmaceutical use, zveg4 proteins, antagonist, and antibodies are formulated for topical or parenteral, particularly intravenous or subcutaneous, delivery according to conventional methods. In general, pharmaceutical formulations will include a zveg4 polypeptide in combination with a pharmaceutically acceptable vehicle, such as saline, buffered saline, 5% dextrose in water, or the like. Formulations may further include one or more excipients, preservatives, solubilizers, buffering agents, albumin to prevent protein loss on vial surfaces, thickeners, gelling agents, etc. Methods of formulation are well known in the art and are disclosed, for example, in *Remington: The Science and Practice of Pharmacy*, Gennaro, ed., Mack Publishing Co., Easton, PA, 19th ed., 1995. Zveg4 will ordinarily be used in a concentration of about 10 to 100 µg/ml of total volume, although concentrations in the range of 1 ng/ml to 1000 µg/ml may be used. For topical application, such as for the promotion of wound healing, the protein will be applied in the range of 0.1-10 µg/cm<sup>2</sup> of wound area, with the exact dose determined by the clinician according to accepted standards, taking into account the nature and severity of the condition to be treated, patient traits, etc. Determination of dose is within the level of ordinary skill in the art. The therapeutic formulations will generally be administered over the period required for neovascularization, typically from one to several months and, in treatment of chronic conditions, for a year or more. Dosing is daily or intermittently over the period of treatment. Intravenous administration will be by bolus injection or infusion over a typical period of one to several hours. Sustained release formulations can also be employed. In general, a therapeutically effective amount of zveg4 is an amount sufficient to produce a clinically significant change in the treated condition, such as a clinically significant reduction in time required by wound closure, a significant reduction in wound area, a significant improvement in vascularization, a significant reduction in morbidity, or a significantly increased histological score.

Proteins of the present invention are useful for modulating the proliferation, differentiation, migration, or metabolism of responsive cell types, which include both primary cells and cultured cell lines. Of particular interest in this regard are hematopoietic cells (including stem cells and mature myeloid and lymphoid cells), endothelial cells, neuronal cells, mesenchymal cells (including fibroblasts, pericytes, stellate cells, mesangial cells, chondrocytes and smooth muscle cells), and bone-derived cells (including osteoblast and osteoclast precursors). Zveg4 polypeptides are added to tissue culture media for these cell types at a concentration of about 10 pg/ml to about 1000 ng/ml. Those skilled in the art will recognize that zveg4 proteins can be advantageously combined with other growth factors in culture media.



Within the laboratory research field, zvegf4 proteins can also be used as molecular weight standards; as reagents in assays for determining circulating levels of the protein, such as in the diagnosis of disorders characterized by over- or under-production of zvegf4 protein; or as standards in the analysis of cell phenotype.

5           Zvegf4 proteins can also be used to identify inhibitors of their activity. Test compounds are added to the assays disclosed above to identify compounds that inhibit the activity of zvegf4 protein. In addition to those assays disclosed above, samples can be tested for inhibition of zvegf4 activity within a variety of assays designed to measure receptor binding or the stimulation/inhibition of zvegf4-dependent  
10 cellular responses. For example, zvegf4-responsive cell lines can be transfected with a reporter gene construct that is responsive to a zvegf4-stimulated cellular pathway. Reporter gene constructs of this type are known in the art, and will generally comprise a zvegf4-activated serum response element (SRE) operably linked to a gene encoding an assayable protein, such as luciferase. Candidate compounds, solutions, mixtures or  
15 extracts are tested for the ability to inhibit the activity of zvegf4 on the target cells as evidenced by a decrease in zvegf4 stimulation of reporter gene expression. Assays of this type will detect compounds that directly block zvegf4 binding to cell-surface receptors, as well as compounds that block processes in the cellular pathway subsequent to receptor-ligand binding. In the alternative, compounds or other samples can be  
20 tested for direct blocking of zvegf4 binding to receptor using zvegf4 tagged with a detectable label (e.g., <sup>125</sup>I, biotin, horseradish peroxidase, FITC, and the like). Within assays of this type, the ability of a test sample to inhibit the binding of labeled zvegf4 to the receptor is indicative of inhibitory activity, which can be confirmed through secondary assays. Receptors used within binding assays may be cellular receptors or  
25 isolated, immobilized receptors.

The activity of zvegf4 proteins can be measured with a silicon-based biosensor microphysiometer that measures the extracellular acidification rate or proton excretion associated with receptor binding and subsequent physiologic cellular responses. An exemplary such device is the Cytosensor™ Microphysiometer  
30 manufactured by Molecular Devices, Sunnyvale, CA. A variety of cellular responses, such as cell proliferation, ion transport, energy production, inflammatory response, regulatory and receptor activation, and the like, can be measured by this method. See, for example, McConnell et al., *Science* 257:1906-1912, 1992; Pitchford et al., *Meth. Enzymol.* 228:84-108, 1997; Arimilli et al., *J. Immunol. Meth.* 212:49-59, 1998; and  
35 Van Liefde et al., *Eur. J. Pharmacol.* 346:87-95, 1998. The microphysiometer can be used for assaying adherent or non-adherent eukaryotic or prokaryotic cells. By measuring extracellular acidification changes in cell media over time, the

microphysiometer directly measures cellular responses to various stimuli, including zveg4 proteins, their agonists, and antagonists. The microphysiometer can be used to measure responses of a zveg4-responsive eukaryotic cell, compared to a control eukaryotic cell that does not respond to zveg4 polypeptide. Zveg4-responsive eukaryotic cells comprise cells into which a receptor for zveg4 has been transfected creating a cell that is responsive to zveg4, as well as cells naturally responsive to zveg4 such as cells derived from vascular or neural tissue. Differences, measured by a change in extracellular acidification, in the response of cells exposed to zveg4 polypeptide relative to a control not exposed to zveg4, are a direct measurement of zveg4-modulated cellular responses. Moreover, such zveg4-modulated responses can be assayed under a variety of stimuli. The present invention thus provides methods of identifying agonists and antagonists of zveg4 proteins, comprising providing cells responsive to a zveg4 polypeptide, culturing a first portion of the cells in the absence of a test compound, culturing a second portion of the cells in the presence of a test compound, and detecting a change in a cellular response of the second portion of the cells as compared to the first portion of the cells. The change in cellular response is shown as a measurable change in extracellular acidification rate. Culturing a third portion of the cells in the presence of a zveg4 protein and the absence of a test compound provides a positive control for the zveg4-responsive cells and a control to compare the agonist activity of a test compound with that of the zveg4 polypeptide. Antagonists of zveg4 can be identified by exposing the cells to zveg4 protein in the presence and absence of the test compound, whereby a reduction in zveg4-stimulated activity is indicative of antagonist activity in the test compound.

Zveg4 proteins can also be used to identify cells, tissues, or cell lines that respond to a zveg4-stimulated pathway. The microphysiometer, described above, can be used to rapidly identify ligand-responsive cells, such as cells responsive to zveg4 proteins. Cells are cultured in the presence or absence of zveg4 polypeptide. Those cells that elicit a measurable change in extracellular acidification in the presence of zveg4 are responsive to zveg4. Responsive cells can then be used to identify antagonists and agonists of zveg4 polypeptide as described above.

Inhibitors of zveg4 activity (zveg4 antagonists) include anti-zveg4 antibodies and soluble zveg4 receptors, as well as other peptidic and non-peptidic agents, including ribozymes, small molecule inhibitors, and angiogenically or mitogenically inactive receptor-binding fragments of zveg4 polypeptides. Such antagonists can be used to block biological activities of zveg4, including mitogenic, chemotactic, or angiogenic effects. These antagonists are therefore useful in reducing the growth of solid tumors by inhibiting neovascularization of the developing tumor or

by directly blocking tumor cell growth; in the treatment of diabetic retinopathy, psoriasis, arthritis, and scleroderma; and in reducing fibrosis, including scar formation. Inhibitors of zveg4 may also be useful in the treatment of proliferative vascular disorders wherein zveg4 activity is pathogenic. Such disorders may include  
5 atherosclerosis and intimal hyperplastic restenosis following angioplasty, endarterectomy, vascular grafting, organ transplant, or vascular stent emplacement. These conditions involve complex growth factor-mediated responses wherein certain factors may be beneficial to the clinical outcome and others may be pathogenic.

Inhibitors of zveg4 may also prove useful in the treatment of ocular  
10 neovascularization, including diabetic retinopathy and age-related macular degeneration. Experimental evidence suggests that these conditions result from the expression of angiogenic factors induced by hypoxia in the retina.

Zveg4 antagonists are also of interest in the treatment of inflammatory disorders, such as rheumatoid arthritis and psoriasis. In rheumatoid arthritis, studies  
15 suggest that VEGF plays an important role in the formation of pannus, an extensively vascularized tissue that invades and destroys cartilage. Psoriatic lesions are hypervascular and overexpress the angiogenic polypeptide IL-8.

Zveg4 antagonists may also prove useful in the treatment of infantile hemangiomas, which exhibit overexpression of VEGF and bFGF during the  
20 proliferative phase.

Inhibitors are formulated for pharmaceutical use as generally disclosed above, taking into account the precise chemical and physical nature of the inhibitor and the condition to be treated. The relevant determinations are within the level of ordinary skill in the formulation art. Other angiogenic and vasculogenic factors, including  
25 VEGF and bFGF, have been implicated in pathological neovascularization. In such instances it may be advantageous to combine a zveg4 inhibitor with one or more inhibitors of these other factors.

The polypeptides, nucleic acids, and antibodies of the present invention may be used in diagnosis or treatment of disorders associated with cell loss or abnormal  
30 cell proliferation (including cancer), including impaired or excessive vasculogenesis or angiogenesis, and diseases of the nervous system. Labeled zveg4 polypeptides may be used for imaging tumors or other sites of abnormal cell proliferation. Because angiogenesis in adult animals is generally limited to wound healing and the female reproductive cycle, it is a very specific indicator of pathological processes.  
35 Angiogenesis is indicative of, for example, developing solid tumors, retinopathies, and arthritis.

Zveg4 polypeptides and anti-zveg4 antibodies can be directly or indirectly conjugated to drugs, toxins, radionuclides and the like, and these conjugates used for *in vivo* diagnostic or therapeutic applications. For instance, polypeptides or antibodies of the present invention may be used to identify or treat tissues or organs that  
5 express a corresponding anti-complementary molecule (receptor or antigen, respectively, for instance). More specifically, zveg4 polypeptides or anti-zveg4 antibodies, or bioactive fragments or portions thereof, can be coupled to detectable or cytotoxic molecules and delivered to a mammal having cells, tissues, or organs that express the anti-complementary molecule. For example, the CUB domain of zveg4  
10 can be used to target peptidic and non-peptidic moieties to semaphorins as disclosed above. In another embodiment, polypeptide-toxin fusion proteins or antibody/fragment-toxin fusion proteins may be used for targeted cell or tissue inhibition or ablation, such as in cancer therapy. Of particular interest in this regard are conjugates of a zveg4 polypeptide and a cytotoxin, which can be used to target the  
15 cytotoxin to a tumor or other tissue that is undergoing undesired angiogenesis or neovascularization.

In another embodiment, zveg4-cytokine fusion proteins or antibody/fragment-cytokine fusion proteins may be used for enhancing *in vitro* cytotoxicity (for instance, that mediated by monoclonal antibodies against tumor  
20 targets) and for enhancing *in vivo* killing of target tissues (for example, blood and bone marrow cancers). See, generally, Hornick et al., *Blood* 89:4437-4447, 1997). In general, cytokines are toxic if administered systemically. The described fusion proteins enable targeting of a cytokine to a desired site of action, such as a cell having binding sites for zveg4, thereby providing an elevated local concentration of cytokine. Suitable  
25 cytokines for this purpose include, for example, interleukin-2 and granulocyte-macrophage colony-stimulating factor (GM-CSF). Such fusion proteins may be used to cause cytokine-induced killing of tumors and other tissues undergoing angiogenesis or neovascularization.

In yet another embodiment, a zveg4 polypeptide or anti-zveg4 antibody  
30 can be conjugated with a radionuclide, particularly with a beta-emitting or gamma-emitting radionuclide, and used to reduce restenosis. For instance, iridium-192 impregnated ribbons placed into stented vessels of patients until the required radiation dose was delivered resulted in decreased tissue growth in the vessel and greater luminal diameter than the control group, which received placebo ribbons. Further,  
35 revascularisation and stent thrombosis were significantly lower in the treatment group. Similar results are predicted with targeting of a bioactive conjugate containing a radionuclide, as described herein.

The bioactive polypeptide or antibody conjugates described herein can be delivered intravenously, intra-arterially or intraductally, or may be introduced locally at the intended site of action.

Polynucleotides encoding zveg4 polypeptides are useful within gene therapy applications where it is desired to increase or inhibit zveg4 activity. For example, Isner et al., *The Lancet* (ibid.) reported that VEGF gene therapy promoted blood vessel growth in an ischemic limb. Additional applications of zveg4 gene therapy include stimulation of wound healing, repopulation of vascular grafts, stimulation of neurite growth, and inhibition of cancer growth and metastasis. Gene delivery systems useful in this regard include adenovirus, adeno-associated virus, and naked DNA vectors.

The present invention also provides polynucleotide reagents for diagnostic use. For example, a zveg4 gene, a probe comprising zveg4 DNA or RNA, or a subsequence thereof can be used to determine if a mutation has occurred at the zveg4 locus on human chromosome 11. Detectable chromosomal aberrations at the zveg4 gene locus include, but are not limited to, aneuploidy, gene copy number changes, insertions, deletions, restriction site changes and rearrangements. Such aberrations can be detected using polynucleotides of the present invention by employing molecular genetic techniques, such as restriction fragment length polymorphism (RFLP) analysis, short tandem repeat (STR) analysis employing PCR techniques, and other genetic linkage analysis techniques known in the art (Sambrook et al., *ibid.*; Ausubel et. al., *ibid.*; A.J. Marian, *Chest* 108:255-265, 1995).

The invention is further illustrated by the following non-limiting examples.

## EXAMPLES

### Example 1

Human Multiple Tissue Northern Blots I, II, III and Human RNA Master  
5 Blots (Clontech Laboratories, Inc., Palo Alto, CA) were probed to determine the tissue  
expression of zveg4. Blots were prehybridized for 3 hours at 65 degrees in 10 ml of a  
hybridization solution (ExpressHyb™ Hybridization Solution; Clontech Laboratories,  
Inc.) containing 1 mg of salmon sperm DNA that had been boiled 5 minutes, then iced  
1 minute. The probe used was a 251-bp PCR fragment generated with 20 pmole each  
10 of primers ZC21,119 (SEQ ID NO:25) and ZC21,120 (SEQ ID NO:26), and 5 µl of a  
heart cDNA library prepared from heart RNA using a commercially available kit  
(Marathon™ cDNA Amplification Kit from Clontech Laboratories, Inc.). The reaction  
was run as follows: 94 degrees for 1 minute; then 30 cycles of 94 degrees, 20 seconds;  
67 degrees, 1 minute; and ended with a 5-minute incubation at 72 degrees. The PCR  
15 product was gel-purified, and the DNA was eluted from the gel slab with a spin column  
containing a silica gel membrane (QIAquick™ Gel Extraction Kit; Qiagen, Inc.,  
Valencia, CA).

51 ng of the resulting zveg4 fragment was labeled with <sup>32</sup>P using a  
commercially available kit (Rediprime™ II random-prime labeling system; Amersham  
20 Pharmacia Biotech, Piscataway, N.J.). Unincorporated radioactivity was removed with  
a push column (NucTrap® column; Stratagene, La Jolla, CA; see U.S. Patent No.  
5,336,412). 10 x 10<sup>6</sup> cpm of the resulting labeled probe and 1 mg of salmon sperm  
DNA were boiled 5 minutes, iced 1 minute, then mixed with 10 ml hybridization  
solution (ExpressHyb™) and added to blots. Hybridization took place overnight at 65  
25 degrees, followed by a wash in 2 x SSC, 0.1% SDS at room temperature, followed by a  
wash in 0.1 x SSC, 0.1% SDS at 50 degrees. Blots were exposed to film at -80 degrees  
overnight.

There was an approximately 4.4 kb transcript in every tissue except bone  
marrow. Heart, pancreas, stomach and adrenal gland showed the strongest zveg4  
30 expression on the Northern blots, and the dot blot additionally showed strong  
expression in the pituitary gland and the ovary.

### Example 2

Zveg4 was identified from the sequence of a clone from a human  
35 chronic myelogenous leukemia cell (K562) library by its homology to the VEGF  
family. Additional sequence was elucidated from a long sequence read of a clone from  
a pituitary library. An antisense expressed sequence tag (EST) for zveg4 was found,

for which its 5' partner was identified. This 5' EST (EST448186; GenBank) appeared to contain the 5' untranslated sequence for zveg4. A primer was designed from EST448186 to close the gap in the sequence. 20 pm each of ZC21,987 (SEQ ID NO:27) and ZC21,120 (SEQ ID NO:26) and 1.93 µg of a thyroid library were used in the PCR reaction. It was a modified PCR reaction using 5% DMSO and 1/10 volume of a commercial reagent (GC-Melt™; Clontech Laboratories, Inc.). The reaction was run for 1 minute at 94 degrees; then 30 cycles of 94 degrees, 20 seconds; 67 degrees, 1 minute; then a final 5-minute incubation at 72 degrees. A resulting 833-bp product was sequenced and found to be a zveg4 fragment containing the remainder of the coding sequence with an initiation MET codon, upstream stop codon, and 5' untranslated sequence. The composite sequence included an open reading frame of 1,110 bp (SEQ ID NO:1).

### Example 3

To make transgenic animals expressing zveg4 genes requires adult, fertile males (studs) (B6C3f1, 2-8 months of age (Taconic Farms, Germantown, NY)), vasectomized males (duds) (B6D2f1, 2-8 months, (Taconic Farms)), prepubescent fertile females (donors) (B6C3f1, 4-5 weeks, (Taconic Farms)) and adult fertile females (recipients) (B6D2f1, 2-4 months, (Taconic Farms)).

The donors are acclimated for 1 week, then injected with approximately 8 IU/mouse of Pregnant Mare's Serum gonadotrophin (Sigma, St. Louis, MO) I.P., and 46-47 hours later, 8 IU/mouse of human Chorionic Gonadotropin (hCG (Sigma)) I.P. to induce superovulation. Donors are mated with studs subsequent to hormone injections. Ovulation generally occurs within 13 hours of hCG injection. Copulation is confirmed by the presence of a vaginal plug the morning following mating.

Fertilized eggs are collected under a surgical scope (Leica MZ12 Stereo Microscope; Leica, Wetzlar, Germany). The oviducts are collected and eggs are released into urinanalysis slides containing hyaluronidase (Sigma Chemical Co.). Eggs are washed once in hyaluronidase, and twice in Whitten's W640 medium (Table 7; all reagents available from Sigma Chemical Co.) that has been incubated with 5% CO<sub>2</sub>, 5% O<sub>2</sub>, and 90% N<sub>2</sub> at 37°C. The eggs are stored in a 37°C/5% CO<sub>2</sub> incubator until microinjection.

Table 7

	<u>mgs/200 ml</u>	<u>mgs/500 ml</u>
NaCl	1280	3200
KCl	72	180
KH <sub>2</sub> PO <sub>4</sub>	32	80
MgSO <sub>4</sub> · 7H <sub>2</sub> O	60	150
Glucose	200	500
Ca <sup>2+</sup> Lactate	106	265
Benzylpenicillin	15	37.5
Streptomycin SO <sub>4</sub>	10	25
NaHCO <sub>3</sub>	380	950
Na Pyruvate	5	12.5
H <sub>2</sub> O	200 ml	500 ml
500 mM EDTA	100 µl	250 µl
5% Phenol Red	200 µl	500 µl
BSA	600	1500

Zveg4 cDNA is inserted into the expression vector pHB12-8 (see Fig. 2). Vector pHB12-8 was derived from p2999B4 (Palmiter et al., *Mol. Cell Biol.* 13:5266-5275, 1993) by insertion of a rat insulin II intron (ca. 200 bp) and polylinker (Fse I/Pme I/Asc I) into the Nru I site. The vector comprises a mouse metallothionein (MT-1) promoter (ca. 750 bp) and human growth hormone (hGH) untranslated region and polyadenylation signal (ca. 650 bp) flanked by 10 kb of MT-1 5' flanking sequence and 7 kb of MT-1 3' flanking sequence. The cDNA is inserted between the insulin II and hGH sequences.

10-20 micrograms of plasmid DNA is linearized, gel-purified, and resuspended in 10 mM Tris pH 7.4, 0.25 mM EDTA pH 8.0, at a final concentration of 5-10 nanograms per microliter for microinjection.

Plasmid DNA is microinjected into harvested eggs contained in a drop of W640 medium overlaid by warm, CO<sub>2</sub>-equilibrated mineral oil. The DNA is drawn into an injection needle (pulled from a 0.75mm ID, 1mm OD borosilicate glass capillary) and injected into individual eggs. Each egg is penetrated with the injection needle into one or both of the haploid pronuclei.

Picoliters of DNA are injected into the pronuclei, and the injection needle is withdrawn without coming into contact with the nucleoli. The procedure is repeated until all the eggs are injected. Successfully microinjected eggs are transferred



into an organ tissue-culture dish with pregassed W640 medium for storage overnight in a 37°C/5% CO<sub>2</sub> incubator.

The following day, 2-cell embryos are transferred into pseudopregnant recipients. The recipients are identified by the presence of copulation plugs, after  
5 copulating with vasectomized duds. Recipients are anesthetized and shaved on the dorsal left side and transferred to a surgical microscope. A small incision is made in the skin and through the muscle wall in the middle of the abdominal area outlined by the ribcage, the saddle, and the hind leg, midway between knee and spleen. The reproductive organs are exteriorized onto a small surgical drape. The fat pad is  
10 stretched out over the surgical drape, and a baby serrefine (Roboz, Rockville, MD) is attached to the fat pad and left hanging over the back of the mouse, preventing the organs from sliding back in.

With a fine transfer pipette containing mineral oil followed by alternating W640 and air bubbles, 12-17 healthy 2-cell embryos from the previous day's  
15 injection are transferred into the recipient. The swollen ampulla is located, and, holding the oviduct between the ampulla and the bursa, a nick in the oviduct is made with a 28 g needle close to the bursa, making sure not to tear the ampulla or the bursa.

The pipette is transferred into the nick in the oviduct, and the embryos are blown in, allowing the first air bubble to escape the pipette. The fat pad is gently  
20 pushed into the peritoneum, and the reproductive organs are allowed to slide in. The peritoneal wall is closed with one suture, and the skin is closed with a wound clip. The mice recuperate on a 37°C slide warmer for a minimum of 4 hours.

The recipients are returned to cages in pairs, and allowed 19-21 days gestation. After birth, 19-21 days postpartum is allowed before weaning. The  
25 weanlings are sexed and placed into separate sex cages, and a 0.5 cm biopsy (used for genotyping) is snipped off the tail with clean scissors.

Genomic DNA is prepared from the tail snips using a commercially available kit (DNeasy™ 96 Tissue Kit; Qiagen, Valencia, CA) following the manufacturer's instructions. Genomic DNA is analyzed by PCR using primers  
30 designed to the human growth hormone (hGH) 3' UTR portion of the transgenic vector. The use of a region unique to the human sequence (identified from an alignment of the human and mouse growth hormone 3' UTR DNA sequences) ensures that the PCR reaction does not amplify the mouse sequence. Primers ZC17,251 (SEQ ID NO:28) and ZC17,252 (SEQ ID NO:29) amplify a 368-base-pair fragment of hGH. In addition,  
35 primers ZC17,156 (SEQ ID NO:30) and ZC17,157 (SEQ ID NO:31), which hybridize to vector sequences and amplify the cDNA insert, may be used along with the hGH primers. In these experiments, DNA from animals positive for the transgene will

generate two bands, a 368-base-pair band corresponding to the hGH 3' UTR fragment and a band of variable size corresponding to the cDNA insert.

Once animals are confirmed to be transgenic (TG), they are back-crossed into an inbred strain by placing a TG female with a wild-type male, or a TG male with one or two wild-type female(s). As pups are born and weaned, the sexes are separated, and their tails snipped for genotyping.

To check for expression of a transgene in a live animal, a partial hepatectomy is performed. A surgical prep is made of the upper abdomen directly below the xiphoid process. Using sterile technique, a small 1.5-2 cm incision is made below the sternum, and the left lateral lobe of the liver is exteriorized. Using 4-0 silk, a tie is made around the lower lobe securing it outside the body cavity. An atraumatic clamp is used to hold the tie while a second loop of absorbable Dexon (American Cyanamid, Wayne, N.J.) is placed proximal to the first tie. A distal cut is made from the Dexon tie, and approximately 100 mg of the excised liver tissue is placed in a sterile petri dish. The excised liver section is transferred to a 14-ml polypropylene round bottom tube, snap frozen in liquid nitrogen, and stored on dry ice. The surgical site is closed with suture and wound clips, and the animal's cage is placed on a 37°C heating pad for 24 hours post-operatively. The animal is checked daily post-operatively, and the wound clips are removed 7-10 days after surgery.

Analysis of the mRNA expression level of each transgene is done using an RNA solution hybridization assay or real-time PCR on an ABI Prism 7700 (PE Applied Biosystems, Inc., Foster City, CA) following the manufacturer's instructions.

#### Example 4

An expression plasmid containing all or part of a polynucleotide encoding zveg4 is constructed via homologous recombination. A fragment of zveg4 cDNA is isolated by PCR using the polynucleotide sequence of SEQ ID NO: 1 with flanking regions at the 5' and 3' ends corresponding to the vector sequences flanking the zveg4 insertion point. The primers for PCR each include from 5' to 3' end: 40 bp of flanking sequence from the vector and 17 bp corresponding to the amino and carboxyl termini from the open reading frame of zveg4.

Ten µl of the 100 µl PCR reaction is run on a 0.8% LMP agarose gel (Seaplaque GTG) with 1 x TBE buffer for analysis. The remaining 90 µl of PCR reaction is precipitated with the addition of 5 µl 1 M NaCl and 250 µl of absolute ethanol. The plasmid pZMP6, which has been cut with SmaI, is used for recombination with the PCR fragment. Plasmid pZMP6 was constructed from pZP9 (deposited at the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-

2209, under Accession No. 98668) with the yeast genetic elements taken from pRS316 (deposited at the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209, under Accession No. 77145), an internal ribosome entry site (IRES) element from poliovirus, and the extracellular domain of CD8 truncated at the C-terminal end of the transmembrane domain. pZMP6 is a mammalian expression vector containing an expression cassette having the mouse metallothionein-1 promoter, multiple restriction sites for insertion of coding sequences, a stop codon, and a human growth hormone terminator. The plasmid also contains an *E. coli* origin of replication; a mammalian selectable marker expression unit comprising an SV40 promoter, enhancer and origin of replication, a DHFR gene, and the SV40 terminator; as well as the URA3 and CEN-ARS sequences required for selection and replication in *S. cerevisiae*.

One hundred microliters of competent yeast cells (*S. cerevisiae*) are independently combined with 10  $\mu$ l of the various DNA mixtures from above and transferred to a 0.2-cm electroporation cuvette. The yeast/DNA mixtures are electropulsed at 0.75 kV (5 kV/cm),  $\infty$  ohms, 25  $\mu$ F. To each cuvette is added 600  $\mu$ l of 1.2 M sorbitol, and the yeast is plated in two 300- $\mu$ l aliquots onto two URA-D plates and incubated at 30°C. After about 48 hours, the Ura<sup>+</sup> yeast transformants from a single plate are resuspended in 1 ml H<sub>2</sub>O and spun briefly to pellet the yeast cells. The cell pellet is resuspended in 1 ml of lysis buffer (2% Triton X-100, 1% SDS, 100 mM NaCl, 10 mM Tris, pH 8.0, 1 mM EDTA). Five hundred microliters of the lysis mixture is added to an Eppendorf tube containing 300  $\mu$ l acid-washed glass beads and 200  $\mu$ l phenol-chloroform, vortexed for 1 minute intervals two or three times, and spun for 5 minutes in an Eppendorf centrifuge at maximum speed. Three hundred microliters of the aqueous phase is transferred to a fresh tube, and the DNA is precipitated with 600  $\mu$ l ethanol (EtOH), followed by centrifugation for 10 minutes at 4°C. The DNA pellet is resuspended in 10  $\mu$ l H<sub>2</sub>O.

Transformation of electrocompetent *E. coli* host cells (Electromax DH10B™ cells; obtained from Life Technologies, Inc., Gaithersburg, MD) is done with 0.5-2 ml yeast DNA prep and 40  $\mu$ l of cells. The cells are electropulsed at 1.7 kV, 25  $\mu$ F, and 400 ohms. Following electroporation, 1 ml SOC (2% Bacto™ Tryptone (Difco, Detroit, MI), 0.5% yeast extract (Difco), 10 mM NaCl, 2.5 mM KCl, 10 mM MgCl<sub>2</sub>, 10 mM MgSO<sub>4</sub>, 20 mM glucose) is plated in 250- $\mu$ l aliquots on four LB AMP plates (LB broth (Lennox), 1.8% Bacto™ Agar (Difco), 100 mg/L Ampicillin).

Individual clones harboring the correct expression construct for zveg4 are identified by restriction digest to verify the presence of the zveg4 insert and to confirm that the various DNA sequences have been joined correctly to one another.

The inserts of positive clones are subjected to sequence analysis. Larger scale plasmid DNA is isolated using a commercially available kit (QIAGEN Plasmid Maxi Kit, Qiagen, Valencia, CA) according to manufacturer's instructions. The correct construct is designated zveg4/pZMP6.

5

#### Example 5

CHO DG44 cells (Chasin et al., *Som. Cell. Molec. Genet.* 12:555-566, 1986) are plated in 10-cm tissue culture dishes and allowed to grow to approximately 50% to 70% confluency overnight at 37°C, 5% CO<sub>2</sub>, in Ham's F12/FBS media (Ham's F12 medium, Life Technologies), 5% fetal bovine serum (Hyclone, Logan, UT), 1% L-glutamine (JRH Biosciences, Lenexa, KS), 1% sodium pyruvate (Life Technologies). The cells are then transfected with the plasmid zveg4/pZMP6 by liposome-mediated transfection using a 3:1 (w/w) liposome formulation of the polycationic lipid 2,3-dioleoyloxy-N-[2(sperminecarboxamido)ethyl]-N,N-dimethyl-1-propaniminium-

15 trifluoroacetate and the neutral lipid dioleoyl phosphatidylethanolamine in membrane-filtered water (Lipofectamine™ Reagent, Life Technologies), in serum free (SF) media formulation (Ham's F12, 10 mg/ml transferrin, 5 mg/ml insulin, 2 mg/ml fetuin, 1% L-glutamine and 1% sodium pyruvate). Zveg4/pZMP6 is diluted into 15-ml tubes to a total final volume of 640 µl with SF media. 35 µl of Lipofectamine™ is mixed with

20 605 µl of SF medium. The Lipofectamine™ mixture is added to the DNA mixture and allowed to incubate approximately 30 minutes at room temperature. Five ml of SF media is added to the DNA:Lipofectamine™ mixture. The cells are rinsed once with 5 ml of SF media, aspirated, and the DNA:Lipofectamine™ mixture is added. The cells are incubated at 37°C for five hours, then 6.4 ml of Ham's F12/10% FBS, 1% PSN

25 media is added to each plate. The plates are incubated at 37°C overnight, and the DNA:Lipofectamine™ mixture is replaced with fresh 5% FBS/Ham's media the next day. On day 3 post-transfection, the cells are split into T-175 flasks in growth medium. On day 7 post-transfection, the cells are stained with FITC-anti-CD8 monoclonal antibody (PharMingen, San Diego, CA) followed by anti-FITC-conjugated magnetic

30 beads (Miltenyi Biotec, Auburn, CA). The CD8-positive cells are separated using commercially available columns (MiniMACS Separation Unit; Miltenyi Biotec) according to the manufacturer's directions and put into DMEM/Ham's F12/5% FBS without nucleosides but with 50 nM methotrexate (selection medium).

Cells are plated for subcloning at a density of 0.5, 1 and 5 cells per well

35 in 96-well dishes in selection medium and allowed to grow out for approximately two weeks. The wells are checked for evaporation of medium and brought back to 200 µl

per well as necessary during this process. When a large percentage of the colonies in the plate are near confluency, 100  $\mu$ l of medium is collected from each well for analysis by dot blot, and the cells are fed with fresh selection medium. The supernatant is applied to a nitrocellulose filter in a dot blot apparatus, and the filter is treated at 100°C in a vacuum oven to denature the protein. The filter is incubated in 625 mM Tris-glycine, pH 9.1, 5mM  $\beta$ -mercaptoethanol, at 65°C, 10 minutes, then in 2.5% non-fat dry milk Western A Buffer (0.25% gelatin, 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 5 mM EDTA, 0.05% Igepal CA-630) overnight at 4°C on a rotating shaker. The filter is incubated with the anti-CD8 antibody-HRP conjugate in 2.5% non-fat dry milk Western A buffer for 1 hour at room temperature on a rotating shaker. The filter is then washed three times at room temperature in PBS plus 0.01% Tween 20, 15 minutes per wash. The filter is developed with chemiluminescence reagents (ECL™ direct labeling kit; Amersham Corp., Arlington Heights, IL) according to the manufacturer's directions and exposed to film (Hyperfilm ECL, Amersham) for approximately 5 minutes. Positive clones are trypsinized from the 96-well dish and transferred to 6-well dishes in selection medium for scaleup and analysis by Western blot.

#### Example 6

The protein coding region of zveg4 is amplified by PCR using primers that add FseI and AscI restriction sites at the 5' and 3' termini, respectively. PCR primers are used with a template containing the full-length zveg4 cDNA in a PCR reaction as follows: one cycle at 95°C for 5 minutes; followed by 15 cycles at 95°C for 1 min., 58°C for 1 min., and 72°C for 1.5 min.; followed by 72°C for 7 min.; followed by a 4°C soak. The PCR reaction product is loaded onto a 1.2 % (low melt) (SeaPlaque GTG™; FMC, Rockland, ME) gel in TAE buffer. The zveg4 PCR product is excised from the gel and purified using a spin column containing a silica gel membrane (QIAquick™ Gel Extraction Kit; Qiagen, Inc., Valencia, CA) as per kit instructions. The PCR product is then digested, phenol/chloroform extracted, EtOH precipitated, and rehydrated in 20ml TE (Tris/EDTA pH 8). The zveg4 fragment is then ligated into the cloning sites of the transgenic vector pHB12-8 and transformed into *E. coli* host cells (Electromax DH10B™ cells; obtained from Life Technologies, Inc., Gaithersburg, MD) by electroporation. Clones containing zveg4 DNA are identified by restriction analysis. A positive clone is confirmed by direct sequencing.

The zveg4 cDNA is released from the pTG12-8 vector using FseI and AscI enzymes. The cDNA is isolated on a 1% low melt agarose gel, and is then excised from the gel. The gel slice is melted at 70°C, extracted twice with an equal volume of Tris buffered phenol, and EtOH precipitated. The DNA is resuspended in 10 $\mu$ l H<sub>2</sub>O.

The zveg4 cDNA is cloned into the FseI-AscI sites of a modified pAdTrack CMV (He et al., *Proc. Natl. Acad. Sci. USA* 95:2509-2514, 1998). This construct contains a GFP marker gene. The CMV promoter driving GFP expression has been replaced with the SV40 promoter, and the SV40 polyadenylation signal has been replaced with the human growth hormone polyadenylation signal. In addition, the native polylinker has been replaced with FseI, EcoRV, and AscI sites. This modified form of pAdTrack CMV was named pZyTrack. Ligation is performed using a DNA ligation and screening kit (Fast-Link™; Epicentre Technologies, Madison, WI). In order to linearize the plasmid, approximately 5 µg of the pZyTrack zveg4 plasmid is digested with PmeI. Approximately 1 µg of the linearized plasmid is cotransformed with 200ng of supercoiled pAdEasy (He et al., *ibid.*) into BJ5183 cells. The co-transformation is done using a Bio-Rad Gene Pulser at 2.5kV, 200 ohms and 25 µF. The entire co-transformation is plated on 4 LB plates containing 25 µg/ml kanamycin. The smallest colonies are picked and expanded in LB/kanamycin, and recombinant adenovirus DNA identified by standard DNA miniprep procedures. Digestion of the recombinant adenovirus DNA with FseI-AscI confirms the presence of zveg4 DNA. The recombinant adenovirus miniprep DNA is transformed into *E. coli* DH10B competent cells, and DNA is prepared therefrom.

Approximately 5 µg of recombinant adenoviral DNA is digested with PacI enzyme (New England Biolabs) for 3 hours at 37°C in a reaction volume of 100 µl containing 20-30U of PacI. The digested DNA is extracted twice with an equal volume of phenol/chloroform and precipitated with ethanol. The DNA pellet is resuspended in 10 µl distilled water. A T25 flask of QBI-293A cells (Quantum Biotechnologies, Inc., Montreal, Canada), inoculated the day before and grown to 60-70% confluence, are transfected with the PacI digested DNA. The PacI-digested DNA is diluted up to a total volume of 50 µl with sterile HBS (150 mM NaCl, 20 mM HEPES). In a separate tube, 20 µl of 1mg/ml N-[1-(2,3-Dioleoyloxy)propyl]-N,N,N-trimethyl-ammonium methylsulfate (DOTAP; Boehringer Mannheim) is diluted to a total volume of 100 µl with HBS. The DNA is added to the DOTAP, mixed gently by pipeting up and down, and left at room temperature for 15 minutes. The media is removed from the 293A cells and washed with 5 ml serum-free MEM-alpha (Life Technologies, Gaithersburg, MD) containing 1 mM sodium pyruvate (Life Technologies), 0.1 mM MEM non-essential amino acids (Life Technologies) and 25 mM HEPES buffer (Life Technologies). 5 ml of serum-free MEM is added to the 293A cells and held at 37°C. The DNA/lipid mixture is added drop-wise to the T25 flask of 293A cells, mixed gently, and incubated at 37°C for 4 hours. After 4 hours the media containing the DNA/lipid mixture is aspirated off and replaced with 5 ml complete MEM containing

5% fetal bovine serum. The transfected cells are monitored for Green Fluorescent Protein (GFP) expression and formation of foci (viral plaques).

Seven days after transfection of 293A cells with the recombinant adenoviral DNA, the cells expressing the GFP protein start to form foci. These foci are  
5 viral "plaques" and the crude viral lysate is collected by using a cell scraper to collect all of the 293A cells. The lysate is transferred to a 50ml conical tube. To release most of the virus particles from the cells, three freeze/thaw cycles are done in a dry ice/ethanol bath and a 37°C waterbath.

Ten 10-cm plates of nearly confluent (80-90%) 293A cells are set up 20  
10 hours prior to infection. The crude lysate is amplified (primary amplification) to obtain a working "stock" of zvegf4 rAdV lysate. 200 ml of crude rAdV lysate is added to each 10-cm plate, and the plates are monitored for 48 to 72 hours looking for cytopathic effect (CPE) under the white light microscope and expression of GFP under the fluorescent microscope. When all of the 293A cells show CPE, this 1° stock lysate is  
15 collected, and freeze/thaw cycles performed as described above.

Secondary (2°) amplification of zvegf4 rAdV is obtained from twenty 15-cm tissue culture dishes of 80-90% confluent 293A cells. All but 20 ml of 5% MEM media is removed, and each dish is inoculated with 300-500 ml of 1° amplified rAdV lysate. After 48 hours the 293A cells are lysed from virus production, the lysate is  
20 collected into 250 ml polypropylene centrifuge bottles, and the rAdV is purified.

NP-40 detergent is added to a final concentration of 0.5% to the bottles of crude lysate to lyse all cells. Bottles are placed on a rotating platform for 10 minutes and agitated as fast as possible. The debris is pelleted by centrifugation at 20,000 X G for 15 minutes. The supernatant is transferred to 250-ml polycarbonate centrifuge  
25 bottles, and 0.5 volume of 20% PEG8000/2.5M NaCl solution is added. The bottles are shaken overnight on ice. The bottles are centrifuged at 20,000 X G for 15 minutes, and the supernatants are discarded into a bleach solution. A white precipitate (precipitated virus/PEG) forms in two vertical lines along the walls of the bottles on either side of the spin mark. Using a sterile cell scraper, the precipitate from 2 bottles is resuspended in  
30 2.5 ml PBS. The virus solution is placed in 2-ml microcentrifuge tubes and centrifuged at 14,000 X G in a microcentrifuge for 10 minutes to remove any additional cell debris. The supernatants from the 2-ml microcentrifuge tubes are transferred into a 15-ml polypropylene snapcap tube and adjusted to a density of 1.34 g/ml with CsCl. The volume of the virus solution is estimated, and 0.55 g/ml of CsCl added. The CsCl is  
35 dissolved, and 1 ml of this solution weighed. The solution is transferred to polycarbonate, thick-walled, 3.2 ml centrifuge tubes (Beckman) and spun at 348,000 X

G for 3-4 hours at 25°C. The virus forms a white band. Using wide-bore pipette tips, the virus band is collected.

The virus from the gradient will have a large amount of CsCl, which must be removed before it can be used on cells. Pharmacia PD-10 columns prepacked with Sephadex® G-25M (Pharmacia) are used to desalt the virus preparation. The column is equilibrated with 20 ml of PBS. The virus is loaded and allowed to run into the column. 5 ml of PBS is added to the column, and fractions of 8-10 drops collected. The optical density of 1:50 dilutions of each fraction is determined at 260 nm on a spectrophotometer, and a clear absorbance peak is identified. These fractions are pooled, and the optical density (OD) of a 1:25 dilution is determined. OD is converted into virus concentration using the formula  $(OD \text{ at } 260\text{nm})(25)(1.1 \times 10^{12}) = \text{virions/ml}$ .

To store the virus, glycerol is added to the purified virus to a final concentration of 15%, mixed gently and stored in aliquots at -80°C.

A protocol developed by Quantum Biotechnologies, Inc. (Montreal, Canada) is followed to measure recombinant virus infectivity. Briefly, two 96-well tissue culture plates are seeded with  $1 \times 10^4$  293A cells per well in MEM containing 2% fetal bovine serum for each recombinant virus to be assayed. After 24 hours, 10-fold dilutions of each virus from  $1 \times 10^{-2}$  to  $1 \times 10^{-14}$  are made in MEM containing 2% fetal bovine serum. 100µl of each dilution is placed in each of 20 wells. After 5 days at 37°C, wells are read either positive or negative for CPE and PFU/ml is calculated.

TCID<sub>50</sub> formulation used is as per Quantum Biotechnologies, Inc., above. The titer (T) is determined from a plate where virus used is diluted from  $10^{-2}$  to  $10^{-14}$ , and read 5 days after the infection. At each dilution a ratio (R) of positive wells for CPE per the total number of wells is determined. The titer of the undiluted sample is  $T = 10^{(1+F)} = \text{TCID}_{50}/\text{ml}$ , where  $F = 1+d(S-0.5)$ , S is the sum of the ratios (R), and d is Log<sub>10</sub> of the dilution series (e.g., d = 1 for a ten-fold dilution series). To convert TCID<sub>50</sub>/ml to pfu/ml, 0.7 is subtracted from the exponent in the calculation for titer (T).

#### Example 7

Recombinant zveg4 having a carboxyl-terminal Glu-Glu affinity tag was produced in a baculovirus expression system according to conventional methods. The culture was harvested, and the cells were lysed with a solution of 0.02 M Tris-HCl, pH 8.3, 1 mM EDTA, 1 mM DTT, 1 mM 4-(2-Aminoethyl)-benzenesulfonyl fluoride hydrochloride (Pefabloc® SC; Boehringer-Mannheim), 0.5 µM aprotinin, 4 mM leupeptin, 4 mM E-64, 1% NP-40 at 4°C for 15 minutes on a rotator. The solution was centrifuged, and the supernatant was recovered. Twenty ml of extract was combined with 50 µl of anti-Glu-Glu antibody conjugated to Sepharose® beads in 50 µl buffer.



The mixture was incubated on a rotator at 4°C overnight. The beads were recovered by centrifugation and washed 3 x 15 minutes at 4°C. Pellets were combined with sample buffer containing reducing agent and heated at 98°C for five minutes. The protein was analyzed by polyacrylamide gel electrophoresis under reducing conditions followed by western blotting on a PVDF membrane using an antibody to the affinity tag. Two bands were detected, one a  $M_r \approx 49$  kD and the other at  $M_r \approx 21$  kD. Sequence analysis showed the larger band to comprise two sequences, one beginning at Arg-19 of SEQ ID NO:2 and the other beginning at Asn-35 of SEQ ID NO:2. The asparagine residue appeared to have been deamidated to an aspartic acid. The smaller band began at Ser-250 of SEQ ID NO:2.

#### Example 8

The *zveg4* cDNA was cloned into the EcoRV-AscI sites of a modified pAdTrack-CMV (He et al., *Proc. Natl. Acad. Sci. USA* 95:2509-2514, 1998). This construct contains the green fluorescent protein (GFP) marker gene. The CMV promoter driving GFP expression was replaced with the SV40 promoter, and the SV40 polyadenylation signal was replaced with the human growth hormone polyadenylation signal. In addition, the native polylinker was replaced with FseI, EcoRV, and AscI sites. This modified form of pAdTrack-CMV was named pZyTrack. Ligation was performed using a commercially available DNA ligation and screening kit (Fast-Link™ kit; Epicentre Technologies, Madison, WI).

*Zveg4* was assayed in an aortic ring outgrowth assay (Nicosia and Ottinetti, *ibid.*; Villaschi and Nicosia, *ibid.*). Thoracic aortas were isolated from 1-2 month old SD male rats and transferred to petri dishes containing HANK's buffered salt solution. The aortas were flushed with additional HANK's buffered salt solution to remove blood, and adventitial tissue surrounding the aorta was carefully removed. Cleaned aortas were transferred to petri dishes containing EBM basal media, serum free (Clonetics, San Diego, CA). Aortic rings were obtained by slicing approximately 1-mm sections using a scalpel blade. The ends of the aortas used to hold the aorta in place were not used. The rings were rinsed in fresh EBM basal media and placed individually in a wells of a 24-well plate coated with basement membrane matrix (Matrigel®; Becton Dickinson, Franklin Lakes, NJ). The rings were overlaid with an additional 50 µl of the matrix solution and placed at 37°C for 30 minutes to allow the matrix to gel. Test samples were diluted in EBM basal serum-free media supplemented with 100 units/ml penicillin, 100 µg/ml streptomycin and HEPES buffer and added at 1 ml/well. Background control was EBM basal serum-free media alone. Basic FGF (R&D Systems, Minneapolis, MN) at 20 ng/ml was used as a positive control. *Zveg4*

adenovirus was added to wells, assuming a cell count of 500,000 cells and a multiplicity of infection of 5000 particles/cell. A null adenovirus (designated "zPar") was used as a control. Samples were added in a minimum of quadruplets. Rings were incubated for 5-7 days at 37°C and analyzed for growth. Aortic outgrowth was scored by multiple, blinded observers using 0 as no growth and 4 as maximum growth. Zveg4 adenovirus produced a significant increase in outgrowth, comparable to the most potent control (bFGF).

#### Example 9

10 Polyclonal anti-peptide antibodies were prepared by immunizing 2 female New Zealand white rabbits with the peptides huzveg4-1 (CGHKEVPPRIKSRTNQIK; SEQ ID NO:39), huzveg4-2 (ESWQEDLENMYLDTPRYRGRSYHDC; SEQ ID NO:40), or huzveg4-3 (CFEPGHIKRRGRAKTMALVDIQLD; SEQ ID NO:41). The peptides were synthesized using an Applied Biosystems Model 431A peptide synthesizer (Applied Biosystems, Inc., Foster City, CA) according to the manufacturer's instructions. The peptides were conjugated to keyhole limpet hemocyanin (KLH) with maleimide activation. The rabbits were each given an initial intraperitoneal (ip) injection of 200 µg of peptide in Complete Freund's Adjuvant followed by booster ip injections of 100 µg peptide in Incomplete Freund's Adjuvant every three weeks. Seven to ten days after the administration of the second booster injection (3 total injections), the animals were bled, and the sera were collected. The animals were then boosted and bled every three weeks.

25 The zveg4 peptide-specific rabbit sera were characterized by an ELISA titer check using 1 µg/ml of the peptide used to make the antibody as an antibody target. The 2 rabbit sera to the huzveg4-1 peptide had titer to their specific peptide at a dilution of 1:5,000,000. The 2 rabbit sera to the huzveg4-2 peptide had titer to their specific peptide at a dilution of 1:5,000,000. The 2 rabbit sera to the huzveg4-3 peptide had titer to their specific peptide at a dilution of 1:500,000.

30 The zveg4 peptide-specific polyclonal antibodies were affinity purified from the sera using CNBr-SEPHAROSE 4B protein columns (Pharmacia LKB) that were prepared using 10 mg of the specific peptide per gram CNBr-SEPHAROSE, followed by 20X dialysis in PBS overnight. Zveg4-specific antibodies were characterized by an ELISA titer check using 1 µg/ml of the appropriate peptide antigens as antibody targets. The lower limit of detection (LLD) of the anti-huzveg4-1 affinity purified antibody on its specific antigen (huzveg4-1 peptide) was a dilution of 0.1

pg/ml. The LLD of the anti-huzveg4-2 affinity purified antibody on its specific antigen (huzveg4-2 peptide) was a dilution of 5 ng/ml. The LLD of the rabbit anti-huzveg4-3 affinity purified antibody on its specific antigen (huzveg4-3 peptide) was a dilution of 5 ng/ml.

5

#### Example 10

Recombinant carboxyl-terminal Glu-Glu tagged zveg4 (zveg4-cee) was produced from recombinant baculovirus-infected insect cells. Two-liter cultures were harvested, and the media were sterile-filtered using a 0.2 µm filter.

10 Protein was purified from the conditioned media by a combination of anti-Glu-Glu (anti-EE) peptide antibody affinity chromatography and S-200 gel exclusion chromatography. Culture media (pH 6.0, conductivity 7 mS) was directly loaded onto a 20 x 80 mm (25-ml bed volume) anti-EE antibody affinity column at a flow of 6 ml/minute. The column was washed with ten column volumes of PBS, then  
15 bound protein was eluted with two column volumes of 0.4 mg/ml EYMPTD peptide (SEQ ID NO:42) (Princeton BioMolecules Corp., Langhorne, PA). Five-ml fractions were collected. Samples from the anti-EE antibody affinity column were analyzed by SDS-PAGE with silver staining and western blotting (as disclosed below) for the presence of zveg4-cee. Zveg4-cee-containing fractions were pooled and concentrated  
20 to 3.8 ml by filtration using a Biomax™ -5 concentrator (Millipore Corp., Bedford, MA), and loaded onto a 16 x 1000 mm gel filtration column (Sephacryl™ S-200 HR; Amersham Pharmacia Biotech, Piscataway, NJ). The fractions containing purified zveg4-cee were pooled, filtered through a 0.2 µm filter, aliquoted into 100 µl each, and frozen at -80°C. The concentration of the final purified protein was determined by  
25 colorimetric assay (BCA assay reagents; Pierce, Rockford, IL) and HPLC-amino acid analysis.

Recombinant zveg4-cee was analyzed by SDS-PAGE (NuPAGE™ 4-12% gel; Novex, San Diego, CA) with silver staining (FASTsilver™, Geno Technology, Inc., Maplewood, MO) and Western blotting using antibodies to the huzveg4-1,  
30 huzveg4-2, and huzveg4-3 peptides, and anti-EE antibody. Either the conditioned media or purified protein was electrophoresed using an electrophoresis mini-cell (XCell II™ mini-cell; Novex, San Diego, CA) and transferred to nitrocellulose (0.2 µm; Bio-Rad Laboratories, Hercules, CA) at room temperature using an XCell II™ blot module (Novex) with stirring according to directions provided in the instrument manual. The  
35 transfer was run at 500 mA for one hour in a buffer containing 25 mM Tris base, 200 mM glycine, and 20% methanol. The filters were then blocked with 10% non-fat dry

milk in PBS for 10 minutes at room temperature. The nitrocellulose was quickly rinsed, then primary antibody was added in PBS containing 2.5% non-fat dry milk. The blots were incubated for two hours at room temperature or overnight at 4°C with gentle shaking. Following the incubation, blots were washed three times for 10 minutes each  
5 in PBS. Secondary antibody (goat anti-rabbit IgG conjugated to horseradish peroxidase; obtained from Rockland Inc., Gilbertsville, PA) diluted 1:2000 in PBS containing 2.5% non-fat dry milk was added, and the blots were incubated for two hours at room temperature with gentle shaking. The blots were then washed three times, 10 minutes each, in PBS, then quickly rinsed in H<sub>2</sub>O. The blots were developed  
10 using commercially available chemiluminescent substrate reagents (SuperSignal® ULTRA reagents 1 and 2 mixed 1:1; reagents obtained from Pierce), and the signal was captured using image analysis software (Lumi-Imager™ Lumi Analyst 3.0; Roche Molecular Biochemicals, Indianapolis, IN) for times ranging from 10 seconds to 5 minutes or as necessary.

15 The purified zvefg4-ccc appeared as a single band at about 85 kDa under non-reducing conditions with silver staining, but at about 50 kDa under reducing conditions, suggesting a dimeric form of zvefg4-ccc under non-reducing conditions.

Using either 4-1, 4-3 or anti-EE antibody, the purified zvefg4-ccc showed the same result as silver staining gel; the 4-3 antibody gave a much weaker  
20 signal. However, in addition to recognizing the 85-kDa band under non-reducing conditions and the 50-kDa band under reducing conditions, the 4-2 antibody recognized two bands at 35 kDa and 32 kDa under non-reducing conditions, and two bands at 38 kDa and 35 kDa under reducing conditions. While not wishing to be bound by theory, the smaller bands are likely to be cleaved forms of zvefg4-ccc missing the N-  
25 terminal portion of the protein that is recognized by the 4-1 antibody.

#### Example 11

Recombinant zvegf4 was analyzed for mitogenic activity on rat liver stellate cells (obtained from N. Fausto, University of Washington), human aortic  
30 smooth muscle cells (Clonetics Corp., Walkersville, MD), human retinal pericytes (Clonetics Corp.) and human hepatic fibroblasts (Clonetics Corp.). Test samples consisted of conditioned media (CM) from adenovirally infected HaCaT human keratinocyte cells (Boukamp et al., *J. Cell. Biol.* 106:761-771, 1988; Skobe and Fusenig, *Proc. Natl. Acad. Sci. USA* 95:1050-1055, 1998; obtained from Dr. Norbert E.  
35 Fusenig, Deutsches Krebsforschungszentrum, Heidelberg, Germany) expressing full length zvegf-4. Control CM was generated from HaCaT cells infected with a parental GFP-expressing adenovirus (zPar). The CM were concentrated 10-fold using a 15 ml

centrifugal filter device with a 10K membrane filter (Ultrafree®; Millipore Corp., Bedford, MA), then diluted back to 1x with ITS media (serum-free DMEM/Ham's F-12 medium containing 5 µg/ml insulin, 20 µg/ml transferrin, and 16 pg/ml selenium). Cells were plated at a density of 2,000 cells/well in 96-well culture plates and grown for approximately 72 hours in DMEM containing 10% fetal calf serum at 37°C. Cells were quiesced by incubating them for approximately 20 hours in serum-free DMEM/Ham's F-12 medium containing insulin (5 µg/ml), transferrin (20 µg/ml), and selenium (16 pg/ml) (ITS). At the time of the assay, the medium was removed, and test samples were added to the wells in triplicate. For measurement of [<sup>3</sup>H]thymidine incorporation, 20 µl of a 50 µCi/ml stock in DMEM was added directly to the cells, for a final activity of 1 µCi/well. After another 24-hour incubation, media were removed and cells were incubated with 0.1 ml of trypsin until cells detached. Cells were harvested onto 96-well filter plates using a sample harvester (FilterMate™ harvester; Packard Instrument Co., Meriden, CT). The plates were then dried at 65°C for 15 minutes, sealed after adding 40 µl/well scintillation cocktail (Microscint™ O; Packard Instrument Co.) and counted on a microplate scintillation counter (Topcount®; Packard Instrument Co.). Results, presented in Table 8, demonstrated that zveg4 CM had approximately 7-fold higher mitogenic activity than control CM on pericytes cells and approximately a 1.5 - 2.4-fold higher mitogenic activity on the other cell types tested.

Table 8

Sample	CPM incorporated			
	Zveg4 (1x CM)		zPar (1xCM )	
	Mean	St. dev.	Mean	St. dev.
Human retinal pericytes	3621	223	523	306
Human hepatic fibroblasts	7757	753	3232	264
Human aortic SMC	2009	37	1263	51
Rat liver stellate cells	34707	1411	14413	1939

Example 12

Recombinant, C terminal glu-glu tagged zveg4 was analyzed for mitogenic activity on human aortic smooth muscle cells (HAoSMC) (Clonetics), human retinal pericytes (Clonetics) and human aortic adventitial fibroblasts (AoAF) (Clonetics). Cells were plated at a density of 2,000 cells/well in 96-well culture plates and grown for approximately 72 hours in DMEM containing 10% fetal calf serum at 37°C. Cells were quiesced by incubating them for 20 hours in ITS medium. At the

time of the assay, the medium was removed, and test samples were added to the wells in triplicate. Test samples consisted of purified, full-length, tagged zveg4 expressed in baculovirus-infected cells. Purified protein in a buffer containing 0.1% BSA was serially diluted into ITS medium at concentrations of 1 µg/ml to 1 ng/ml and added to the test plate. A control buffer of 0.1% BSA was diluted identically to the highest concentration of zveg4 protein and added to the plate. For measurement of [3H]thymidine incorporation, 20 µl of a 50 µCi/ml stock in DMEM was added directly to the cells, for a final activity of 1 µCi/well. After another 24-hour incubation, mitogenic activity was assessed by measuring the uptake of [3H]thymidine. Media were removed, and cells were incubated with 0.1 ml of trypsin until cells detached. Cells were harvested onto 96-well filter plates using a sample harvester (FilterMate™ harvester; Packard Instrument Co., Meriden, CT). The plates were then dried at 65°C for 15 minutes, sealed after adding 40 µl/well scintillation cocktail (Microscint™ O; Packard Instrument Co.) and counted on a microplate scintillation counter (Topcount®; Packard Instrument Co.). Results, presented in Table 9, demonstrated that 80 ng/ml zveg4 had approximately 1.7-fold higher mitogenic activity on pericytes, 3.2-fold higher activity on aortic SMCs, and 2.6-fold higher activity on aortic fibroblasts as compared to the buffer control.

20

Table 9

Sample	CPM Incorporated					
	Pericytes		HAoSMC		AoAF	
	Mean	St. dev.	Mean	St. dev.	Mean	St. dev.
Zveg4, 80 ng/ml	96.7	18.2	488.7	29.6	177.0	1.0
Zveg4, 20 ng/ml	81.7	11.7	211.7	50.8	107.7	20.1
Zveg4, 5 ng/ml	67.3	6.7	191.7	4.5	123.7	10.5
Buffer control	58.7	8.5	152.3	40.1	68.7	8.3

Example 13

Mice (C57BL6) were infected with a zveg4-encoding adenovirus vector (AdZyveg4) to determine the effects on serum chemistry, complete blood counts (CBC), body and organ weight changes, and histology. On day -1, the mice were tagged, individually weighed, and group normalized for separation into treatment groups (4 mice per cage). Group 1 mice (n=8 females, 7 males) received GFP (control)

adenovirus,  $1 \times 10^{11}$  particles. Group 2 mice (n=8 females, 6 males) received zveg4 adenovirus,  $1 \times 10^{11}$  particles. Group 3 mice (n=8 females, 8 males) were untreated controls. On day 0, the mice received injections of the appropriate adenovirus solution. On day 10, blood was collected (under ether anesthesia) for CBCs and clinical chemistry measurements. On day 20, mice were weighed and sacrificed by cervical dislocation after collecting blood (under ether anesthesia) for CBCs and clinical chemistry measurements. Tissues were collected for histopathology. Observations were as follows:

10 Serum chemistry changes: AdZyveg4 treated mice were hypoglycemic. This effect increased in magnitude over time (day 10 vs. day 20). Serum cholesterol levels were significantly increased (2-fold) at both time points. Serum levels of albumin and the enzymes ALT, AST and alkaline phosphatase were all significantly increased in AdZyveg4 treated mice. Serum calcium and total bilirubin were also significantly increased, and became more elevated over time.

15 CBC changes: AdZyveg4-treated mice had significantly higher lymphocyte count at both time points (mean >10). Platelet counts were significantly lower at day 20. Red blood cell count was significantly higher in females at day 10, significantly higher in males at day 20.

20 Body/organ weights: AdZyveg4-treated males lost weight over the course of the experiment. This result was significantly different than control animals, which gained weight. There was no difference among the female mice; all groups gained similar weight. Spleen weight was significantly greater (approximately 4-fold) in all AdZyveg4-treated mice. Liver weight was also significantly greater in all AdZyveg4-treated mice. There was no significant difference in kidney weight between groups.

25 Histology: In the liver, proliferation of sinusoidal endothelial cells was observed. In the spleen, proliferation of reticuloendothelial cells was observed. In the kidney, proliferative glomerulopathy was observed. While not wishing to be bound by theory, this glomerulopathy may have been due to proliferation of capillary endothelial cells. In the femurs, there was proliferation of endosteal bone (mostly in trabecular bone), which in some cases replaced most of the bone marrow. Proliferation of stromal cells was also observed in bone. In the lung, there was increased frequency of brochoaveolar lymphoid tissue.

#### 35 Example 14

90  $\mu$ g of recombinant zveg4 protein was dissolved in 500  $\mu$ l PBS containing 2 mCi Na-  $^{125}$ I (Amersham Corp.). One derivatized, nonporous polystyrene

bead (ODO-Beads®; Pierce, Rockford, IL) was added, and the reaction mixture was incubated one minute on ice. The iodinated protein was separated from unincorporated  $^{125}\text{I}$  by gel filtration using an elution buffer of 10% acetic acid, 150 mM NaCl, and 0.25% gelatin. The active fraction contained 29  $\mu\text{g/ml}$   $^{125}\text{I}$ -zveg4 with a specific activity of  $3.0 \times 10^4$  cpm/ng.

The following cell lines were plated into the wells of a 24-well tissue culture dish and cultured in growth medium for three days:

1. Human retinal pericytes, passage 6 (pericytes).
2. Rat stellate cells, passage 8.
- 10 3. Human umbilical vein endothelial cells, passage 4 (HUVEC).
4. Human aortic adventitial fibroblasts, passage 5 (AoAF).
5. Human aortic smooth muscle cells, passage 2 (AoSMC).

Cells were washed once with ice-cold binding buffer (HAM'S F-12 containing 2.5 mg/ml BSA, 20 mM HEPES, pH 7.2), then 250  $\mu\text{l}$  of the following solutions was added to each of three wells of the culture dishes containing the test cells. Binding solutions were prepared in 5 mL of binding buffer with 250 pM  $^{125}\text{I}$ -zveg4 and:

1. No addition.
2. 25 nM zveg4.
3. 25 nM zveg3.
- 20 4. 25 nM PDGF-AA.
5. 25 nM PDGF-BB

The reaction mixtures were incubated on ice for 2 hours, then washed three times with one ml of ice-cold binding buffer. The bound  $^{125}\text{I}$ -zveg4 was quantitated by gamma counting a Triton-X 100 extract of the cells.

Results, shown in Table 10, indicate binding of zveg4 to pericytes, stellate cells, AoAF, and AoSMC, but not to HUVEC. The first column represents total CPM  $^{125}\text{I}$ -zveg4 bound/well. The second column is  $^{125}\text{I}$ -zveg4 bound/well when blocked with cold ligand. The difference between the two numbers represents specific binding.

30



Table 10

Cell Type	<sup>125</sup> I-zveg4 Bound (CPM)	<sup>125</sup> I-zveg4 Bound w/cold zveg4 (CPM)
1. Pericytes	3083 +/- 864	623 +/- 60
2. Stellate Cells	2131 +/- 450	413 +/- 164
3. HUVEC	485 +/- 91	227 +/- 13
4. AoAF	1544 +/- 131	300 +/- 15
5. AoSMC	1628 +/- 203	440 +/- 46

Example 15

The zveg4 gene was mapped to human chromosome 11 using the commercially available version of the Stanford G3 Radiation Hybrid Mapping Panel (Research Genetics, Inc., Huntsville, AL). This panel contains PCRable DNAs from each of 83 radiation hybrid clones of the whole human genome, plus two control DNAs (the RM donor and the A3 recipient). A publicly available WWW server (<http://shgc-www.stanford.edu>) allows chromosomal localization of markers. 20- $\mu$ l reaction mixtures were set up in a PCRable 96-well microtiter plate (Stratagene, La Jolla, CA) and used in a thermal cycler (RoboCycler® Gradient 96; Stratagene). Each of the 85 PCR mixtures consisted of 2  $\mu$ l buffer (10X KlenTaq PCR reaction buffer, Clontech Laboratories, Inc., Palo Alto, CA), 1.6  $\mu$ l dNTPs mix (2.5 mM each, PERKIN-ELMER, Foster City, CA), 1  $\mu$ l sense primer, ZC22,685 (SEQ ID NO:37), 1  $\mu$ l antisense primer, ZC22,686 (SEQ ID NO:38), 2  $\mu$ l of a density increasing agent and tracking dye (RediLoad, Research Genetics, Inc., Huntsville, AL), 0.4  $\mu$ l of a commercially available DNA polymerase/antibody mix (50X Advantage™ KlenTaq Polymerase Mix, obtained from Clontech Laboratories, Inc., Palo Alto, CA), 25 ng of DNA from an individual hybrid clone or control, and x  $\mu$ l ddH<sub>2</sub>O for a total volume of 20  $\mu$ l. The reaction mixtures were overlaid with an equal amount of mineral oil and sealed. The PCR cycler conditions were an initial 5-minute denaturation at 94°C; 35 cycles of 45 seconds denaturation at 94°C, 45 seconds annealing at 64°C. and 75 seconds extension at 72°C; followed by a final extension for 7 minutes at 72°C. The reaction products were separated by electrophoresis on a 2% agarose gel. The results showed linkage of zveg4 to the human chromosome 11 framework marker SHGC-34226 with a LOD score of 14.90 and at a distance of 0 cR\_10000 from the marker. The use of surrounding genes/markers positions Zveg4 in the 11q22.3-q23.1 chromosomal region.

Example 16

The structure of recombinant zveg4 was analyzed by Western blotting using conventional techniques. Protein produced in the HaCaT human keratinocyte cell line was electrophoresed under reducing and non-reducing conditions, transferred to  
5 filters, and probed with antibodies to the interdomain and CUB domain regions of the protein. Reduced protein appeared as a single band having an apparent  $M_r$  of approximately 53 kD, consistent with a glycosylated, monomeric protein. Non-reduced protein appeared as a single band having an apparent  $M_r$  of approximately 85 kD, consistent with a disulfide-linked dimer.

10

Example 17

An expression plasmid containing full-length zveg4 was constructed, using the expression vector pEZE2. pEZE2 is derived from pDC312 by the addition of additional restriction enzyme recognition sites to the multiple cloning site. pDC312  
15 and pEZE2 contain an EASE segment, as described in WO 97/25420, which can improve expression of recombinant proteins two to eight fold in mammalian cells, preferably Chinese Hamster Ovary (CHO) cells. The pEZE2 expression unit contains the CMV enhancer/promoter, the adenovirus tripartite leader sequence, a multiple cloning site for insertion of the coding region for the recombinant protein, an  
20 encephalomyocarditis virus internal ribosome entry site, a coding segment for mouse dihydrofolate reductase, and the SV40 transcription terminator. In addition, pEZE2 contains an *E. coli* origin of replication and a bacterial beta-lactamase gene.

A zveg4 DNA fragment was generated by PCR (Advantage2 PCR Kit, Clontech, Palo Alto, CA) with 5' FseI and 3' AscI sites for direct cloning into the  
25 expression vector. The 5' primer contained an FseI site, Kozak sequence, and the first 21 basepairs of the native leader sequence for zveg4 (ZC26,136; SEQ ID NO:43). The 3' primer contained the last 21 basepairs of zveg4, a stop codon, and an AscI site (ZC26,137; SEQ ID NO:44). The PCR reaction included 1 :L of template (ESTEP plasmid zveg4pcrf1#3) and was run as follows: 94°C, 1 minute, 1 cycle; then 25 cycles  
30 of 94°C, 30 seconds; 55°C, 30 seconds; 68°C, 1 minute; then a final extension cycle of 72°C for 7 minutes.

The ESTEP plasmid zveg4pcrf1#3 contains the full-length human zveg4 fragment. This fragment was generated by PCR using 20 pm each of ZC22,341 (SEQ ID NO:45) and ZC22,342 (SEQ ID NO:46) primers and 3 :L of a thyroid library.  
35 The reaction was run as follows: 94°C, 1 minute, 1 cycle; then 30 cycles of 94°C, 20 seconds; 66°C, 1.5 minutes; then a final extension cycle of 72°C for 5 minutes. The 1,272 bp product was gel purified on a 1% TBE gel, and the DNA was extracted from

the gel slab using the QIAquick Gel Extraction Kit (Qiagen, Valencia, CA). This 1,272 bp fragment was subcloned into pCR2.1 vector (Invitrogen, Carlsbad, CA), and designated zveg4pcrf1#3.

The PCR generated fragment was purified (Qiaquick PCR clean-up kit, Qiagen, Valencia, CA) and digested with restriction enzymes AscI and FseI (New England Biolabs, Beverly, MA) in a single 100 :L reaction. Five micrograms of the expression vector pEZE2 were also digested with FseI and AscI in a single 100 :L reaction. The digested DNA was fractionated by agarose gel electrophoresis and the DNA fragments were isolated and purified (Qiaquick Gel Extraction Kit, Qiagen).

Five microliters of the zveg4 DNA fragment and 1 :L of the pEZE2 vector fragment were ligated overnight at room temperature (New England Biolabs High Concentrated Ligase and supplied buffer). One microliter of the ligation reaction was added to 25 :L of electrocompetant *E. coli* strain DH10B (Life Technologies) in a 0.2 cm cuvette. The mixture was electroporated (BioRad *E. coli* Pulser) at 2.3 kv. To the cuvette, 1 mL of LB broth was added, and 100 :L of the mix was plated onto LB/Ampicillin agar plates. The plates were incubated overnight at 37°C, and 8 isolated colonies were picked for DNA mini prep (Qiaquick Mini-Prep Kit, Qiagen). Individual clones were screened by PCR for the presence of zveg4 DNA, using the above-mentioned primers. DNA sequencing was performed on clones #1-6, to verify the correct full-length sequence. One clone contained the correct expected sequence and a Maxi prep of DNA was made (Qiagen Plasmid Maxi Kit, Qiagen).

CHO DG44 (Chasin et al., *Som. Cell. Molec. Genet.* 12:555-666, 1986) were plated and allowed to grow to approximately 50% to 70% confluency over night at 37°C in MEM alpha media (JRH Biosciences, Lenexa, KS), 7.5% fetal bovine serum (Hyclone, Logan, UT), 1% L-glutamine (Life Technologies), 1% sodium pyruvate (Life Technologies), 1% HT solution (Life Technologies), and 1% Penicillin/Streptomycin (Life Technologies). The cells were then transfected with the plasmid pEZE2/zveg4 by liposome-mediated transfection, using a 10:1 (w/w) liposome formulation of the polycationic lipid dioctaldecylamidoglycyl spermine, in serum-free (SF) medium formulation DMEM/F12 – Life Technologies, Non-Essential Amino Acids-Life Technologies, 1% L-glutamine, 1% sodium pyruvate. The plasmid pEZE2/zveg4 was diluted in a final volume of 500 :L of SF medium in a 15 mL conical tube, and 20 :L of Transfectam (Promega, Madison, WI) reagent was added, mixed well and incubated at room temperature for 10 minutes. After incubation, 4.5 mL of SF medium was added to the DNA mixture and mixed well using a 5 mL pipette. The cells were rinsed 3 times with SF medium, and the 5 mL of DNA solution was overlayed upon the cell monolayer. The cells were incubated at 37°C, 5% CO<sub>2</sub> for 2 hours. Then 6 mL of

complete medium (MEM alpha, 7.5% FBS, 1% L-glutamine, 1% sodium pyruvate, 1% HT, 1% Pen/Strep) and the cells were incubated for a further 48 hours. After 48 hours, the cells were trypsinized from the plate with 1 mL of 0.25% Trypsin/ 1 mM EDTA (Life Technologies) and quenched with 4 mL of complete medium without nucleosides  
5 (MEM alpha, 7.5% Dialysed FBS, 1% L-glutamine, 1% sodium pyruvate, 1% Pen/Strep). Five hundred microliters of the cell suspension were transferred to plates containing 10 mL of complete medium without nucleosides. The cultures were grown for 14 days, until single colonies that were approximately 0.25 cm in diameter were present. Cloning rings (Bellco Glass, Inc., Vineland, NJ) were used to isolate 24 single  
10 colonies, which were removed with trypsin, transferred to 6 well cell cluster plates (Costar, Corning, NY), and incubated 4 days.

The cell wells were rinsed with SF medium and 2 mL of SF medium was added, and the culture was incubated for 24 hours. The conditioned SF medium was concentrated approximately 20-fold using a 10K centrifuge device (Millipore  
15 Corporation, Bedford, MA). Twenty-five microliters of the concentrate was added to 15  $\mu$ L of 4X Sample Buffer (Novex, San Diego, CA) with 50 mM beta-Mercaptoethanol, and the mixture was run on a 4-12% NuPAGE gel (Novex). The proteins from the gel were transferred to nitrocellulose membranes (Novex) and the blot was blocked with 10% non-fat dry milk in Western A (0.25% gelatin, 50 mM Tris-HCl  
20 pH 7.4, 150 mM NaCl, 5 mM EDTA, 0.05% Igepal CA-630) overnight at room temperature on a rotating shaker platform. The membrane was rinsed 3 times in Western A. An antibody to the N-terminus of the zveg4 protein was diluted at 1:3000 in 50 mL 5% non-fat milk in Western A. The antibody solution was overlayed on the membrane and incubated at room temperature on a rocking platform for 1 hour. After  
25 the 1 hour incubation, the solution was discarded and the membrane rinsed 3 times with Western A and once with Western B (50 mM Tris pH 7.4, 5 mM EDTA, 0.05% Igepal CA-630, 1 M NaCl, 0.25 % Gelatin). The secondary antibody, an F(ab')<sub>2</sub> fragment of Donkey-Anti-Rabbit-HRP (Amersham Corp., Arlington Heights, IL), was diluted in Western A at 1:3000, overlayed on the membrane, and incubated 1 hour at room  
30 temperature on a rocking platform. The secondary antibody solution was discarded, and the membrane was washed 3 times in Western A and 3 times in Western B. Chemiluminescence was used to detect the full-length or protease-digested N-terminus of zveg4 according to the manufacturer's instructions (Pierce, Rockford, IL), and was analysed by LumiAnalyser (Roche/Boehringer Mannheim, Mannheim, Germany). Four  
35 of the 12 clones were positive for zveg4, and numbers 7 and 12 were trypsinized and transferred to T175 flasks (Costar, Corning, NY) in complete medium without nucleosides.

Example 18

An expression construct encoding the growth factor domain of zveg4 is prepared. A PCR fragment was generated (Clontech Advantage 2 PCR Kit) that  
5 contained a 5' BamHI restriction site, an N-terminal EE tag, and zveg4 amino acid residues 258-381 (stop codon included). The 5' oligo primer contains the BamHI site, an N-terminal EE tag sequence, and zveg4 basepairs corresponding to the N-terminus of the growth factor domain (ZC27,116; SEQ ID NO:47). The 3' oligo primer contains the last 21 basepairs of zveg4 (stop codon included) (ZC27,137; SEQ ID NO:48). The  
10 expression vector pZMP20 was used, which contains the CMV immediate early promoter, a consensus intron from the variable region of mouse immunoglobulin heavy chain locus, Kozak sequences, an optimized t-PA secretory signal sequence (U.S. Patent No. 5,641,655), multiple restriction sites for insertion of coding sequences, a stop codon, and a human growth hormone terminator. The plasmid also contains an  
15 IRES element from poliovirus, the extracellular domain of CD8 truncated at the C-terminal end of the transmembrane domain, an *E. coli* origin of replication, a DHFR gene, the SV40 terminator, and the URA3 and CEN-ARS sequences required for replication in *S. cerevisiae*. The resulting plasmid is designated pZMP20/GFD.NEE. A 504 basepair fragment with a 5' FseI site and 3' AscI site is isolated from this plasmid  
20 for ligation into the pEZE2 expression vector (5' FseI, 3' AscI) for expression in CHO DG44 cells.

Example 19A. Mouse genomic library screen

25 A partial mouse zveg4 sequence was obtained by probing a mouse genomic library with a human zveg4 restriction digest fragment containing the entire coding sequence. The probe was generated by digesting 8 :g of a full-length human zveg4 plasmid with EcoRI (Gibco BRL, Gaithersburg, MD). The 1,289 bp fragment was gel purified on a 2.3% TBE gel and the cDNA was extracted from the agarose slab  
30 using the QIAquick Gel Extraction Kit (Qiagen). The mouse genomic library was an emb13 SP6/T7 lambda BamHI cloned library (Clontech, Palo Alto, CA) plated on a K802 host lawn on 24 NZY plates, and represented  $7.2 \times 10^5$  pfus.

Twenty four filter lifts were prehybridized in EXPRESSHYB solution (Clontech) containing 0.1 mg/ml salmon sperm DNA which had been boiled 5 minutes,  
35 then iced. Hybridization took place overnight at 50°C. Sixty three ng of the human fragment mentioned above were labeled with  $^{32}$ P using the Rediprime II Random Prime Labeling System (Amersham Pharmacia, Buckinghamshire, England).

Unincorporated radioactivity was removed using a NucTrap push column (Stratagene, La Jolla, CA). Filters were hybridized in EXPRESSHYB solution containing  $1.0 \times 10^6$  cpm/ml zveg4 probe, 0.1 mg/ml salmon sperm DNA, and 0.5 :g/ml murine cot-1 DNA which had been boiled 5 minutes, then iced. Hybridization took place overnight at 50°C. Filter lifts were washed in 2 x SSC, 0.1% SDS at room temperature for 2 hours, then the temperature was raised to 60°C for one hour. Overnight exposure at -80°C showed 7 putative primary hits.

A K802 host culture was prepared to plate the primary hits for a secondary screen. The 7 primary hits were picked with a Pasteur pipet and eluted in 1 ml SM (0.1 M NaCl, 50 mM Tris pH 7.5, 10 mM MgSO<sub>4</sub>, 0.02% gelatin) with a few drops of chloroform overnight at 4°C. After plating to determine titers, 10 times the number of plaques in the original pfu were plated on NZY maxi plates with 10 mM MgSO<sub>4</sub> /NZY top agarose and a lawn of K802 cells for four of the primary hits and grown overnight at 37°C. Lifts were done using Hybond-N filters (Amersham Pharmacia). The filters were marked for orientation with a hot needle, denatured in 1.5 M NaCl and 0.5 M NaOH for 10 minutes, then neutralized in 1.5 M NaCl and 0.5 M Tris-HCl pH 7.2 for 10 minutes. The DNA was affixed to the filter using a STRATALINKER UV crosslinker (Stratagene, La Jolla, CA) at 1200 joules, and prewashed at 65°C in prewash buffer consisting of 0.25 x SSC, 0.25% SDS and 1 mM EDTA, changing solution three times for a total of 45 minutes to remove cell debris. Five lifts were put in each vial, three vials total. Each vial of lifts was prehybridized overnight at 50°C in 13 ml of EXPRESSHYB Hybridization Solution (Clontech) mixed with 1.3 mg salmon sperm DNA which had been boiled 5 minutes, then iced.

Sixty three ng of the human zveg4 fragment was labeled for a probe as described above. Each vial of filters was hybridized in 9 ml of EXPRESSHYB Hybridization Solution mixed with 0.99 to  $1.1 \times 10^6$  human zveg4 probe, 0.5 :g/ml murine cot-1 DNA, and 0.9 mg/ml salmon sperm DNA which had been boiled 5 minutes, then iced. Hybridization took place overnight at 50°C. Wash conditions described above for the primary screen were repeated for this secondary screen. Two of the 4 primary putative hits that were tested came up positive after an overnight exposure at -80°C.

Isolated plaques #7c1 and #18b2 were eluted in 200 :l SM overnight at 4°C, and fresh host K802 cells were prepared. Serial dilutions ranging from  $10^{-2}$  to  $10^{-3}$  were plated to obtain a titer estimate. Only #18b2 gave any plaques (for a titer of 2.6 to  $3.0 \times 10^3$  phage per :l), and this plaque was further pursued. Two plates with  $10^5$  pfus per plate were prepared for a phage DNA prep from plate lysates. Plates were grown at 37°C for 6 hours, until the phage were starting to get confluent, and then 12

ml of SM per plate was added to elute the phage overnight at 4°C. At this point, plates were shaken at room temperature one hour, the supernatant was removed, 1% chloroform was added, and supernatant was shaken for 15 minutes. The DNA was prepped using the Wizard Lambda Preps DNA Purification System (Promega), sections  
5 IV and VI.

Plaque #18b2 DNA was cut with several restriction enzymes to generate fragments to run on a Southern gel. Digests were run on a 1% TBE agarose gel. The gel was soaked in 0.25 M HCl for 30 minutes, rinsed in distilled H<sub>2</sub>O, soaked in 0.5 M NaOH and 1.5 M NaCl for 40 minutes with one solution change, and neutralized in 1.5  
10 M NaCl and 0.5 M Tris-HCl (pH 7.2) for 40 minutes with one solution change. A TURBOBLOTTER Rapid Downward Transfer System (Schleicher & Schuell, Keene, NH) was set up to transfer the DNA onto a Nytran/BA-S membrane (Schleicher & Schuell) overnight. The DNA was affixed to the Nytran using a STRATALINKER UV crosslinker (Stratagene) at 1200 joules. The blot was prehybridized overnight at 50°C  
15 in 12 ml EXPRESSHYB Hybridization Solution (Clontech) mixed with 1.2 mg salmon sperm DNA which had been boiled 5 minutes, then iced. Fifty nine ng of the human zveg4 fragment was labeled for a probe, as described above. Unincorporated radioactivity was removed by chromatography using a commercially available push column (NUCTRAP column, Stratagene). Ten ml of EXPRESSHYB Hybridization  
20 Solution was mixed with  $1.0 \times 10^6$  cpm/ml of human zveg4 probe, 0.5 :g/ml murine cot-1 DNA, and 0.1 mg/ml salmon sperm DNA which had been boiled 5 minutes, then iced, and then added to the blot. Hybridization took place overnight at 50°C. The blot was washed as described above, and exposed to film overnight at -80°C.

The Southern gel had a fragment from the BamH1/Pst1 digest which  
25 hybridized to the probe in the size range of 2.0 to 2.9 kb, which was pursued. Plaque 18b2 lambda DNA (2.8 :g) was cut with 20 units of BamH1 (Boehringer Mannheim, Indianapolis, IN), and 20 :l Pst1 (Life Technologies) for 2 hours at 37°C. The digest was run on a 1% TBE gel, and a 2.0 kb doublet, as well as 2.7 kb/2.9 kb bands, were excised from the gel. The DNA was extracted from the agarose using the Qiaquick Gel  
30 Extraction Kit (Qiagen). The 18b2 fragments were ligated into a pbluescriptIIKS+ vector (Stratagene) cut with BamH1, Pst1 and BamH1/Pst1. Three clones with a Pst1 insert, and 4 clones with a BamH1/Pst1 insert, from these ligations were digested with their respective insert site restriction enzymes for another Southern blot to determine which was the original hybridizing fragment. The 1% TBE gel was treated and the  
35 DNA was transferred to the Nytran blot as described above. The blot was prehybridized as above in 13 ml of hybridization solution. Fifty nine ng of the human zveg4 fragment was labeled and unincorporated radioactivity was removed as

described above. Human zveg4 probe ( $8.4 \times 10^5$ /ml cpm), 0.1 mg/ml of salmon sperm DNA, and 0.5 :g/ml of mouse cot-1 DNA were boiled 5 minutes, iced 1 minute, and mixed with 7 ml of EXPRESSHYB hybridization solution, then added to the blot. Hybridization took place overnight at 50°C. The same washing procedure was used as mentioned above. The blot was exposed to film for 3 hours at -80°C, and both 2.0 kb band inserts strongly hybridized to the probe. These clones were sequenced and found to contain part of the murine zveg4 cub domain. Primers were designed from this sequence for a PCR cDNA screen.

10                    B. PCR screen of mouse cDNA panel

A panel of available in house and commercial mouse cDNAs were screened with 20 pm each of ZG26,317 (SEQ ID NO:49) and ZG26,318 (SEQ ID NO:50) primers. The PCR reaction conditions were as follows: 94°C, 2 minutes; then 35 cycles of 94°C, 10 seconds; 65°C, 20 seconds; 72°C, 30 seconds; then ended with a 15 5 minute extension at 72°C. Embryo, salivary gland, neonatal skin and testis showed strong products of the predicted 200 bp size.

C. Full Length Mouse zveg4 Sequence

The in house mouse testis arrayed library representing  $9.6 \times 10^5$  clones was screened by PCR using primers ZG26,317 (SEQ ID NO:49) and ZG26,318 (SEQ ID NO:50) according to conditions specified above. This library was deconvoluted down to a positive pool of 250 clones. *E. coli* DH10B cells (Gibco BRL) were transformed with this pool by electroporation following the manufacturer's protocol. The transformed culture was titered and arrayed out to 96 wells at ~20 cells/well. The cells were grown up in LB+amp overnight at 37°C. An aliquot of the cells was pelleted and PCR was used to identify a positive pool. Thermocycler conditions were as described above. The remaining cells from a positive pool were plated, and colonies were screened by PCR to identify a positive clone. Sequence analysis indicated that this clone, named "zveg4mpzp7x-6", was incomplete at the 5' end and appeared to 30 contain an intron at the 5' end.

The mouse salivary gland library representing  $9.6 \times 10^5$  clones was then screened by PCR using primers ZG26,317 (SEQ ID NO:49) and ZG26,318 (SEQ ID NO:50) according to conditions specified above. The library was deconvoluted down to a positive pool of 250 clones. This 250 clonal pool was verified as having the 5' end by RACE. Twenty pm each of ZG26,318 (SEQ ID NO:50) and ZG14,063 (SEQ ID NO:51) primers and 3 :l of that pool was used. The reaction was run as follows: 94°C, 2 minutes, then 5 cycles of 94°C, 15 seconds; 70°C, 30 seconds; 30 cycles of 94°C, 15



seconds, 62°C, 20 seconds; 7°C, 30 seconds, and a final extension at 72°C for 7 minutes. The RACE product obtained upon sequencing confirmed that this pool contained the initiation Met. The same protocol as described above was carried out to isolate a single clone from the pool. Sequence analysis revealed that this clone, named  
5 "zveg4mpzp7x-7", had a 225 bp deletion in coding compared to clone #6 (bp 865 to bp 1079 in the final sequence).

The sequences derived from zveg4mpzp7x-6 and from zveg4mpzp7x-7 were combined to obtain a full-length mouse zveg4 polynucleotide sequence (SEQ ID NO:52) and mouse zveg4 polypeptide sequence (SEQ ID NO:53).

10

#### D. Full Length Mouse zveg4 Clone

The full length cDNA clone was generated by a two step ligation of fragments from clone #6 and clone #7 from above. An EcoR1/Hind3 three prime fragment was generated from clone #6 first. Nine :g of clone #6 were digested with 15  
15 units of EcoR1 (Gibco BRL, Gaithersburg, MD) and 15 units of Hind3 (Gibco BRL) for 2 hours at 37°C. The 528 bp fragment was gel purified on a 1% TBE gel, and the cDNA was extracted from the gel slab using the QIAquick Gel Extraction Kit (Qiagen). It was ligated into pbluescriptIIKS+ (Stratagene) digested with EcoR1 and Hind3. Three :g of a clone with this zveg4 insert was digested with 15 units of EcoR1 (Gibco  
20 BRL), gel purified on a 1% TBE gel, and the DNA was extracted using the kit mentioned above. The 5' EcoR1 zveg4 fragment from clone #7 was ligated into the EcoR1-digested clone mentioned above. This EcoR1 fragment was generated by digesting 8 :g of clone #7 with 30 units of EcoR1 (Gibco BRL) for 2 hours at 37°C. The 754 bp fragment was gel purified on a 1% TBE gel, and the DNA was extracted  
25 from the gel slab as mentioned above.

#### Example 20

##### A. Treatment of Naïve PC12 Cells with zveg4 Conditioned Medium

HaCat cells were infected with a null adenovirus (zPar) as a control, or  
30 with adenovirus expressing zveg4. Conditioned medium (CM) from these transfected cells was assayed for its ability to induce neurite outgrowth in the PC12 Pheochromocytoma cell line (see Banker and Goslin, in Culturing Nerve Cells, chapter 6, "Culture and experimental use of the PC12 rat Pheochromocytoma cell line"; also, see Rydel and Greene, J. Neuroscience 7(11): 3639-53, November 1987).

35 Briefly, PC12 cell cultures (ATCC# CRL 1721) were propagated with RPMI 1640 medium (Gibco/BRL, Gaithersburg, MD), 10% horse serum (Sigma, St. Louis, MO), and 5% fetal bovine serum (FBS; Hyclone, Logan, UT). Plastic culture

dishes (Beckton Dickinson, Bedford, MA) were coated with rat tail collagen type I, and PC12 cells were plated into 24 well plates at  $2 \times 10^4$  cells/ml in RPMI + 1% FBS and incubated overnight at 37°C in 5% CO<sub>2</sub>. The PC12 culture medium was then removed, and replaced with either zveg4-CM or control-CM added in 2-fold dilutions (starting at 5x dilution). Recombinant human NGF (R+D, Minneapolis, MN) was added as a positive control at concentrations of 100 or 30 ng/ml. As a negative control, CM of the null adenovirus (zpar) was used. To test for synergy of zveg4 and NGF, additional wells of PC12 cells were treated with zveg4-CM in combination with a suboptimal concentration of NGF (3 ng/ml). The culture medium was replaced every second day with RPMI + 1% FBS, until the total length of incubation reached 7 days.

The NGF-treated PC12 cells exhibited stable neurite outgrowth and neuronal differentiation. PC12 cells exposed to zveg4-CM exhibited morphological changes, such as cell flattening and the appearance of cells with short processes, suggesting differentiation into neuronal lineage. For PC12 cells incubated with a suboptimal dose of NGF plus zveg4-CM, an increase in a population of cells bearing short processes was observed.

#### B. Treatment of Primed, Neurite-Bearing PC12 Cells with zveg4 Conditioned Medium

Zveg4-CM and a control-CM (zpar) (as described in Example 20.A., above) were assayed for their ability to promote survival of differentiated PC12 neurons (see Banker and Goslin, *supra*, Rydel and Greene, *supra*).

Briefly, PC12 cells were maintained as described in Example 20.A., above, and were treated with appropriate doses of NGF to induce differentiation into cells that express the properties of post-mitotic sympathetic neurons. More specifically, PC12 cells were treated with recombinant human NGF (R+D, Minneapolis, MN) at a concentration of 50 ng/ml for 6 days, with a change of medium every other day. Cells were plated into 24 well plates overnight, and the culture medium was replaced with zveg4-CM or control-CM (in 2-fold dilutions, starting at 5x), or with NGF as a positive control (starting with 100 ng/ml in 3-fold dilutions).

Cultures were set up either with 1% FBS or serum-free culture (SF) medium. Cells were propagated over 9 days, with medium changes on every second day. Continuous treatment with NGF alone promoted the survival of the entire neuronal population and produced increasing neurite outgrowth. Zveg4-CM promoted the survival of a subpopulation of neurons, but did not induce additional neurite outgrowth. Cells cultured in control-CM degenerated.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

## CLAIMS

We claim:

1. An isolated polypeptide of at least 15 amino acid residues comprising an epitope-bearing portion of a protein of SEQ ID NO:2.
2. The polypeptide of claim 1 wherein said polypeptide comprises a segment that is at least 70% identical to a sequence selected from the group consisting of:
  - (a) residues 52-179 of SEQ ID NO:2; and
  - (b) residues 258-370 of SEQ ID NO:2.
3. The isolated polypeptide according to claim 1 wherein said polypeptide is selected from the group consisting of:
  - residues 19-179 of SEQ ID NO:2;
  - residues 52-179 of SEQ ID NO:2;
  - residues 19-253 of SEQ ID NO:2;
  - residues 52-253 of SEQ ID NO:2;
  - residues 19-255 of SEQ ID NO:2;
  - residues 52-255 of SEQ ID NO:2;
  - residues 19-257 of SEQ ID NO:2;
  - residues 52-257 of SEQ ID NO:2;
  - residues 19-253 of SEQ ID NO:2;
  - residues 52-253 of SEQ ID NO:2;
  - residues 19-370 of SEQ ID NO:2;
  - residues 52-370 of SEQ ID NO:2;
  - residues 180-370 of SEQ ID NO:2; and
  - residues 258-370 of SEQ ID NO:2.
4. An isolated polypeptide comprising a sequence of amino acids of the formula  $R1_x-R2_y-R3_z$ , wherein:
  - R1 is a polypeptide of from 100 to 130 residues in length, is at least 70% identical to residues 52-179 of SEQ ID NO:2, and comprises a sequence motif C[KR]Y[DNE][WYF]X{11,15}G[KR][WYF]C (SEQ ID NO:4) corresponding to residues 109-131 of SEQ ID NO:2;
  - R2 is a polypeptide at least 90% identical to residues 180-257 of SEQ ID NO:2;

R3 is a polypeptide at least 70% identical in amino acid sequence to residues 258-370 of SEQ ID NO:2 and comprises cysteine residues at positions corresponding to residues 272, 302, 306, 318, 360, and 362 of SEQ ID NO:2; a glycine residue at a position corresponding to residue 304 of SEQ ID NO:2; and a sequence motif CX{18,33}CXGXCX{6,33}CX{20,50}CXC (SEQ ID NO:3) corresponding to residues 272-362 of SEQ ID NO:2; and

each of x, y, and z is individually 0 or 1, subject to the limitations that:

at least one of x and z is 1; and

if x and z are each 1, then y is 1.

5. The isolated polypeptide of claim 4 wherein x=1.
6. The isolated polypeptide of claim 5 wherein R1 is at least 90% identical to residues 52-179 of SEQ ID NO:2.
7. The isolated polypeptide of claim 5 wherein R1 comprises residues 52-179 of SEQ ID NO:2.
8. The isolated polypeptide of claim 5 wherein R1 is at least 90% identical to residues 19-179 of SEQ ID NO:2.
9. The isolated polypeptide of claim 5 wherein y=1.
10. The isolated polypeptide of claim 9 wherein z=1.
11. The isolated polypeptide of claim 10 wherein R3 is at least 90% identical to residues 258-370 of SEQ ID NO:2.
12. The isolated polypeptide of claim 4 wherein said polypeptide comprises residues 52-253 of SEQ ID NO:2, residues 180-370 of SEQ ID NO:2, or residues 52-370 of SEQ ID NO:2.
13. The isolated polypeptide of claim 4 wherein z=1.
14. The isolated polypeptide of claim 13 wherein R3 is at least 90% identical to residues 258-370 of SEQ ID NO:2.

15. The isolated polypeptide of claim 13 wherein R3 comprises residues 258-370 of SEQ ID NO:2.

16. The isolated polypeptide of claim 13 wherein  $y=1$ .

17. The isolated polypeptide of claim 16 wherein  $x=1$  and R1 is at least 90% identical to residues 52-179 of SEQ ID NO:2.

18. The isolated polypeptide of claim 17 wherein R3 is at least 90% identical to residues 258-370 of SEQ ID NO:2.

19. The isolated polypeptide of claim 13, further comprising cysteine residues at positions corresponding to residues 308 and 316 of SEQ ID NO:2.

20. The isolated polypeptide of claim 4, further comprising an affinity tag.

21. The isolated polypeptide of claim 4, further comprising an immunoglobulin constant domain.

22. An isolated protein comprising a first polypeptide operably linked to a second polypeptide, wherein said first polypeptide comprises a sequence of amino acids of the formula  $R1_x-R2_y-R3_z$ , wherein:

R1 is a polypeptide of from 100 to 130 residues in length, is at least 70% identical to residues 52-179 of SEQ ID NO:2, and comprises a sequence motif  $C[KR]Y[DNE][WYF]X\{11,15\}G[KR][WYF]C$  (SEQ ID NO:4) corresponding to residues 109-131 of SEQ ID NO:2;

R2 is a polypeptide at least 90% identical to residues 180-257 of SEQ ID NO:2;

R3 is a polypeptide at least 70% identical in amino acid sequence to residues 258-370 of SEQ ID NO:2 and comprises cysteine residues at positions corresponding to residues 272, 302, 306, 318, 360, and 362 of SEQ ID NO:2; a glycine residue at a position corresponding to residue 304 of SEQ ID NO:2; and a sequence motif  $CX\{25,33\}CXGXCX\{10,33\}CX\{20,50\}CXC$  (SEQ ID NO:3) corresponding to residues 272-362 of SEQ ID NO:2; and

each of  $x$ ,  $y$ , and  $z$  is individually 0 or 1, subject to the limitations that:

at least one of  $x$  and  $z$  is 1; and

if  $x$  and  $z$  are each 1, then  $y$  is 1,

and wherein said protein modulates cell proliferation, apoptosis, differentiation, metabolism, or migration.

23. The isolated protein of claim 22 wherein said protein is a heterodimer.

24. The isolated protein of claim 23 wherein z is 1 and said second polypeptide is selected from the group consisting of VEGF, VEGF-B, VEGF-C, VEGF-D, zveg3, PlGF, PDGF-A, and PDGF-B.

25. The isolated protein of claim 23 wherein:

x=1, z=1, and said second polypeptide comprises residues 46-345 of SEQ ID NO:32;

x=1 and said second polypeptide comprises residues 46-170 of SEQ ID NO:32; or

z=1 and said second polypeptide comprises residues 235-345 of SEQ ID NO:32.

26. The isolated protein of claim 22 wherein said protein is a homodimer.

27. The isolated protein of claim 26 wherein z=1.

28. The isolated protein according to claim 27 wherein each of said first and second polypeptides comprises residues 258-370 of SEQ ID NO:2.

29. The isolated protein of claim 26 wherein x=1.

30. The isolated protein of claim 29 wherein each of said first and second polypeptides comprises residues 52-179 of SEQ ID NO:2.

31. An isolated protein produced by a method comprising:

(a) culturing a host cell containing an expression vector comprising the following operably linked elements:

a transcription promoter;

a DNA segment encoding a polypeptide comprising a sequence of amino acid residues selected from the group consisting of:

i) residues 52-370 of SEQ ID NO:2;

ii) residues 52-253 of SEQ ID NO:2;

iii) residues 180-370 of SEQ ID NO:2; and  
iii) residues 258-370 of SEQ ID NO:2; and  
a transcription terminator,  
under conditions whereby the DNA segment is expressed; and  
(b) recovering from the cell the protein product of expression of the DNA construct.

32. An isolated polynucleotide of up to approximately 4.4 kb in length, wherein said polynucleotide encodes a polypeptide comprising a sequence of amino acids of the formula  $R1_x-R2_y-R3_z$ , wherein:

R1 is a polypeptide of from 100 to 130 residues in length, is at least 70% identical to residues 52-179 of SEQ ID NO:2, and comprises a sequence motif C[KR]Y[DNE][WYF]X{11,15}G[KR][WYF]C (SEQ ID NO:4) corresponding to residues 109-131 of SEQ ID NO:2;

R2 is a polypeptide at least 90% identical to residues 180-257 of SEQ ID NO:2;

R3 is a polypeptide at least 70% identical in amino acid sequence to residues 258-370 of SEQ ID NO:2 and comprises cysteine residues at positions corresponding to residues 272, 302, 306, 318, 360, and 362 of SEQ ID NO:2; a glycine residue at a position corresponding to residue 304 of SEQ ID NO:2; and a sequence motif CX{25,33}CXGXCX{10,33}CX{20,50}CXC (SEQ ID NO:3) corresponding to residues 272-362 of SEQ ID NO:2; and

each of x, y, and z is individually 0 or 1, subject to the limitations that:

at least one of x and z is 1; and

if x and z are each 1, then y is 1.

33. The polynucleotide of claim 32, wherein said polynucleotide is DNA.

34. The polynucleotide of claim 33 comprising nucleotides 1 through 1110 of SEQ ID NO:6.

35. The polynucleotide of claim 33 comprising nucleotides 226 through 1335 of SEQ ID NO:1.

36. An expression vector comprising the following operably linked elements:

a transcription promoter;



a DNA polynucleotide according to claim 32; and  
a transcription terminator.

37. The expression vector of claim 36, further comprising a secretory signal sequence operably linked to the DNA polynucleotide.

38. A cultured cell into which has been introduced an expression vector according to claim 36, wherein said cell expresses the polypeptide encoded by the DNA polynucleotide.

39. A pharmaceutical composition comprising a protein according to claim 22 in combination with a pharmaceutically acceptable vehicle.

40. The pharmaceutical composition of claim 39 wherein  $x=1$ .

41. The pharmaceutical composition of claim 39 wherein  $z=1$ .

42. A method of producing a protein comprising:  
culturing a cell into which has been introduced an expression vector according to claim 36, whereby said cell expresses the polypeptide encoded by the DNA segment; and  
recovering the expressed protein.

43. An antibody that specifically binds to an epitope of a polypeptide according to claim 4.

44. The antibody of claim 43 which is a monoclonal antibody.

45. The antibody of claim 43 which is a single-chain antibody.

46. The antibody of claim 43 operably linked to a reporter molecule.

47. A method for detecting a genetic abnormality in a patient, comprising:  
obtaining a genetic sample from a patient;  
incubating the genetic sample with a polynucleotide comprising at least 14 contiguous nucleotides of SEQ ID NO:1 or the complement of SEQ ID NO:1, under conditions wherein said polynucleotide will hybridize to a complementary polynucleotide sequence, to produce a first reaction product; and

comparing said first reaction product to a control reaction product, wherein a difference between said first reaction product and said control reaction product is indicative of a genetic abnormality in the patient.

48. A polypeptide comprising a sequence selected from the group consisting of:

residues 46-234 of SEQ ID NO:33 operably linked to residues 258-370 of SEQ ID NO:2;

residues 46-170 of SEQ ID NO:33 operably linked to residues 180-370 of SEQ ID NO:2;

residues 52-257 of SEQ ID NO:2 operably linked to residues 235-345 of SEQ ID NO:33; and

residues 52-179 of SEQ ID NO:2 operably linked to residues 171-345 of SEQ ID NO:33.

49. A method of activating a cell-surface PDGF receptor, comprising exposing a cell comprising a cell-surface PDGF receptor to the polypeptide or protein of any of claims 1-31, whereby the polypeptide or protein binds to and activates the receptor.

50. The method of claim 49 wherein the receptor is a PDGF alpha-receptor.


51. The method of claim 49 wherein the receptor is a PDGF beta-receptor.

52. A method of inhibiting a PDGF receptor mediated cellular process, comprising exposing a cell comprising a cell-surface PDGF receptor to a compound that inhibits binding of the polypeptide or protein of any of claims 1-31 to the receptor.

53. A method of stimulating the growth of bone tissue, comprising applying to bone a growth-stimulating amount of the polypeptide or protein of any of claims 1-31.

54. A method of modulating the proliferation, differentiation, migration, or metabolism of bone cells, comprising exposing bone cells to an effective amount of the polypeptide or protein of any of claims 1-31.

*1/8*



*Fig. 1*

2/8

49	0.75		L	=====
50	1.32		Y	=====
51	1.32		R	=====
52	1.55		R	=====
53	1.63		D	=====
54	1.17		E	=====
55	0.42		T	=====
56	0.42		I	=====
57	-0.08		= Q	
58	0.02		V	
59	0.32		K	=====
60	-0.10		= G	
61	-0.10		= N	
62	-0.57		===== G	
63	-0.52		===== Y	
64	-0.55		===== V	
65	-0.05		Q	
66	-0.08		= S	
67	0.17		P	=====
68	0.17		R	=====
69	0.17		F	=====
70	0.18		P	=====
71	-0.32		===== N	
72	0.60		S	=====
73	0.63		Y	=====
74	0.30		P	=====
75	-0.05		R	
76	-0.37		===== N	
77	-0.43		===== L	
78	-1.50	=====	L	
79	-1.03	=====	L	
80	-1.03	=====	T	
81	-0.82	=====	W	
82	-0.47	=====	R	
83	-0.37	=====	L	
84	0.70		H	=====
85	0.23		S	=====
86	0.47		Q	=====
87	1.05		E	=====
88	0.70		N	=====
89	0.70		T	=====
90	-0.10		= R	
91	-0.38		===== I	
92	-0.73	=====	Q	
93	-0.73	=====	L	
94	-0.40	=====	V	
95	-0.40	=====	F	
96	-0.52	=====	D	
97	-0.27	=====	N	
98	-0.15		= Q	
99	-0.15		= F	
100	0.32		G	=====
101	0.20		L	=====
102	1.12		E	=====
103	1.15		E	=====

3/8

104	1.95	A	=====
105	1.15	E	=====
106	0.48	N	=====
107	1.07	D	=====
108	0.18	I	==
109	0.65	C	=====
110	-0.27	== R	
111	-0.22	== Y	
112	0.45	D	=====
113	-0.30	== F	
114	0.58	V	=====
115	0.58	E	=====
116	0.70	V	=====
117	1.00	E	=====
118	1.00	D	=====
119	1.18	I	=====
120	0.73	S	=====
121	0.17	E	==
122	0.17	T	==
123	-0.18	== S	
124	-0.18	== T	
125	-0.12	= I	
126	0.33	I	=====
127	-0.17	== R	
128	-0.03	G	
129	0.27	R	=====
130	-0.32	== W	
131	0.18	C	==
132	0.18	G	==
133	0.50	H	=====
134	0.67	K	=====
135	0.67	E	=====
136	1.25	V	=====
137	0.45	P	=====
138	0.45	P	=====
139	0.75	R	=====
140	1.25	I	=====
141	1.18	K	=====
142	0.72	S	=====
143	1.05	R	=====
144	0.25	T	=====
145	0.70	N	=====
146	-0.10	= Q	
147	-0.10	= I	
148	-0.55	==== K	
149	-0.08	= I	
150	0.27	T	=====
151	0.27	F	=====
152	1.07	K	=====
153	0.75	S	=====
154	0.75	D	=====
155	0.00	D	
156	-0.13	= Y	
157	-0.13	= F	
158	-0.63	==== V	

4/8

159	-0.25	===	A
160	-0.25	===	K
161	0.50		P =====
162	0.28		G ===
163	-0.60	=====	F
164	-0.98	=====	K
165	-0.93	=====	I
166	-0.82	=====	Y
167	-1.62	=====	Y
168	-0.82	=====	S
169	0.07		L =
170	0.03		L
171	0.02		E
172	0.32		D ===
173	0.53		F =====
174	-0.05	=	Q
175	-0.63	=====	P
176	-0.17	==	A
177	0.30		A ===
178	0.23		A ==
179	0.35		S =====
180	0.27		E ===
181	0.85		T =====
182	0.85		N =====
183	0.10		w =
184	0.10		E =
185	0.12		S =
186	0.33		V ===
187	-0.47	=====	T
188	-1.08	=====	S
189	-0.83	=====	S
190	-1.02	=====	I
191	-1.02	=====	s
192	-1.45	=====	G
193	-1.12	=====	V
194	-0.50	=====	S
195	-0.50	=====	Y
196	-0.20	==	N
197	-0.50	=====	S
198	-0.18	==	P
199	0.28		S ===
200	0.23		V ==
201	0.17		T ==
202	-0.18	==	D
203	-0.23	==	P
204	-0.25	===	T
205	-0.25	===	L
206	-0.33	===	I
207	-0.57	=====	A
208	0.23		D ==
209	1.03		A =====
210	1.62		L =====
211	0.82		D =====
212	0.82		K =====
213	1.62		K =====

5/8


214	0.70	I	=====
215	0.70	A	=====
216	0.13	E	=
217	0.18	F	==
218	0.77	D	=====
219	0.77	T	=====
220	0.88	V	=====
221	0.08	E	=
222	0.65	D	=====
223	0.52	L	=====
224	-0.40	=====	L
225	-0.87	=====	K
226	-0.57	=====	Y
227	0.23		F ==
228	-0.22	==	N
229	0.00		P
230	0.45		E =====
231	0.92		S =====
232	1.42		w =====
233	0.62		Q =====
234	1.07		E =====
235	1.27		D =====
236	1.02		L =====
237	0.13		E =
238	-0.67	=====	N
239	0.13		M =
240	-0.43	=====	Y
241	-0.47	=====	L
242	0.25		D ===
243	0.65		T =====
244	1.45		P =====
245	0.95		R =====
246	1.52		Y =====
247	1.57		R =====
248	1.08		G =====
249	0.98		R =====
250	0.98		S =====
251	1.48		Y =====
252	1.48		H =====
253	1.48		D =====
254	1.97		R =====
255	1.80		K =====
256	1.80		S =====
257	1.00		K =====
258	1.00		V =====
259	1.45		D =====
260	0.65		L =====
261	0.93		D =====
262	0.93		R =====
263	1.73		L =====
264	1.15		N =====
265	1.15		D =====
266	1.95		D =====
267	1.93		A =====
268	1.48		K =====

6/8

269	0.82		R	=====
270	0.83		Y	=====
271	0.33		S	===
272	0.33		C	===
273	0.35		T	=====
274	0.32		P	===
275	0.53		R	=====
276	0.35		N	=====
277	0.38		Y	=====
278	-0.42	=====	S	
279	0.05		V	=
280	0.53		N	=====
281	0.98		I	=====
282	0.93		R	=====
283	1.40		E	=====
284	1.40		E	=====
285	0.82		L	=====
286	0.35		K	=====
287	-0.40	=====	L	
288	-0.35	=====	A	
289	-1.27	=====	N	
290	-1.38	=====	V	
291	-1.30	=====	V	
292	-0.83	=====	F	
293	-0.75	=====	F	
294	-0.80	=====	P	
295	-0.68	=====	R	
296	-0.52	=====	C	
297	-0.48	=====	L	
298	-0.48	=====	L	
299	-0.48	=====	V	
300	-0.18	==	Q	
301	0.12		R	=
302	0.40		C	=====
303	0.20		G	==
304	-0.30	===	G	
305	-0.30	===	N	
306	-0.87	=====	C	
307	-0.93	=====	G	
308	-1.22	=====	C	
309	-1.02	=====	g	
310	-1.58	=====	T	
311	-0.92	=====	V	
312	-0.30	===	N	
313	-0.40	=====	W	
314	-0.22	==	R	
315	-0.42	=====	S	
316	0.18		C	==
317	-0.27	===	T	
318	-0.32	===	C	
319	0.35		N	=====
320	0.35		S	=====
321	0.27		G	===
322	0.73		K	=====
323	1.18		T	=====



7/8



8/8




Fig. 2

## SEQUENCE LISTING

<110> ZymoGenetics, Inc.

<120> GROWTH FACTOR HOMOLOG ZVEGF4

<130> 99-19PC

<150> US 09/304,216

<151> 1999-05-03

<150> US 60/164,463

<151> 1999-11-10

<150> US 60/180,169

<151> 2000-02-04

<160> 53

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1882

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (226)...(1338)

<400> 1

```
ccgtcaccat ttatcagctc agcaccacaa ggaagtgcgg caccacacacg cgctcggaaa      60
gttcagcatg caggaagttt ggggagagct cggcgattag cacagcgacc cgggccagcg      120
cagggcgagc gcaggcggcg agagcgagg gcggcgcggc gtcggtcccg ggagcagaac      180
ccggcttttt cttggagcga cgctgtctct agtcgctgat cccaa atg cac cgg ctc      237
                                   Met His Arg Leu
```

1

```
atc ttt gtc tac act cta atc tgc gca aac ttt tgc agc tgt cgg gac      285
Ile Phe Val Tyr Thr Leu Ile Cys Ala Asn Phe Cys Ser Cys Arg Asp
  5                      10                      15                      20
```

act tct gca acc ccg cag agc gca tcc atc aaa gct ttg cgc aac gcc	333
Thr Ser Ala Thr Pro Gln Ser Ala Ser Ile Lys Ala Leu Arg Asn Ala	
25 30 35	
aac ctc agg cga gat gag agc aat cac ctc aca gac ttg tac cga aga	381
Asn Leu Arg Arg Asp Glu Ser Asn His Leu Thr Asp Leu Tyr Arg Arg	
40 45 50	
gat gag acc atc cag gtg aaa gga aac ggc tac gtg cag agt cct aga	429
Asp Glu Thr Ile Gln Val Lys Gly Asn Gly Tyr Val Gln Ser Pro Arg	
55 60 65	
ttc ccg aac agc tac ccc agg aac ctg ctc ctg aca tgg cgg ctt cac	477
Phe Pro Asn Ser Tyr Pro Arg Asn Leu Leu Leu Thr Trp Arg Leu His	
70 75 80	
tct cag gag aat aca cgg ata cag cta gtg ttt gac aat cag ttt gga	525
Ser Gln Glu Asn Thr Arg Ile Gln Leu Val Phe Asp Asn Gln Phe Gly	
85 90 95 100	
tta gag gaa gca gaa aat gat atc tgt agg tat gat ttt gtg gaa gtt	573
Leu Glu Glu Ala Glu Asn Asp Ile Cys Arg Tyr Asp Phe Val Glu Val	
105 110 115	
gaa gat ata tcc gaa acc agt acc att att aga gga cga tgg tgt gga	621
Glu Asp Ile Ser Glu Thr Ser Thr Ile Ile Arg Gly Arg Trp Cys Gly	
120 125 130	
cac aag gaa gtt cct cca agg ata aaa tca aga acg aac caa att aaa	669
His Lys Glu Val Pro Pro Arg Ile Lys Ser Arg Thr Asn Gln Ile Lys	
135 140 145	
atc aca ttc aag tcc gat gac tac ttt gtg gct aaa cct gga ttc aag	717
Ile Thr Phe Lys Ser Asp Asp Tyr Phe Val Ala Lys Pro Gly Phe Lys	
150 155 160	
att tat tat tct ttg ctg gaa gat ttc caa ccc gca gca gct tca gag	765
Ile Tyr Tyr Ser Leu Leu Glu Asp Phe Gln Pro Ala Ala Ala Ser Glu	
165 170 175 180	
acc aac tgg gaa tct gtc aca agc tct att tca ggg gta tcc tat aac	813
Thr Asn Trp Glu Ser Val Thr Ser Ser Ile Ser Gly Val Ser Tyr Asn	
185 190 195	

tct cca tca gta acg gat ccc act ctg att gcg gat gct ctg gac aaa	861
Ser Pro Ser Val Thr Asp Pro Thr Leu Ile Ala Asp Ala Leu Asp Lys	
200 205 210	
aaa att gca gaa ttt gat aca gtg gaa gat ctg ctc aag tac ttc aat	909
Lys Ile Ala Glu Phe Asp Thr Val Glu Asp Leu Leu Lys Tyr Phe Asn	
215 220 225	
cca gag tca tgg caa gaa gat ctt gag aat atg tat ctg gac acc cct	957
Pro Glu Ser Trp Gln Glu Asp Leu Glu Asn Met Tyr Leu Asp Thr Pro	
230 235 240	
cgg tat cga ggc agg tca tac cat gac cgg aag tca aaa gtt gac ctg	1005
Arg Tyr Arg Gly Arg Ser Tyr His Asp Arg Lys Ser Lys Val Asp Leu	
245 250 255 260	
gat agg ctc aat gat gat gcc aag cgt tac agt tgc act ccc agg aat	1053
Asp Arg Leu Asn Asp Asp Ala Lys Arg Tyr Ser Cys Thr Pro Arg Asn	
265 270 275	
tac tcg gtc aat ata aga gaa gag ctg aag ttg gcc aat gtg gtc ttc	1101
Tyr Ser Val Asn Ile Arg Glu Glu Leu Lys Leu Ala Asn Val Val Phe	
280 285 290	
ttt cca cgt tgc ctc ctc gtg cag cgc tgt gga gga aat tgt ggc tgt	1149
Phe Pro Arg Cys Leu Leu Val Gln Arg Cys Gly Gly Asn Cys Gly Cys	
295 300 305	
gga act gtc aac tgg agg tcc tgc aca tgc aat tca ggg aaa acc gtg	1197
Gly Thr Val Asn Trp Arg Ser Cys Thr Cys Asn Ser Gly Lys Thr Val	
310 315 320	
aaa aag tat cat gag gta tta cag ttt gag cct ggc cac atc aag agg	1245
Lys Lys Tyr His Glu Val Leu Gln Phe Glu Pro Gly His Ile Lys Arg	
325 330 335 340	
agg ggt aga gct aag acc atg gct cta gtt gac atc cag ttg gat cac	1293
Arg Gly Arg Ala Lys Thr Met Ala Leu Val Asp Ile Gln Leu Asp His	
345 350 355	

cat gaa cga tgc gat tgt atc tgc agc tca aga cca cct cga taa 1338  
 His Glu Arg Cys Asp Cys Ile Cys Ser Ser Arg Pro Pro Arg \*  
                   360                  365                  370

gagaatgtgc acatccttac attaagcctg aaagaacctt tagtttaagg aggggtgagat 1398  
 aagagaccct tttcctacca gcaaccaaac ttactactag cctgcaatgc aatgaacaca 1458  
 agtggttgct gagtctcagc cttgctttgt taatgccatg gcaagtagaa aggtatatca 1518  
 tcaacttcta tacctaagaa tataggattg catttaataa tagtgtttga gggtatatat 1578  
 gcacaaacac acacagaaat atattcatgt ctatgtgtat atagatcaaa tgtttttttt 1638  
 ttttggtata tataaccagg tacaccagag gttacatatg ttgagttag actcttaaaa 1698  
 tcctttgcca aaataaggga tgggtcaaata tatgaaacat gtcttttagaa aatttaggag 1758  
 ataaatttat ttttaaattt tgaaacacga aacaattttg aatcttgctc tcttaaagaa 1818  
 agcatcttgt atattaaaaa tcaaaagatg aggcctttctt acatatacat cttagttgat 1878  
 tatt 1882

<210> 2  
 <211> 370  
 <212> PRT  
 <213> Homo sapiens

<400> 2  
 Met His Arg Leu Ile Phe Val Tyr Thr Leu Ile Cys Ala Asn Phe Cys  
   1                  5                  10                  15  
 Ser Cys Arg Asp Thr Ser Ala Thr Pro Gln Ser Ala Ser Ile Lys Ala  
                   20                  25                  30  
 Leu Arg Asn Ala Asn Leu Arg Arg Asp Glu Ser Asn His Leu Thr Asp  
                   35                  40                  45  
 Leu Tyr Arg Arg Asp Glu Thr Ile Gln Val Lys Gly Asn Gly Tyr Val  
                   50                  55                  60  
 Gln Ser Pro Arg Phe Pro Asn Ser Tyr Pro Arg Asn Leu Leu Leu Thr  
   65                  70                  75                  80  
 Trp Arg Leu His Ser Gln Glu Asn Thr Arg Ile Gln Leu Val Phe Asp  
                   85                  90                  95  
 Asn Gln Phe Gly Leu Glu Glu Ala Glu Asn Asp Ile Cys Arg Tyr Asp  
                   100                  105                  110  
 Phe Val Glu Val Glu Asp Ile Ser Glu Thr Ser Thr Ile Ile Arg Gly  
                   115                  120                  125  
 Arg Trp Cys Gly His Lys Glu Val Pro Pro Arg Ile Lys Ser Arg Thr  
                   130                  135                  140  
 Asn Gln Ile Lys Ile Thr Phe Lys Ser Asp Asp Tyr Phe Val Ala Lys  
   145                  150                  155                  160  
 Pro Gly Phe Lys Ile Tyr Tyr Ser Leu Leu Glu Asp Phe Gln Pro Ala  
                   165                  170                  175

Ala Ala Ser Glu Thr Asn Trp Glu Ser Val Thr Ser Ser Ile Ser Gly  
                   180                  185                  190  
 Val Ser Tyr Asn Ser Pro Ser Val Thr Asp Pro Thr Leu Ile Ala Asp  
                   195                  200                  205  
 Ala Leu Asp Lys Lys Ile Ala Glu Phe Asp Thr Val Glu Asp Leu Leu  
                   210                  215                  220  
 Lys Tyr Phe Asn Pro Glu Ser Trp Gln Glu Asp Leu Glu Asn Met Tyr  
 225                  230                  235                  240  
 Leu Asp Thr Pro Arg Tyr Arg Gly Arg Ser Tyr His Asp Arg Lys Ser  
                   245                  250                  255  
 Lys Val Asp Leu Asp Arg Leu Asn Asp Asp Ala Lys Arg Tyr Ser Cys  
                   260                  265                  270  
 Thr Pro Arg Asn Tyr Ser Val Asn Ile Arg Glu Glu Leu Lys Leu Ala  
                   275                  280                  285  
 Asn Val Val Phe Phe Pro Arg Cys Leu Leu Val Gln Arg Cys Gly Gly  
                   290                  295                  300  
 Asn Cys Gly Cys Gly Thr Val Asn Trp Arg Ser Cys Thr Cys Asn Ser  
 305                  310                  315                  320  
 Gly Lys Thr Val Lys Lys Tyr His Glu Val Leu Gln Phe Glu Pro Gly  
                   325                  330                  335  
 His Ile Lys Arg Arg Gly Arg Ala Lys Thr Met Ala Leu Val Asp Ile  
                   340                  345                  350  
 Gln Leu Asp His His Glu Arg Cys Asp Cys Ile Cys Ser Ser Arg Pro  
                   355                  360                  365  
 Pro Arg  
 370

&lt;210&gt; 3

&lt;211&gt; 126

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; polypeptide motif

&lt;221&gt; VARIANT

&lt;222&gt; (2)...(19)

&lt;223&gt; Xaa = Any Amino Acid

&lt;221&gt; VARIANT

&lt;222&gt; (20)...(34)

&lt;223&gt; Xaa = Any Amino Acid or is not present

<221> VARIANT

<222> (36)...(45)

<223> Xaa = Any Amino Acid

<221> VARIANT

<222> (46)...(72)

<223> Xaa = Any Amino Acid or is not present

<221> VARIANT

<222> (74)...(93)

<223> Xaa = Any Amino Acid

<221> VARIANT

<222> (94)...(123)

<223> Xaa = Any Amino Acid or is not present

<221> VARIANT

<222> (125)...(125)

<223> Xaa = Any Amino Acid

<400> 3

Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1				5					10						15	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				20					25						30	
Xaa	Xaa	Cys	Xaa	Gly	Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
		35					40							45		
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
		50					55						60			
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
65						70				75						80
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				85						90					95	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				100						105					110	
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys	Xaa	Cys			
				115						120					125	

<210> 4

<211> 24

<212> PRT

<213> Artificial Sequence



&lt;220&gt;

&lt;223&gt; polypeptide motif

&lt;221&gt; VARIANT

&lt;222&gt; (2)...(2)

&lt;223&gt; Xaa = Lys or Arg

&lt;221&gt; VARIANT

&lt;222&gt; (4)...(4)

&lt;223&gt; Xaa = Asp, Asn or Glu

&lt;221&gt; VARIANT

&lt;222&gt; (5)...(5)

&lt;223&gt; Xaa = Trp, Tyr or Phe

&lt;221&gt; VARIANT

&lt;222&gt; (6)...(16)

&lt;223&gt; Xaa = Any Amino Acid

&lt;221&gt; VARIANT

&lt;222&gt; (17)...(20)

&lt;223&gt; Xaa = Any Amino Acid or is not present

&lt;221&gt; VARIANT

&lt;222&gt; (22)...(22)

&lt;223&gt; Xaa = Lys or Arg

&lt;221&gt; VARIANT

&lt;222&gt; (23)...(23)

&lt;223&gt; Xaa = Trp, Tyr or Phe

&lt;400&gt; 4

Cys	Xaa	Tyr	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1				5				10							15	
Xaa	Xaa	Xaa	Xaa	Gly	Xaa	Xaa	Cys									
				20												

&lt;210&gt; 5

&lt;211&gt; 6

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> polypeptide tag

<400> 5

Glu Tyr Met Pro Met Glu  
1 5

<210> 6

<211> 1110

<212> DNA

<213> Artificial Sequence

<220>

<223> degenerate sequence

<221> misc\_feature

<222> (1)...(1110)

<223> n = A,T,C or G

<400> 6

atgcaymgny tnathttygt ntayacnytn athtgygcna ayttytgyws ntgymngay	60
acnwsngcna cncncarws ngcnwsnath aargcnytnm gnaaygcnaa yytnmgnmgn	120
gaygarwsna aycayytnac ngayytntay mgnmngayg aracnathca rgtnaarggn	180
aayggntayg tncarwsncc nmgnattyccn aaywsntayc cnmgnaayyt nytnytnacn	240
tggmgnytn aywsncarga raayacnmgn athcarytng tnttygayaa ycarttyggg	300
ytnrgargarg cngaraayga yathtgymgn taygayttyg tngargtnga rgayathwsn	360
garacnwsna cnathathmg nggnmgntgg tgyggncaya argargtncc nccnmgnath	420
aarwsnmgna cnaaycarat haarathacn ttyaarwsng aygaytaytt ygtngcnaar	480
ccnggnttya arathtayta ywsnytnytn gargayttyc arccngcngc ngcnwsngar	540
acnaaytggg arwsngtnac nwsnwsnath wsngngntnw sntayaayws nccnwsngtn	600
acngayccna cnytnathgc ngaygcnytn gayaaraara thgcngartt ygayacngtn	660
gargayytny tnaartaytt yaayccngar wsntggcarg argayytnga raayatgtay	720
ytngayacnc cnmgntaymg nggnmgnwsn taycaygaym gnaarwsnaa rgtngayytn	780
gaymgnytna aygaygaygc naarmgntay wsntgyacnc cnmgnaayta ywsngtnaay	840
athmgngarg arytnaaryt ngcnaaygtg gtnttyttyc cnmgntgyt nytngtncar	900
mgntgyggng gnaaytgygg ntgyggnacn gtnaaytggm gnwsntgyac ntgyaaywsn	960
ggnaaracng tnaaraarta ycaygargtn ytncarttyg arccnggnca yathaarmgn	1020
mgnggnmgng cnaaracnat ggcnytngtg gayathcary tngaycayca ygarmgntgy	1080
gaytgyatht gywsnwsnmg nccnccnmgn	1110

<210> 7

<211> 17

<212> DNA

<213> Artificial Sequence

<220>  
<223> oligonucleotide primer

<221> misc\_feature  
<222> (1)...(17)  
<223> n = A,T,C or G

<400> 7  
mgntgyggng gnaaytg 17

<210> 8  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer

<221> misc\_feature  
<222> (1)...(17)  
<223> n = A,T,C or G

<400> 8  
mgntgydsng gnwrytg 17

<210> 9  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer

<221> misc\_feature  
<222> (1)...(17)  
<223> n = A,T,C or G

<400> 9  
carywnccns hrcanck 17

<210> 10  
<211> 17

10

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide primer

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(17)

&lt;223&gt; n = A,T,C or G

&lt;400&gt; 10

ttyttyccnm gntgyyt

17

&lt;210&gt; 11

&lt;211&gt; 17

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide primer

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(17)

&lt;223&gt; n = A,T,C or G

&lt;400&gt; 11

ntnddnccnn sntgybt

17

&lt;210&gt; 12

&lt;211&gt; 17

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; oligonucleotide primer

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(17)

&lt;223&gt; n = A,T,C or G

&lt;400&gt; 12

avrcansnng gnhhnan

17

<210> 13  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer

<221> misc\_feature  
<222> (1)...(17)  
<223> n = A,T,C or G

<400> 13  
caygarmgnt gygaytg 17

<210> 14  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer

<221> misc\_feature  
<222> (1)...(17)  
<223> n = A,T,C or G

<400> 14  
caynnnnvnt gyvvntg 17

<210> 15  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer

<221> misc\_feature  
<222> (1)...(17)  
<223> n = A,T,C or G

<400> 15

canbbrcanb nnnnrtg	17
<210> 16	
<211> 17	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> oligonucleotide primer	
<221> misc_feature	
<222> (1)...(17)	
<223> n = A,T,C or G	
<400> 16	
tgyacnccnm gnaayta	17
<210> 17	
<211> 17	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> oligonucleotide primer	
<221> misc_feature	
<222> (1)...(17)	
<223> n = A,T,C or G	
<400> 17	
tgyhnnmcnm knrmndh	17
<210> 18	
<211> 17	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> oligonucleotide primer	
<221> misc_feature	
<222> (1)...(17)	
<223> n = A,T,C or G	

<400> 18 dhnkynmkng knndrca	17
<210> 19 <211> 18 <212> DNA <213> Artificial Sequence	
<220> <223> oligonucleotide primer	
<221> misc_feature <222> (1)...(18) <223> n = A,T,C or G	
<400> 19 ntaygaytwy gtngargt	18
<210> 20 <211> 18 <212> DNA <213> Artificial Sequence	
<220> <223> oligonucleotide primer	
<221> misc_feature <222> (1)...(18) <223> n = A,T,C or G	
<400> 20 natrcdrawr canctyca	18
<210> 21 <211> 18 <212> DNA <213> Artificial Sequence	
<220> <223> oligonucleotide primer	
<221> misc_feature	

<222> (1)...(18)  
<223> n = A,T,C or G

<400> 21  
gntdbccnma ndvntayc 18

<210> 22  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer

<221> misc\_feature  
<222> (1)...(18)  
<223> n = A,T,C or G

<400> 22  
cnahvggnkt nhbnatrg 18

<210> 23  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer

<221> misc\_feature  
<222> (1)...(18)  
<223> n = A,T,C or G

<400> 23  
tnhdnggnmr ntdbtgyg 18

<210> 24  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer



<221> misc\_feature  
<222> (1)...(18)  
<223> n = A,T,C or G

<400> 24  
andhnccnky nahvacrc 18

<210> 25  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer ZC21.119

<400> 25  
aggacgatgg tgtggacaca agga 24

<210> 26  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer ZC21.120

<400> 26  
tccagagcat ccgcaatcag agtg 24

<210> 27  
<211> 24  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer ZC21.987

<400> 27  
caacctgttg tttgtcccgt cacc 24

<210> 28  
<211> 25

<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer ZC17.251

<400> 28  
tctggacgtc ctctgctgg tatag 25

<210> 29  
<211> 25  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer ZC17.252

<400> 29  
ggtatggagc aaggggcaag ttggg 25

<210> 30  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer ZC17.156

<400> 30  
gagtggcaac ttccagggcc aggagag 27

<210> 31  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer ZC17.157

<400> 31  
cttttgctag cctcaaccct gactatc 27

<210> 32

<211> 1760

<212> DNA

<213> Homo sapiens

 $\langle 220 \rangle$ 

<221> CDS

<222> (154)...(1191)

<400> 32

attatgtgga aactaccctg cgattctctg ctgccagagc aggctcggcg cttccacccc	60
agtgcagcct tcccctggcg gtggtgaaag agactcggga gtcgctgctt ccaaagtgcc	120
cgccgtgagt gagctctcac cccagtcagc caa atg agc ctc ttc ggg ctt ctc	174
Met Ser Leu Phe Gly Leu Leu	
1 5	
ctg ctg aca tct gcc ctg gcc ggc cag aga cag ggg act cag gcg gaa	222
Leu Leu Thr Ser Ala Leu Ala Gly Gln Arg Gln Gly Thr Gln Ala Glu	
10 15 20	
tcc aac ctg agt agt aaa ttc cag ttt tcc agc aac aag gaa cag aac	270
Ser Asn Leu Ser Ser Lys Phe Gln Phe Ser Ser Asn Lys Glu Gln Asn	
25 30 35	
gga gta caa gat cct cag cat gag aga att att act gtg tct act aat	318
Gly Val Gln Asp Pro Gln His Glu Arg Ile Ile Thr Val Ser Thr Asn	
40 45 50 55	
gga agt att cac agc cca agg ttt cct cat act tat cca aga aat acg	366
Gly Ser Ile His Ser Pro Arg Phe Pro His Thr Tyr Pro Arg Asn Thr	
60 65 70	
gtc ttg gta tgg aga tta gta gca gta gag gaa aat gta tgg ata caa	414
Val Leu Val Trp Arg Leu Val Ala Val Glu Glu Asn Val Trp Ile Gln	
75 80 85	
ctt acg ttt gat gaa aga ttt ggg ctt gaa gac cca gaa gat gac ata	462
Leu Thr Phe Asp Glu Arg Phe Gly Leu Glu Asp Pro Glu Asp Asp Ile	
90 95 100	
tgc aag tat gat ttt gta gaa gtt gag gaa ccc agt gat gga act ata	510
Cys Lys Tyr Asp Phe Val Glu Val Glu Glu Pro Ser Asp Gly Thr Ile	
105 110 115	

tta ggg cgc tgg tgt ggt tct ggt act gta cca gga aaa cag att tct	558
Leu Gly Arg Trp Cys Gly Ser Gly Thr Val Pro Gly Lys Gln Ile Ser	
120 125 130 135	
aaa gga aat caa att agg ata aga ttt gta tct gat gaa tat ttt cct	606
Lys Gly Asn Gln Ile Arg Ile Arg Phe Val Ser Asp Glu Tyr Phe Pro	
140 145 150	
tct gaa cca ggg ttc tgc atc cac tac aac att gtc atg cca caa ttc	654
Ser Glu Pro Gly Phe Cys Ile His Tyr Asn Ile Val Met Pro Gln Phe	
155 160 165	
aca gaa gct gtg agt cct tca gtg cta ccc cct tca gct ttg cca ctg	702
Thr Glu Ala Val Ser Pro Ser Val Leu Pro Pro Ser Ala Leu Pro Leu	
170 175 180	
gac ctg ctt aat aat gct ata act gcc ttt agt acc ttg gaa gac ctt	750
Asp Leu Leu Asn Asn Ala Ile Thr Ala Phe Ser Thr Leu Glu Asp Leu	
185 190 195	
att cga tat ctt gaa cca gag aga tgg cag ttg gac tta gaa gat cta	798
Ile Arg Tyr Leu Glu Pro Glu Arg Trp Gln Leu Asp Leu Glu Asp Leu	
200 205 210 215	
tat agg cca act tgg caa ctt ctt ggc aag gct ttt gtt ttt gga aga	846
Tyr Arg Pro Thr Trp Gln Leu Leu Gly Lys Ala Phe Val Phe Gly Arg	
220 225 230	
aaa tcc aga gtg gtg gat ctg aac ctt cta aca gag gag gta aga tta	894
Lys Ser Arg Val Val Asp Leu Asn Leu Leu Thr Glu Glu Val Arg Leu	
235 240 245	
tac agc tgc aca cct cgt aac ttc tca gtg tcc ata agg gaa gaa cta	942
Tyr Ser Cys Thr Pro Arg Asn Phe Ser Val Ser Ile Arg Glu Glu Leu	
250 255 260	
aag aga acc gat acc att ttc tgg cca ggt tgt ctc ctg gtt aaa cgc	990
Lys Arg Thr Asp Thr Ile Phe Trp Pro Gly Cys Leu Leu Val Lys Arg	
265 270 275	
tgt ggt ggg aac tgt gcc tgt tgt ctc cac aat tgc aat gaa tgt caa	1038
Cys Gly Gly Asn Cys Ala Cys Cys Leu His Asn Cys Asn Glu Cys Gln	
280 285 290 295	

tgt gtc cca agc aaa gtt act aaa aaa tac cac gag gtc ctt cag ttg 1086  
 Cys Val Pro Ser Lys Val Thr Lys Lys Tyr His Glu Val Leu Gln Leu  
                   300                                  305                                  310

aga cca aag acc ggt gtc agg gga ttg cac aaa tca ctc acc gac gtg 1134  
 Arg Pro Lys Thr Gly Val Arg Gly Leu His Lys Ser Leu Thr Asp Val  
                   315                                  320                                  325

gcc ctg gag cac cat gag gag tgt gac tgt gtg tgc aga ggg agc aca 1182  
 Ala Leu Glu His His Glu Glu Cys Asp Cys Val Cys Arg Gly Ser Thr  
                   330                                  335                                  340

gga gga tag ccgcatcacc accagcagct ctgcccaga gctgtgcagt 1231  
 Gly Gly \*  
           345

gcagtggctg attctattag agaacgtatg cggtatctcc atccttaatc tcagttgttt 1291  
 gcttcaagga cctttcatct tcaggattta cagtgcattc tgaaagagga gacatcaaac 1351  
 agaattagga gttgtgcaac agctcttttg agaggaggcc taaaggacag gagaaaagg 1411  
 cttcaatcgt ggaaagaaaa ttaaattgtg tattaaatag atcaccagct agtttcagag 1471  
 ttaccatgta cgtattccac tagctgggtt ctgtatttca gttctttcga tacggcttag 1531  
 ggtaatgtca gtacaggaaa aaaactgtgc aagtgagcac ctgattccgt tgccttgctt 1591  
 aactctaaag ctccatgtcc tgggcctaaa atcgataaaa atctggattt tttttttttt 1651  
 tttttgctca tattcacata tgtaaaccag aacattctat gtactacaaa cctgggtttt 1711  
 aaaaaggaac tatgttgcta tgaattaaac ttgtgtcgtg ctgatagga 1760

<210> 33

<211> 345

<212> PRT

<213> Homo sapiens

<400> 33

Met Ser Leu Phe Gly Leu Leu Leu Thr Ser Ala Leu Ala Gly Gln  
 1                  5                                  10                                  15  
 Arg Gln Gly Thr Gln Ala Glu Ser Asn Leu Ser Ser Lys Phe Gln Phe  
                   20                                  25                                  30  
 Ser Ser Asn Lys Glu Gln Asn Gly Val Gln Asp Pro Gln His Glu Arg  
                   35                                  40                                  45  
 Ile Ile Thr Val Ser Thr Asn Gly Ser Ile His Ser Pro Arg Phe Pro  
                   50                                  55                                  60  
 His Thr Tyr Pro Arg Asn Thr Val Leu Val Trp Arg Leu Val Ala Val  
 65                                  70                                  75                                  80

Glu Glu Asn Val Trp Ile Gln Leu Thr Phe Asp Glu Arg Phe Gly Leu  
                             85                            90                            95  
 Glu Asp Pro Glu Asp Asp Ile Cys Lys Tyr Asp Phe Val Glu Val Glu  
                             100                            105                            110  
 Glu Pro Ser Asp Gly Thr Ile Leu Gly Arg Trp Cys Gly Ser Gly Thr  
                             115                            120                            125  
 Val Pro Gly Lys Gln Ile Ser Lys Gly Asn Gln Ile Arg Ile Arg Phe  
                             130                            135                            140  
 Val Ser Asp Glu Tyr Phe Pro Ser Glu Pro Gly Phe Cys Ile His Tyr  
 145                            150                            155                            160  
 Asn Ile Val Met Pro Gln Phe Thr Glu Ala Val Ser Pro Ser Val Leu  
                             165                            170                            175  
 Pro Pro Ser Ala Leu Pro Leu Asp Leu Leu Asn Asn Ala Ile Thr Ala  
                             180                            185                            190  
 Phe Ser Thr Leu Glu Asp Leu Ile Arg Tyr Leu Glu Pro Glu Arg Trp  
                             195                            200                            205  
 Gln Leu Asp Leu Glu Asp Leu Tyr Arg Pro Thr Trp Gln Leu Leu Gly  
                             210                            215                            220  
 Lys Ala Phe Val Phe Gly Arg Lys Ser Arg Val Val Asp Leu Asn Leu  
 225                            230                            235                            240  
 Leu Thr Glu Glu Val Arg Leu Tyr Ser Cys Thr Pro Arg Asn Phe Ser  
                             245                            250                            255  
 Val Ser Ile Arg Glu Glu Leu Lys Arg Thr Asp Thr Ile Phe Trp Pro  
                             260                            265                            270  
 Gly Cys Leu Leu Val Lys Arg Cys Gly Gly Asn Cys Ala Cys Cys Leu  
                             275                            280                            285  
 His Asn Cys Asn Glu Cys Gln Cys Val Pro Ser Lys Val Thr Lys Lys  
                             290                            295                            300  
 Tyr His Glu Val Leu Gln Leu Arg Pro Lys Thr Gly Val Arg Gly Leu  
 305                            310                            315                            320  
 His Lys Ser Leu Thr Asp Val Ala Leu Glu His His Glu Glu Cys Asp  
                             325                            330                            335  
 Cys Val Cys Arg Gly Ser Thr Gly Gly  
                             340                            345

&lt;210&gt; 34

&lt;211&gt; 3571

&lt;212&gt; DNA

&lt;213&gt; Mus musculus

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1049)...(2086)

&lt;400&gt; 34

gaattcccgg gtcgaccac gcgtccgggc gcccagggga aaggaagctg ggggccgcct	60
ggcggcattc ctcgccgcag tgtgggctcc gtctgccgcg gggcccgcag tgccccctgt	120
ctgcgccagc acctgttggc ccgccagctg gccgccgcg cccccgcgc cccccgcgc	180
cgcccgccg ccagccccgc gcccgcgcg ccgccgcgtg ggggaaagtg gagacgggga	240
ggggacaaga gcgatcctcc aggccagcca ggccttcct tagccgcccg tgcttagccg	300
ccacctctcc tcagccctgc gtcctgccct gccttagggc aggcattccga gcgctcgcga	360
ctccgagccg cccaagctct cccggcttcc cgcagcactt cgccggtacc cgagggaact	420
tcggtggcca ccgactgcag caaggaggag gctccgcggt ggatccgggc cagtcccgag	480
tcgtccccgc ggctctctg cccgcccggg accgcgcg cactcgcagg gcacggtccc	540
ctccccccag gtgggggtgg ggccgcgcct gccgcccga tcagcagctt tgtcattgat	600
cccaaggtgc tcgcctcgt gccgacctg cttccagtct ggcttgccg gaccccgagt	660
cctcgcctgt gtcctgtccc ccaaactgac aggtgctccc tgcgagtcgc cacgactcat	720
cgccgtccc ccggtcccc accccttctt tcctccctcg cctaccccca cccccgcac	780
ttcggcacag ctcaggattt gtttaaact tgggaaact gtccaggtcc aggttttgct	840
ttgatcctt tcaaaaactg gagacacaga agagggtctt aggaaaaact tttggatggg	900
attatgtgga aactaccctg cgattctctg ctgccagagc cggccaggcg cttccaccgc	960
agcgcagcct ttccccggct gggctgagcc ttggagtcgt cgcttcccca gtgcccgcg	1020
cgagtgcgcc ctcgccccag tcagccaa atg ctc ctc ctc ggc ctc ctc ctg	1072
Met Leu Leu Leu Gly Leu Leu Leu	
1 5	

ctg aca tct gcc ctg gcc ggc caa aga acg ggg act cgg gct gag tcc	1120
Leu Thr Ser Ala Leu Ala Gly Gln Arg Thr Gly Thr Arg Ala Glu Ser	
10 15 20	

aac ctg agc agc aag ttg cag ctc tcc agc gac aag gaa cag aac gga	1168
Asn Leu Ser Ser Lys Leu Gln Leu Ser Ser Asp Lys Glu Gln Asn Gly	
25 30 35 40	

gtg caa gat ccc cgg cat gag aga gtt gtc act ata tct ggt aat ggg	1216
Val Gln Asp Pro Arg His Glu Arg Val Val Thr Ile Ser Gly Asn Gly	
45 50 55	

agc atc cac agc ccg aag ttt cct cat aca tac cca aga aat atg gtg	1264
Ser Ile His Ser Pro Lys Phe Pro His Thr Tyr Pro Arg Asn Met Val	
60 65 70	

ctg gtg tgg aga tta gtt gca gta gat gaa aat gtg cgg atc cag ctg	1312
Leu Val Trp Arg Leu Val Ala Val Asp Glu Asn Val Arg Ile Gln Leu	
75 80 85	

aca ttt gat gag aga ttt ggg ctg gaa gat cca gaa gac gat ata tgc	1360
Thr Phe Asp Glu Arg Phe Gly Leu Glu Asp Pro Glu Asp Asp Ile Cys	
90 95 100	
aag tat gat ttt gta gaa gtt gag gag ccc agt gat gga agt gtt tta	1408
Lys Tyr Asp Phe Val Glu Val Glu Glu Pro Ser Asp Gly Ser Val Leu	
105 110 115 120	
gga cgc tgg tgt ggt tct ggg act gtg cca gga aag cag act tct aaa	1456
Gly Arg Trp Cys Gly Ser Gly Thr Val Pro Gly Lys Gln Thr Ser Lys	
125 130 135	
gga aat cat atc agg ata aga ttt gta tct gat gag tat ttt cca tct	1504
Gly Asn His Ile Arg Ile Arg Phe Val Ser Asp Glu Tyr Phe Pro Ser	
140 145 150	
gaa ccc gga ttc tgc atc cac tac agt att atc atg cca caa gtc aca	1552
Glu Pro Gly Phe Cys Ile His Tyr Ser Ile Ile Met Pro Gln Val Thr	
155 160 165	
gaa acc acg agt cct tcg gtg ttg ccc cct tca tct ttg tca ttg gac	1600
Glu Thr Thr Ser Pro Ser Val Leu Pro Pro Ser Ser Leu Ser Leu Asp	
170 175 180	
ctg ctc aac aat gct gtg act gcc ttc agt acc ttg gaa gag ctg att	1648
Leu Leu Asn Asn Ala Val Thr Ala Phe Ser Thr Leu Glu Glu Leu Ile	
185 190 195 200	
cgg tac cta gag cca gat cga tgg cag gtg gac ttg gac agc ctc tac	1696
Arg Tyr Leu Glu Pro Asp Arg Trp Gln Val Asp Leu Asp Ser Leu Tyr	
205 210 215	
aag cca aca tgg cag ctt ttg ggc aag gct ttc ctg tat ggg aaa aaa	1744
Lys Pro Thr Trp Gln Leu Leu Gly Lys Ala Phe Leu Tyr Gly Lys Lys	
220 225 230	
agc aaa gtg gtg aat ctg aat ctc ctc aag gaa gag gta aaa ctc tac	1792
Ser Lys Val Val Asn Leu Asn Leu Leu Lys Glu Glu Val Lys Leu Tyr	
235 240 245	
agc tgc aca ccc cgg aac ttc tca gtg tcc ata cgg gaa gag cta aag	1840
Ser Cys Thr Pro Arg Asn Phe Ser Val Ser Ile Arg Glu Glu Leu Lys	
250 255 260	





```

aagcacgtcc ccagaaacct cgaccatttc taggcacagt gttctgggct atgctgcgct 3276
gtatggacat atcctattta tttcaatact agggttttat tacctttaaa ctctgctcca 3336
tacacttgta ttaatacatg gatattttta tgtacagaag tatatcattt aaggagtcca 3396
cttattatac tctttggcaa ttgcaaagaa aatcaacata atacattgct tgtaaagtct 3456
taatctgtgc ccaagttttg tgggtgactat ttgaattaaa atgtattgaa tcatcaaata 3516
aaataatctg gctatttttg ggaaaaaaaa aaaaaaaaaa aaaaagggcg gccgc 3571

```

&lt;210&gt; 35

&lt;211&gt; 345

&lt;212&gt; PRT

&lt;213&gt; Mus musculus

&lt;400&gt; 35

```

Met Leu Leu Leu Gly Leu Leu Leu Leu Thr Ser Ala Leu Ala Gly Gln
 1             5             10             15
Arg Thr Gly Thr Arg Ala Glu Ser Asn Leu Ser Ser Lys Leu Gln Leu
      20             25             30
Ser Ser Asp Lys Glu Gln Asn Gly Val Gln Asp Pro Arg His Glu Arg
      35             40             45
Val Val Thr Ile Ser Gly Asn Gly Ser Ile His Ser Pro Lys Phe Pro
      50             55             60
His Thr Tyr Pro Arg Asn Met Val Leu Val Trp Arg Leu Val Ala Val
65             70             75             80
Asp Glu Asn Val Arg Ile Gln Leu Thr Phe Asp Glu Arg Phe Gly Leu
      85             90             95
Glu Asp Pro Glu Asp Asp Ile Cys Lys Tyr Asp Phe Val Glu Val Glu
      100            105            110
Glu Pro Ser Asp Gly Ser Val Leu Gly Arg Trp Cys Gly Ser Gly Thr
      115            120            125
Val Pro Gly Lys Gln Thr Ser Lys Gly Asn His Ile Arg Ile Arg Phe
      130            135            140
Val Ser Asp Glu Tyr Phe Pro Ser Glu Pro Gly Phe Cys Ile His Tyr
145            150            155            160
Ser Ile Ile Met Pro Gln Val Thr Glu Thr Thr Ser Pro Ser Val Leu
      165            170            175
Pro Pro Ser Ser Leu Ser Leu Asp Leu Leu Asn Asn Ala Val Thr Ala
      180            185            190
Phe Ser Thr Leu Glu Glu Leu Ile Arg Tyr Leu Glu Pro Asp Arg Trp
      195            200            205
Gln Val Asp Leu Asp Ser Leu Tyr Lys Pro Thr Trp Gln Leu Leu Gly
      210            215            220
Lys Ala Phe Leu Tyr Gly Lys Lys Ser Lys Val Val Asn Leu Asn Leu
225            230            235            240

```

25

Leu Lys Glu Glu Val Lys Leu Tyr Ser Cys Thr Pro Arg Asn Phe Ser  
                             245                            250                            255  
 Val Ser Ile Arg Glu Glu Leu Lys Arg Thr Asp Thr Ile Phe Trp Pro  
                             260                            265                            270  
 Gly Cys Leu Leu Val Lys Arg Cys Gly Gly Asn Cys Ala Cys Cys Leu  
                             275                            280                            285  
 His Asn Cys Asn Glu Cys Gln Cys Val Pro Arg Lys Val Thr Lys Lys  
                             290                            295                            300  
 Tyr His Glu Val Leu Gln Leu Arg Pro Lys Thr Gly Val Lys Gly Leu  
 305                            310                            315                            320  
 His Lys Ser Leu Thr Asp Val Ala Leu Glu His His Glu Glu Cys Asp  
                             325                            330                            335  
 Cys Val Cys Arg Gly Asn Ala Gly Gly  
                             340                            345

<210> 36  
 <211> 600  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> CDS  
 <222> (496)...(592)

<400> 36  
 gtgggaagag tcccggccgg cgattaaact gggcatgctc agtggcagag caggtttagg 60  
 cccggcctgg gaaactgggg agctgagggtg ctgcgcgccgc cgctctgagc ccgagtgcgc 120  
 gcctctcagg ggccgcggcc ggggctggag aacgtgtgtg ctccgctcgc ctgccccgct 180  
 agattcggcg ctgcccgcgc cctgcagcct gtgctgcagc tgccggccac cggagggggc 240  
 gaacaaacaa acgtcaacct gttgtttgtc ccgtcaccat ttatcagctc agcaccacaa 300  
 ggaagtgcgg caccacacag cgctcggaag gtgcagcatg caggaagttt ggggagagct 360  
 cggcgattag cacagcgacc cgggccagcg cagggcgagc gcagacggcg agagcgcagg 420  
 gcggcgcggc gtcgggtccg ggagcagaac ccggcttttt cttggagcga cgctgtctct 480  
 agtcgctgat cccaa atg cac cgg ctc atc ttt gtc tac act cta atc tgc 531

Met His Arg Leu Ile Phe Val Tyr Thr Leu Ile Cys  
                             1                            5                            10

gca aac ttt tgc agc tgt cgg gac act tct gca acc ccg cag agc gca 579  
 Ala Asn Phe Cys Ser Cys Arg Asp Thr Ser Ala Thr Pro Gln Ser Ala  
                             15                            20                            25

26

tcc atc aaa gct t gattatc  
Ser Ile Lys Ala  
30

600

<210> 37  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer ZC22.685

<400> 37  
gccgtcacca tttatcag

18

<210> 38  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> oligonucleotide primer ZC22.686

<400> 38  
cgggtcgctg tgctaac

18

<210> 39  
<211> 18  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> peptide

<400> 39  
Cys Gly His Lys Glu Val Pro Pro Arg Ile Lys Ser Arg Thr Asn Gln  
1 5 10 15  
Ile Lys

<210> 40  
<211> 25

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/40047

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C12N15/62 C12N5/10 C07K14/52 C07K16/22  
C12Q1/68 A61K38/18

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K C12Q A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, STRAND, WPI Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E,L	WO 00 34474 A (ZYMOMETICS INC) 15 June 2000 (2000-06-15) the whole document seq. ID 37 L: priorities ---	1-54
X	DATABASE GENBANK 'Online! Accession No. AA488780, 15 August 1997 (1997-08-15) STRAUSBERG R.: "EST; H. sapiens cDNA clone IMAGE:824754" XP002144644 encodes amino acids 305-370 of seq. ID 2. ---	1-3,22, 31-33, 36-38,42
P,X	WO 99 37671 A (NA SONGQING ;SONG HO YEONG (US); DOU SHENSHEN (US); LILLY CO ELI ( ) 29 July 1999 (1999-07-29) the whole document ---	4-54
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 September 2000

Date of mailing of the international search report

25/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Gall, I

# INTERNATIONAL SEARCH REPORT

In. ational Application No

PCT/US 00/40047

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 98 24811 A (ZYMOGENETICS INC)  11 June 1998 (1998-06-11)  the whole document  -----</p>	1-54

Gln Leu Asp His His Glu Arg Cys Asp Cys Ile Cys Ser Ser Arg Pro  
355 360 365  
Pro Arg  
370

Cys Tyr Pro Asp Thr Phe Ala Thr Pro Gln Arg Ala Ser Ile Lys Ala  
 20 25 30  
 Leu Arg Asn Ala Asn Leu Arg Arg Asp Glu Ser Asn His Leu Thr Asp  
 35 40 45  
 Leu Tyr Gln Arg Glu Glu Asn Ile Gln Val Thr Ser Asn Gly His Val  
 50 55 60  
 Gln Ser Pro Arg Phe Pro Asn Ser Tyr Pro Arg Asn Leu Leu Leu Thr  
 65 70 75 80  
 Trp Trp Leu Arg Ser Gln Glu Lys Thr Arg Ile Gln Leu Ser Phe Asp  
 85 90 95  
 His Gln Phe Gly Leu Glu Glu Ala Glu Asn Asp Ile Cys Arg Tyr Asp  
 100 105 110  
 Phe Val Glu Val Glu Glu Val Ser Glu Ser Ser Thr Val Val Arg Gly  
 115 120 125  
 Arg Trp Cys Gly His Lys Glu Ile Pro Pro Arg Ile Thr Ser Arg Thr  
 130 135 140  
 Asn Gln Ile Lys Ile Thr Phe Lys Ser Asp Asp Tyr Phe Val Ala Lys  
 145 150 155 160  
 Pro Gly Phe Lys Ile Tyr Tyr Ser Phe Val Glu Asp Phe Gln Pro Glu  
 165 170 175  
 Ala Ala Ser Glu Thr Asn Trp Glu Ser Val Thr Ser Ser Phe Ser Gly  
 180 185 190  
 Val Ser Tyr His Ser Pro Ser Ile Thr Asp Pro Thr Leu Thr Ala Asp  
 195 200 205  
 Ala Leu Asp Lys Thr Val Ala Glu Phe Asp Thr Val Glu Asp Leu Leu  
 210 215 220  
 Lys His Phe Asn Pro Val Ser Trp Gln Asp Asp Leu Glu Asn Leu Tyr  
 225 230 235 240  
 Leu Asp Thr Pro His Tyr Arg Gly Arg Ser Tyr His Asp Arg Lys Ser  
 245 250 255  
 Lys Val Asp Leu Asp Arg Leu Asn Asp Asp Val Lys Arg Tyr Ser Cys  
 260 265 270  
 Thr Pro Arg Asn His Ser Val Asn Leu Arg Glu Glu Leu Lys Leu Thr  
 275 280 285  
 Asn Ala Val Phe Phe Pro Arg Cys Leu Leu Val Gln Arg Cys Gly Gly  
 290 295 300  
 Asn Cys Gly Cys Gly Thr Val Asn Trp Lys Ser Cys Thr Cys Ser Ser  
 305 310 315 320  
 Gly Lys Thr Val Lys Lys Tyr His Glu Val Leu Lys Phe Glu Pro Gly  
 325 330 335  
 His Phe Lys Arg Arg Gly Lys Ala Lys Asn Met Ala Leu Val Asp Ile  
 340 345 350



aat gat gat gtc aag cgt tac agt tgc act ccc agg aat cac tct gtg 929  
 Asn Asp Asp Val Lys Arg Tyr Ser Cys Thr Pro Arg Asn His Ser Val  
 265 270 275

aac ctc agg gag gag ctg aag ctg acc aat gca gtc ttc ttc cca cga 977  
 Asn Leu Arg Glu Glu Leu Lys Leu Thr Asn Ala Val Phe Phe Pro Arg  
 280 285 290 295

tgc ctc ctc gtg cag cgc tgt ggt ggc aac tgt ggt tgc gga act gtc 1025  
 Cys Leu Leu Val Gln Arg Cys Gly Gly Asn Cys Gly Cys Gly Thr Val  
 300 305 310

aac tgg aag tcc tgc aca tgc agc tca ggg aag aca gtg aag aag tat 1073  
 Asn Trp Lys Ser Cys Thr Cys Ser Ser Gly Lys Thr Val Lys Lys Tyr  
 315 320 325

cat gag gta ttg aag ttt gag cct gga cat ttc aag aga agg ggc aaa 1121  
 His Glu Val Leu Lys Phe Glu Pro Gly His Phe Lys Arg Arg Gly Lys  
 330 335 340

gct aag aat atg gct ctt gtt gat atc cag ctg gat cat cat gag cga 1169  
 Ala Lys Asn Met Ala Leu Val Asp Ile Gln Leu Asp His His Glu Arg  
 345 350 355

tgt gac tgt atc tgc agc tca aga cca cct cga taa aacactatgc 1215  
 Cys Asp Cys Ile Cys Ser Ser Arg Pro Pro Arg \*  
 360 365 370

acatctgtac ttgattatg aaaggacctt taggttacaa aaaccctaag aagcttctaa 1275  
 tctcagtgc atgaatgcat atggaaatgt tgctttgtta gtgccatggc aagaagaagc 1335  
 aaatatcatt aatttctata tacataaaca taggaattca cttatcaata gtatgtgaag 1395  
 atatgtatat atacttatat acatgactag ctctatgtat gtaaatagat taaatacttt 1455  
 attcagtata ttactg 1472

&lt;210&gt; 53

&lt;211&gt; 370

&lt;212&gt; PRT

&lt;213&gt; Mus musculus

&lt;400&gt; 53

Met Gln Arg Leu Val Leu Val Ser Ile Leu Leu Cys Ala Asn Phe Ser  
 1 5 10 15

aaa aca cgg ata caa ctg tcc ttt gac cat caa ttc gga cta gag gaa Lys Thr Arg Ile Gln Leu Ser Phe Asp His Gln Phe Gly Leu Glu Glu 90 95 100	401
gca gaa aat gac att tgt agg tat gac ttt gtg gaa gtt gaa gaa gtc Ala Glu Asn Asp Ile Cys Arg Tyr Asp Phe Val Glu Val Glu Glu Val 105 110 115	449
tca gag agc agc act gtt gtc aga gga aga tgg tgt ggc cac aag gag Ser Glu Ser Ser Thr Val Val Arg Gly Arg Trp Cys Gly His Lys Glu 120 125 130 135	497
atc cct cca agg ata acg tca aga aca aac cag att aaa atc aca ttt Ile Pro Pro Arg Ile Thr Ser Arg Thr Asn Gln Ile Lys Ile Thr Phe 140 145 150	545
aag tct gat gac tac ttt gtg gca aaa cct gga ttc aag att tat tat Lys Ser Asp Asp Tyr Phe Val Ala Lys Pro Gly Phe Lys Ile Tyr Tyr 155 160 165	593
tca ttt gtg gaa gat ttc caa ccg gaa gca gcc tca gag acc aac tgg Ser Phe Val Glu Asp Phe Gln Pro Glu Ala Ala Ser Glu Thr Asn Trp 170 175 180	641
gaa tca gtc aca agc tct ttc tct ggg gtg tcc tat cac tct cca tca Glu Ser Val Thr Ser Ser Phe Ser Gly Val Ser Tyr His Ser Pro Ser 185 190 195	689
ata acg gac ccc act ctc act gct gat gcc ctg gac aaa act gtc gca Ile Thr Asp Pro Thr Leu Thr Ala Asp Ala Leu Asp Lys Thr Val Ala 200 205 210 215	737
gaa ttc gat acc gtg gaa gat cta ctt aag cac ttc aat cca gtg tct Glu Phe Asp Thr Val Glu Asp Leu Leu Lys His Phe Asn Pro Val Ser 220 225 230	785
tgg caa gat gat ctg gag aat ttg tat ctg gac acc cct cat tat aga Trp Gln Asp Asp Leu Glu Asn Leu Tyr Leu Asp Thr Pro His Tyr Arg 235 240 245	833
ggc agg tca tac cat gat cgg aag tcc aaa gtg gac ctg gac agg ctc Gly Arg Ser Tyr His Asp Arg Lys Ser Lys Val Asp Leu Asp Arg Leu 250 255 260	881

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/40047

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0034474	A	15-06-2000	NONE	
WO 9937671	A	29-07-1999	AU 2562199 A EP 0984063 A	09-08-1999 08-03-2000
WO 9824811	A	11-06-1998	AU 5439998 A	29-06-1998

Form PCT/ISA/210 (patent family annex) (July 1992)



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
9 November 2000 (09.11.2000)

PCT

(10) International Publication Number  
**WO 00/66736 A1**

(51) International Patent Classification<sup>7</sup>: C12N 15/12,  
15/62, 5/10, C07K 14/52, 16/22, C12Q 1/68, A61K 38/18

(21) International Application Number: PCT/US00/40047

(22) International Filing Date: 3 May 2000 (03.05.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/304,216 3 May 1999 (03.05.1999) US  
60/164,463 10 November 1999 (10.11.1999) US  
60/180,169 4 February 2000 (04.02.2000) US

(71) Applicant: ZYMOGENETICS, INC. [US/US]; 1201  
Eastlake Avenue East, Seattle, WA 98102 (US).

(72) Inventors: GILBERT, Teresa; 246 12th Street Southeast,  
Seattle, WA 98002 (US). HART, Charles, E.; 20114 183rd  
Place Northeast, Woodinville, WA 98072 (US). SHEP-  
PARD, Paul, O.; 13532 278th Drive Northeast, Granite  
Falls, WA 98252 (US). GILBERTSON, Debra, G.; 4025  
Northeast 195th Place, Seattle, WA 98155 (US).

(74) Agent: PARKER, Gary, E.; ZymoGenetics, Inc., 1201  
Eastlake Avenue East, Seattle, WA 98102 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,  
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,  
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,  
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ,  
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,  
TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent  
(AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent  
(AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU,  
MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM,  
GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- With international search report.
- With amended claims and statement.

**Date of publication of the amended claims and statement:**  
21 December 2000

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*

(54) Title: GROWTH FACTOR HOMOLOG ZVEGF4

(57) Abstract: Polypeptide growth factors, methods of making them, polynucleotides encoding them, antibodies to them, and meth-  
ods of using them are disclosed. The polypeptides comprise an amino acid segment that is at least 70 % identical to residues 52-179  
of SEQ ID NO:2 or residues 258-370 of SEQ ID NO:2. Multimers of the polypeptides are also disclosed. The polypeptides, mul-  
timeric proteins, and polynucleotides can be used in the study and regulation of cell and tissue development, as components of cell  
culture media, and as diagnostic agents.

WO 00/66736 A1

## AMENDED CLAIMS

[received by the International Bureau on 25 October 2000 (25.10.00);  
original claims 1-54 replaced by amended claims 1-33 (5 pages)]

1. An isolated polypeptide comprising at least 15 contiguous amino acid residues of SEQ ID NO:2.
2. The isolated polypeptide of claim 1 comprising 30 contiguous amino acid residues of SEQ ID NO:2.
3. The isolated polypeptide of claim 1 or claim 2 wherein said polypeptide comprises:
  - residues 19-179 of SEQ ID NO:2;
  - residues 52-179 of SEQ ID NO:2;
  - residues 19-253 of SEQ ID NO:2;
  - residues 52-253 of SEQ ID NO:2;
  - residues 19-255 of SEQ ID NO:2;
  - residues 52-255 of SEQ ID NO:2;
  - residues 19-257 of SEQ ID NO:2;
  - residues 52-257 of SEQ ID NO:2;
  - residues 19-253 of SEQ ID NO:2;
  - residues 52-253 of SEQ ID NO:2;
  - residues 19-370 of SEQ ID NO:2;
  - residues 52-370 of SEQ ID NO:2;
  - residues 180-370 of SEQ ID NO:2;
  - residues 246-370 of SEQ ID NO:2;
  - residues 250-370 of SEQ ID NO:2; or
  - residues 258-370 of SEQ ID NO:2.
4. The isolated polypeptide of any of claims 1-3, further comprising an affinity tag.
5. The isolated polypeptide of any of claims 1-3, further comprising an immunoglobulin constant domain.
6. An isolated protein comprising a first polypeptide operably linked to a second polypeptide, wherein said first polypeptide comprises residues 258-

370 of SEQ ID NO:2, and wherein said protein modulates cell proliferation, apoptosis, differentiation, metabolism, or migration.

7. The isolated protein of claim 6 wherein said first polypeptide comprises:

residues 19-370 of SEQ ID NO:2;  
residues 52-370 of SEQ ID NO:2;  
residues 180-370 of SEQ ID NO:2;  
residues 246-370 of SEQ ID NO:2; or  
residues 250-370 of SEQ ID NO:2.

8. The isolated protein of claim 6 wherein said first polypeptide is from 113 to 120 amino acid residues in length.

9. The isolated protein of any of claims 6-8 wherein said protein is a heterodimer.

10. The isolated protein of claim 9 wherein said second polypeptide is selected from the group consisting of VEGF, VEGF-B, VEGF-C, VEGF-D, zveg3, PlGF, PDGF-A, and PDGF-B.

11. The isolated protein of any of claims 6-8 wherein said protein is a homodimer.

12. An isolated protein produced by a method comprising:  
(a) culturing a host cell containing an expression vector comprising the following operably linked elements:

a transcription promoter;

a DNA segment encoding a polypeptide comprising a sequence of amino acid residues selected from the group consisting of:

- i) residues 52-370 of SEQ ID NO:2;
- ii) residues 52-253 of SEQ ID NO:2;
- iii) residues 180-370 of SEQ ID NO:2; and
- iii) residues 258-370 of SEQ ID NO:2; and

a transcription terminator,

under conditions whereby the DNA segment is expressed; and

(b) recovering from the cell the protein product of expression of the DNA construct.

13. An isolated polynucleotide of up to approximately 4.4 kb in length, wherein said polynucleotide encodes a polypeptide comprising residues 258-370 of SEQ ID NO:2.

14. The polynucleotide of claim 13, wherein said polynucleotide is DNA.

15. The polynucleotide of claim 14 comprising nucleotides 1 through 1110 of SEQ ID NO:6.

16. The polynucleotide of claim 14 comprising nucleotides 226 through 1335 of SEQ ID NO:1.

17. An expression vector comprising the following operably linked elements:

- a transcription promoter;
- a DNA polynucleotide according to claim 14; and
- a transcription terminator.

18. The expression vector of claim 17, further comprising a secretory signal sequence operably linked to the DNA polynucleotide.

19. A cultured cell into which has been introduced an expression vector according to claim 17 or claim 18, wherein said cell expresses the polypeptide encoded by the DNA polynucleotide.

20. A pharmaceutical composition comprising a protein according to any of claims 6-12 in combination with a pharmaceutically acceptable vehicle.

21. A method of producing a protein comprising:  
culturing a cell into which has been introduced an expression vector according to claim 17 or claim 18, whereby said cell expresses the DNA polynucleotide to produce a protein encoded by the polynucleotide; and  
recovering the protein.



22. An antibody that specifically binds to an epitope of a polypeptide according to claim 12.

23. The antibody of claim 22 which is a monoclonal antibody.

24. The antibody of claim 22 which is a single-chain antibody.

25. The antibody of claim 22 operably linked to a reporter molecule.

26. A method for detecting a genetic abnormality in a patient, comprising:

obtaining a genetic sample from a patient;

incubating the genetic sample with a polynucleotide comprising at least 14 contiguous nucleotides of SEQ ID NO:1 or the complement of SEQ ID NO:1, under conditions wherein said polynucleotide will hybridize to a complementary polynucleotide sequence, to produce a first reaction product; and

comparing said first reaction product to a control reaction product, wherein a difference between said first reaction product and said control reaction product is indicative of a genetic abnormality in the patient.

27. A polypeptide comprising a sequence selected from the group consisting of:

residues 46-234 of SEQ ID NO:33 operably linked to residues 258-370 of SEQ ID NO:2;

residues 46-170 of SEQ ID NO:33 operably linked to residues 180-370 of SEQ ID NO:2;

residues 52-257 of SEQ ID NO:2 operably linked to residues 235-345 of SEQ ID NO:33; and

residues 52-179 of SEQ ID NO:2 operably linked to residues 171-345 of SEQ ID NO:33.

28. A method of activating a cell-surface PDGF receptor, comprising exposing a cell comprising a cell-surface PDGF receptor to the polypeptide or protein of any of claims 1-12, whereby the polypeptide or protein binds to and activates the receptor.

29. The method of claim 28 wherein the receptor is a PDGF alpha-receptor.

30. The method of claim 28 wherein the receptor is a PDGF beta-receptor.

31. A method of inhibiting a PDGF receptor-mediated cellular process, comprising exposing a cell comprising a cell-surface PDGF receptor to a compound that inhibits binding of the polypeptide or protein of any of claims 1-12 to the receptor.

32. A method of stimulating the growth of bone tissue, comprising applying to bone a growth-stimulating amount of the polypeptide or protein of any of claims 1-12.

33. A method of modulating the proliferation, differentiation, migration, or metabolism of bone cells, comprising exposing bone cells to an effective amount of the polypeptide or protein of any of claims 1-12.

**STATEMENT UNDER ARTICLE 19**

In response to the Notification of Transmittal of the International Search Report or the Declaration mailed 25 September 2000, please enter the enclosed amendments to the claims. Please replace pages 90 to 96 of the application as filed with replacement pages 90 to 94 enclosed herewith.

Claims 1, 3, 20 to 22, 23, 24, 26, 31 to 39, and 42 to 54 have been replaced by amended claims 1, 3, 4 to 6, 9, 10, 11, 12 to 20, and 21 to 33, respectively; claims 2, 4 to 19, 25, 27-30, 40, and 41 have been cancelled; and new claims 2, 7, and 8 have been added.



<220>

<223> oligonucleotide primer ZC27116

<400> 47

attataggat ccgagtatat gcctatggag gttgacctgg ataggctcaa tgatgatgcc 60

<210> 48

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide primer ZC26137

<400> 48

attatatggc gcgccttatc gaggtggtct tgagct 36

<210> 49

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide primer ZC26317

<400> 49

atcacctcac agacttgtag cagag 25

<210> 50

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide primer ZC26318

<400> 50

cctacaaatg tcattttctg cttcc 25

<210> 51

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide primer ZC14063

<400> 51

caccagacat aatagctgac agact

25

<210> 52

<211> 1472

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (93)...(1205)

<400> 52

agggactgtg cagtagaaat cgcgcgactc aaccctttgg gctttattta tttacttttg 60  
gagcaacgcg atccctaggt cgctgagccc aa atg caa cgg ctc gtt tta gtc 113  
Met Gln Arg Leu Val Leu Val  
1 5

tcc att ctc ctg tgc gcg aac ttt agc tgc tat ccg gac act ttt gcg 161  
Ser Ile Leu Leu Cys Ala Asn Phe Ser Cys Tyr Pro Asp Thr Phe Ala  
10 15 20

act ccg cag aga gca tcc atc aaa gct ttg cgc aat gcc aac ctc agg 209  
Thr Pro Gln Arg Ala Ser Ile Lys Ala Leu Arg Asn Ala Asn Leu Arg  
25 30 35

aga gat gag agc aat cac ctc aca gac ttg tac cag aga gag gag aac 257  
Arg Asp Glu Ser Asn His Leu Thr Asp Leu Tyr Gln Arg Glu Glu Asn  
40 45 50 55

att cag gtg aca agc aat ggc cat gtg cag agt cct cgc ttc ccg aac 305  
Ile Gln Val Thr Ser Asn Gly His Val Gln Ser Pro Arg Phe Pro Asn  
60 65 70

agc tac cca agg aac ctg ctt ctg aca tgg tgg ctc cgt tcc cag gag 353  
Ser Tyr Pro Arg Asn Leu Leu Leu Thr Trp Trp Leu Arg Ser Gln Glu  
75 80 85